

Advanced Calculus

Tougaloo College

Miraj Samarakkody

Basic Topology

Finite, Countable, and Uncountable Sets

Definition 1.

For any positive integer n , let J_n be the set whose elements are the integers $1, 2, 3, \dots, n$.

Let J be the set consisting of all positive integers.

For any set A , we say,

1. A is finite, if $A \sim J_n$.
2. A is infinite if A is not finite. (or A is infinite if A is equivalent to one of its proper subsets.)
3. A is countable if $A \sim J$.
4. A is uncountable if A is neither finite nor countable.
5. A is at most countable if A is finite or countable.

Finite, Countable, and Uncountable Sets (ii)

Example: Let A be the set of all integers (of course not finite). Prove that A is countable.

Finite, Countable, and Uncountable Sets (iii)

Definition 2: Sequence.

A sequence is a function f defined on the set of all positive integers. If $f(n) = x_n$, for $n \in \mathbb{Z}^+$ denote the sequence f by the symbol $\{x_n\}$. The values of f (the elements x_n), are called the **terms** of the sequence.

If A is a set and if $x_n \in A$ for all $n \in \mathbb{Z}^+$, then $\{x_n\}$ is said to be a sequence in A .

Examples:

$$\{2, 4, 6, 8, \dots\}$$

$$\{1, 1, 1, \dots\}$$

Finite, Countable, and Uncountable Sets (iv)

Theorem 1.

Every infinite subset of a countable set A is countable.

Definition 3.

Let A and Ω be sets, and suppose that with each element α of A there is associated a subset E_α of Ω .

The set whose elements are the sets E_α is denoted by $\{E_\alpha\}$.

This is set of sets.

The union of the sets E_α is defined to be the set S such that $x \in S$ if and only if $x \in E_\alpha$ for at least on $\alpha \in A$.

$$S = \cup_{\alpha \in A} E_\alpha$$

If A consists of the integers $1, 2, \dots, n$

$$S = \cup_{m=1}^n E_m = E_1 \cup E_2 \cup E_n$$

Finite, Countable, and Uncountable Sets (v)

If A is the set of all positive integers,

$$S = \bigcup_{m=1}^{\infty} E_m$$

The intersection of the sets E_{α} is the set P such that $x \in P$ if and only if $x \in E_{\alpha}$ for every $\alpha \in A$.

$$P = \bigcap_{\alpha \in A} E_{\alpha}$$

or

$$P = \bigcap_{m=1}^n E_m = E_1 \cap E_2 \cap \dots \cap E_m$$

or

$$P = \bigcap_{m=1}^{\infty} E_m$$

If $A \cap B$ is empty, we say that A and B are disjoint.

Finite, Countable, and Uncountable Sets (vi)

Example:

Suppose

$$E_1 = \{1, 2, 3\}$$

and

$$E_2 = \{2, 3, 4\}$$

1. What is $E_1 \cup E_2$?
2. What is $E_1 \cap E_2$?

Finite, Countable, and Uncountable Sets (vii)

Example:

Let A be the set of real numbers x such that $0 < x \leq 1$. For every $x \in A$, let E_x be the set of real numbers y such that $0 < y < x$. Prove that

1. $E_x \subset E_z$ if and only if $0 < x \leq z \leq 1$.
2. $\bigcup_{x \in A} E_x = E_1$
3. $\bigcap_{x \in A} E_x$ is empty.