3. The Classical Runge–Kutta Method (RK4)

Miraj Samarakkody

2026-10-14

Table of contents

3.1 Derivation from Simpson's rule . . . 2
Worked example: RK4 by hand, first step with $h = 0.1$ . . . 2
3.2 Seeing the slopes . . . 3
3.3 Why the order is what it is: matching the Taylor series . . . 4
3.4 All three methods side by side . . . 4
3.5 Butcher tableaux . . . 5
3.6 Order conditions . . . 7
3.7 Checking the orders numerically . . . 8
3.8 Large steps on a smooth problem . . . 8
3.9 Summary . . . 9
Exercises . . . 9

RK4 uses four slopes per step and has order 4. For smooth, non-stiff problems it gives an excellent balance of accuracy and simplicity, which is why it is the method most people mean by "Runge–Kutta". This section derives it, introduces the **Butcher tableau** notation that describes every Runge–Kutta method, and explains the **order conditions** that determine a method's order.

Test problem: $y' = x + y$, $y(0) = 1$, exact solution $y = 2e^x - x - 1$.

```
import numpy as np
import matplotlib.pyplot as plt

plt.rcParams["figure.figsize"] = (7, 4.5)
plt.rcParams["axes.grid"] = True
np.set_printoptions(precision=6, suppress=True)

f = lambda x, y: x + y
exact = lambda x: 2 * np.exp(x) - x - 1

def euler(f, x0, y0, h, n):
    x, y = np.zeros(n + 1), np.zeros(n + 1)
    x[0], y[0] = x0, y0
    for i in range(n):
        y[i + 1] = y[i] + h * f(x[i], y[i])
        x[i + 1] = x[i] + h
    return x, y

def heun(f, x0, y0, h, n):
    x, y = np.zeros(n + 1), np.zeros(n + 1)
    x[0], y[0] = x0, y0
    for i in range(n):
```

```
        k1 = f(x[i], y[i])
        k2 = f(x[i] + h, y[i] + h * k1)
        y[i + 1] = y[i] + h / 2 * (k1 + k2)
        x[i + 1] = x[i] + h
    return x, y
```

3.1 Derivation from Simpson's rule

Simpson's rule weights the midpoint four times as heavily as the ends:

$$\int_{x_n}^{x_{n+1}} f\,dx \approx \frac{h}{6}\Big[f_{\text{start}} + 4f_{\text{mid}} + f_{\text{end}}\Big].$$

RK4 estimates the midpoint slope **twice**, each time more accurately than the last, and splits the weight 4 into $2+2$:

$$\begin{aligned}
k_1 &= f(x_n,\ y_n) && \text{slope at the start}\\
k_2 &= f\big(x_n + \tfrac{h}{2},\ y_n + \tfrac{h}{2}k_1\big) && \text{slope at the midpoint, reached using } k_1\\
k_3 &= f\big(x_n + \tfrac{h}{2},\ y_n + \tfrac{h}{2}k_2\big) && \text{slope at the midpoint, reached using the better } k_2\\
k_4 &= f(x_n + h,\ y_n + hk_3) && \text{slope at the end, reached using } k_3\\
y_{n+1} &= y_n + \tfrac{h}{6}(k_1 + 2k_2 + 2k_3 + k_4)
\end{aligned}$$

The weights $\frac{1}{6}, \frac{2}{6}, \frac{2}{6}, \frac{1}{6}$ add up to 1, so $\frac{1}{6}(k_1 + 2k_2 + 2k_3 + k_4)$ is a weighted *average slope*. The local error is $O(h^5)$ and the global error $O(h^4)$: RK4 is **order 4**, and halving h divides the error by about 16. Each step costs **4** evaluations of f.

When f depends on x only, $y' = f(x)$, the stages k_2 and k_3 coincide and RK4 **is** Simpson's rule.

```
def rk4(f, x0, y0, h, n):
    x, y = np.zeros(n + 1), np.zeros(n + 1)
    x[0], y[0] = x0, y0
    for i in range(n):
        k1 = f(x[i], y[i])
        k2 = f(x[i] + h / 2, y[i] + h / 2 * k1)
        k3 = f(x[i] + h / 2, y[i] + h / 2 * k2)
        k4 = f(x[i] + h, y[i] + h * k3)
        y[i + 1] = y[i] + h / 6 * (k1 + 2 * k2 + 2 * k3 + k4)
        x[i + 1] = x[i] + h
    return x, y
```

Worked example: RK4 by hand, first step with $h = 0.1$

stage	evaluation point (x, y)	slope
k_1	$(0,\ 1)$	1
k_2	$(0.05,\ 1 + 0.05 \cdot 1) = (0.05,\ 1.05)$	1.1
k_3	$(0.05,\ 1 + 0.05 \cdot 1.1) = (0.05,\ 1.055)$	1.105
k_4	$(0.1,\ 1 + 0.1 \cdot 1.105) = (0.1,\ 1.1105)$	1.2105

$$y_1 = 1 + \frac{0.1}{6}(1 + 2(1.1) + 2(1.105) + 1.2105) = 1 + \frac{0.1}{6}(6.6205) = 1.1103417.$$

Exact: $y(0.1) = 1.1103418$. The error is 1.7×10^{-7}, about 60,000 times smaller than Euler's first step.

3.2 Seeing the slopes

One large step ($h = 0.5$) from $(0, 1)$ shows where each method samples the slope. Short arrows mark each sampled slope; the dashed line is the step actually taken.

```
h, x0, y0 = 0.5, 0.0, 1.0
xf = np.linspace(0, 0.55, 100)

def slope_mark(ax, x, y, s, color, label=None):
    d = 0.06
    ax.plot([x - d, x + d], [y - d * s, y + d * s], color=color, lw=3, label=label)
    ax.plot(x, y, "o", color=color, ms=5)

fig, axes = plt.subplots(1, 3, figsize=(14, 4.2), sharey=True)
for ax in axes:
    ax.plot(xf, exact(xf), "k", lw=2, label="exact")
    ax.set_xlabel("x")
axes[0].set_ylabel("y")


# Euler
k1 = f(x0, y0)
y_e = y0 + h * k1
slope_mark(axes[0], x0, y0, k1, "C0", "k ")
axes[0].plot([x0, h], [y0, y_e], "C0--")
axes[0].set_title(f"Euler: 1 slope\ny  = {y_e:.4f}, error {exact(h) - y_e:.4f}")

# Heun
k2 = f(x0 + h, y0 + h * k1)
y_h = y0 + h / 2 * (k1 + k2)
slope_mark(axes[1], x0, y0, k1, "C0", "k  (start)")
slope_mark(axes[1], x0 + h, y0 + h * k1, k2, "C1", "k  (predicted end)")
axes[1].plot([x0, h], [y0, y_h], "C2--", label="average slope")
axes[1].set_title(f"Heun: 2 slopes\ny  = {y_h:.4f}, error {exact(h) - y_h:.4f}")

# RK4
r1 = f(x0, y0)
r2 = f(x0 + h / 2, y0 + h / 2 * r1)
r3 = f(x0 + h / 2, y0 + h / 2 * r2)
r4 = f(x0 + h, y0 + h * r3)
y_r = y0 + h / 6 * (r1 + 2 * r2 + 2 * r3 + r4)
slope_mark(axes[2], x0, y0, r1, "C0", "k ")
slope_mark(axes[2], x0 + h / 2, y0 + h / 2 * r1, r2, "C1", "k ")
slope_mark(axes[2], x0 + h / 2, y0 + h / 2 * r2, r3, "C4", "k ")
slope_mark(axes[2], x0 + h, y0 + h * r3, r4, "C3", "k ")
axes[2].plot([x0, h], [y0, y_r], "C2--", label="weighted average")
axes[2].set_title(f"RK4: 4 slopes\ny  = {y_r:.4f}, error {exact(h) - y_r:.5f}")

for ax in axes:
    ax.legend(loc="upper left", fontsize=8)
plt.tight_layout()
print(f"exact y(0.5) = {exact(h):.6f}")
```

```
exact y(0.5) = 1.797443
```



3.3 Why the order is what it is: matching the Taylor series

For the test problem, the exact solution after one step from $(0, 1)$ has the Taylor series

$$y(h) = 2e^h - h - 1 = 1 + h + h^2 + \frac{h^3}{3} + \frac{h^4}{12} + \frac{h^5}{60} + \cdots$$

Expanding each method's formula for one step, symbolically in h:

Method	One step gives	First term missed (local error)	Predicted at $h = 0.1$	Actual
Euler	$1 + h$	h^2	1.0×10^{-2}	1.03×10^{-2}
Heun	$1 + h + h^2$	$h^3/3$	3.3×10^{-4}	3.42×10^{-4}
RK4	$1 + h + h^2 + \frac{h^3}{3} + \frac{h^4}{12}$	$h^5/60$	1.7×10^{-7}	1.69×10^{-7}

(For RK4: $k_1 = 1$, $k_2 = 1 + h$, $k_3 = 1 + h + \frac{h^2}{2}$, $k_4 = 1 + 2h + h^2 + \frac{h^3}{2}$; substituting gives the expression in the table.)

This is the real meaning of order: **a method of order p reproduces the Taylor series of the true solution through the h^p term.** The first term it misses, of size h^{p+1}, is the local error. Over $1/h$ steps these add up to a global error of h^p.

3.4 All three methods side by side


```
h, n = 0.1, 10
x, ye = euler(f, 0, 1, h, n)
_, yh = heun(f, 0, 1, h, n)
_, yr = rk4(f, 0, 1, h, n)
yx = exact(x)

print(f"{'x':>5} {'Euler':>10} {'Heun':>10} {'RK4':>12} {'Exact':>12} "
      f"{'|err| Eul':>10} {'|err| Heun':>10} {'|err| RK4':>10}")
for i in range(n + 1):
    print(f"{x[i]:5.1f} {ye[i]:10.6f} {yh[i]:10.6f} {yr[i]:12.8f} {yx[i]:12.8f} "
          f"{abs(ye[i]-yx[i]):10.2e} {abs(yh[i]-yx[i]):10.2e} {abs(yr[i]-yx[i]):10.2e}")

xf = np.linspace(0, 1, 200)
plt.plot(xf, exact(xf), "k", lw=2, label="Exact  2e  - x - 1")
plt.plot(x, ye, "o--", label="Euler")
plt.plot(x, yh, "s--", label="Heun")
```

```
plt.plot(x, yr, "^", ms=8, label="RK4")
plt.title("y' = x + y,  y(0) = 1,  h = 0.1")
plt.xlabel("x"); plt.ylabel("y"); plt.legend();
```

x	Euler	Heun	RK4	Exact	|err| Eul	|err| Heun	|err| RK4
0.0	1.000000	1.000000	1.00000000	1.00000000	0.00e+00	0.00e+00	0.00e+00
0.1	1.100000	1.110000	1.11034167	1.11034184	1.03e-02	3.42e-04	1.69e-07
0.2	1.220000	1.242050	1.24280514	1.24280552	2.28e-02	7.56e-04	3.75e-07
0.3	1.362000	1.398465	1.39971699	1.39971762	3.77e-02	1.25e-03	6.21e-07
0.4	1.528200	1.581804	1.58364848	1.58364940	5.54e-02	1.85e-03	9.15e-07
0.5	1.721020	1.794894	1.79744128	1.79744254	7.64e-02	2.55e-03	1.26e-06
0.6	1.943122	2.040857	2.04423592	2.04423760	1.01e-01	3.38e-03	1.68e-06
0.7	2.197434	2.323147	2.32750325	2.32750541	1.30e-01	4.36e-03	2.16e-06
0.8	2.487178	2.645578	2.65107913	2.65108186	1.64e-01	5.50e-03	2.73e-06
0.9	2.815895	3.012364	3.01920283	3.01920622	2.03e-01	6.84e-03	3.39e-06
1.0	3.187485	3.428162	3.43655949	3.43656366	2.49e-01	8.40e-03	4.17e-06



3.5 Butcher tableaux

Every explicit Runge–Kutta method with s stages has the form

$$k_i = f\Big(x_n + c_i h,\;\; y_n + h\sum_{j=1}^{i-1} a_{ij}\,k_j\Big), \qquad i = 1, \dots, s,$$

$$y_{n+1} = y_n + h\sum_{i=1}^{s} b_i\,k_i,$$

and is completely described by the numbers c_i, a_{ij} and b_i. They are written in a **Butcher tableau**:

$$
\begin{array}{c|ccccc}
c_1 & & & & & \\
c_2 & a_{21} & & & & \\
\vdots & \vdots & \ddots & & & \\
c_s & a_{s1} & \cdots & a_{s,s-1} & & \\
\hline
 & b_1 & \cdots & b_{s-1} & b_s
\end{array}
$$

- "Explicit" means the matrix $A = (a_{ij})$ is strictly lower triangular: stage i uses only stages before it.
- Almost all methods satisfy $c_i = \sum_j a_{ij}$: each stage is evaluated at the x value that its y value corresponds to.
- The b_i are the weights of the final average; they add up to 1.

Euler

$$
\begin{array}{c|c}
0 & \\
\hline
 & 1
\end{array}
$$

Heun

$$
\begin{array}{c|cc}
0 & & \\
1 & 1 & \\
\hline
 & \frac{1}{2} & \frac{1}{2}
\end{array}
$$

Midpoint

$$
\begin{array}{c|cc}
0 & & \\
\frac{1}{2} & \frac{1}{2} & \\
\hline
 & 0 & 1
\end{array}
$$

Kutta's 3rd order

$$
\begin{array}{c|ccc}
0 & & & \\
\frac{1}{2} & \frac{1}{2} & & \\
1 & -1 & 2 & \\
\hline
 & \frac{1}{6} & \frac{2}{3} & \frac{1}{6}
\end{array}
$$

Classical RK4

$$
\begin{array}{c|cccc}
0 & & & & \\
\frac{1}{2} & \frac{1}{2} & & & \\
\frac{1}{2} & 0 & \frac{1}{2} & & \\
1 & 0 & 0 & 1 & \\
\hline
 & \frac{1}{6} & \frac{1}{3} & \frac{1}{3} & \frac{1}{6}
\end{array}
$$

3/8 rule

$$
\begin{array}{c|cccc}
0 & & & & \\
\frac{1}{3} & \frac{1}{3} & & & \\
\frac{2}{3} & -\frac{1}{3} & 1 & & \\
1 & 1 & -1 & 1 & \\
\hline
 & \frac{1}{8} & \frac{3}{8} & \frac{3}{8} & \frac{1}{8}
\end{array}
$$

One function can now run **any** explicit Runge–Kutta method, given its tableau.

```
TABLEAUX = {
    "Euler":   dict(c=[0], A=[[0]], b=[1]),
    "Heun":    dict(c=[0, 1], A=[[0, 0], [1, 0]], b=[1/2, 1/2]),
    "Midpoint": dict(c=[0, 1/2], A=[[0, 0], [1/2, 0]], b=[0, 1]),
    "Ralston": dict(c=[0, 2/3], A=[[0, 0], [2/3, 0]], b=[1/4, 3/4]),
    "Kutta3":  dict(c=[0, 1/2, 1], A=[[0, 0, 0], [1/2, 0, 0], [-1, 2, 0]], b=[1/6, 2/3, 1/6]),
    "RK4":     dict(c=[0, 1/2, 1/2, 1],
                    A=[[0, 0, 0, 0], [1/2, 0, 0, 0], [0, 1/2, 0, 0], [0, 0, 1, 0]],
                    b=[1/6, 1/3, 1/3, 1/6]),
    "3/8 rule": dict(c=[0, 1/3, 2/3, 1],
                     A=[[0, 0, 0, 0], [1/3, 0, 0, 0], [-1/3, 1, 0, 0], [1, -1, 1, 0]],
                     b=[1/8, 3/8, 3/8, 1/8]),
}
TABLEAUX = {name: {k: np.array(v, dtype=float) for k, v in t.items()} for name, t in TABLEAUX.items()}


def explicit_rk(tableau, f, x0, y0, h, n):
    A, b, c = tableau["A"], tableau["b"], tableau["c"]
    s = len(b)
    x, y = np.zeros(n + 1), np.zeros(n + 1)
    x[0], y[0] = x0, y0
    for i in range(n):
        k = np.zeros(s)
        for j in range(s):
            k[j] = f(x[i] + c[j] * h, y[i] + h * (A[j, :j] @ k[:j]))
        y[i + 1] = y[i] + h * (b @ k)
        x[i + 1] = x[i] + h
    return x, y
```

```
print("max difference between explicit_rk(RK4) and rk4:",
      np.max(np.abs(explicit_rk(TABLEAUX["RK4"], f, 0, 1, 0.1, 10)[1] - rk4(f, 0, 1, 0.1, 10)[1])))
```

```
max difference between explicit_rk(RK4) and rk4: 8.881784197001252e-16
```

3.6 Order conditions

Section 2.3 found the conditions for order 2 by matching Taylor series. Doing the same at higher order gives one condition for each distinct term ("elementary differential") in the Taylor series of the exact solution. In tableau notation, with $\mathbf{c} = (c_i)$, $\mathbf{1} = (1, \dots, 1)$ and powers of vectors taken componentwise:

Order	New conditions
1	$\mathbf{b} \cdot \mathbf{1} = 1$
2	$\mathbf{b} \cdot \mathbf{c} = \frac{1}{2}$
3	$\mathbf{b} \cdot \mathbf{c}^2 = \frac{1}{3}$, $\mathbf{b} \cdot A\mathbf{c} = \frac{1}{6}$
4	$\mathbf{b} \cdot \mathbf{c}^3 = \frac{1}{4}$, $\mathbf{b} \cdot (\mathbf{c} * A\mathbf{c}) = \frac{1}{8}$, $\mathbf{b} \cdot A\mathbf{c}^2 = \frac{1}{12}$, $\mathbf{b} \cdot A^2\mathbf{c} = \frac{1}{24}$

(Here $*$ is the componentwise product.) A method has order p if it satisfies every condition up to and including order p.

The code checks every tableau against these conditions.

```
def order_conditions(t):
    A, b, c = t["A"], t["b"], t["c"]
    one = np.ones_like(b)
    return {
        1: [(b @ one, 1)],
        2: [(b @ c, 1/2)],
        3: [(b @ c**2, 1/3), (b @ (A @ c), 1/6)],
        4: [(b @ c**3, 1/4), (b @ (c * (A @ c)), 1/8), (b @ (A @ c**2), 1/12), (b @ (A @ A @ c), 1/24)],
    }

def order_of(t):
    p = 0
    for order, conds in order_conditions(t).items():
        if all(abs(value - target) < 1e-12 for value, target in conds):
            p = order
        else:
            break
    return p

for name, t in TABLEAUX.items():
    print(f"{name:9s} stages = {len(t['b'])}   order (from conditions) = {order_of(t)}")
```

```
Euler     stages = 1   order (from conditions) = 1
Heun      stages = 2   order (from conditions) = 2
Midpoint  stages = 2   order (from conditions) = 2
Ralston   stages = 2   order (from conditions) = 2
Kutta3    stages = 3   order (from conditions) = 3
RK4       stages = 4   order (from conditions) = 4
3/8 rule  stages = 4   order (from conditions) = 4
```

How fast the conditions multiply. The number of order conditions grows quickly with the order:

Order p	1	2	3	4	5	6	7	8
Conditions (total)	1	2	4	8	17	37	85	200
Minimum stages	1	2	3	4	6	7	9	11

The second row is Butcher's **barrier** result: up to order 4 you need exactly p stages, but beyond that the number of stages exceeds the order. Order 5 costs 6 evaluations per step. This is why RK4 is such a sweet spot: it is the highest order you get "at the price of the order".

3.7 Checking the orders numerically

The order conditions say what *should* happen. To confirm it, run each tableau on a **nonlinear** problem (as Section 2.4 showed, a linear test problem can hide differences between methods). Use the logistic equation

$$y' = y(1-y), \qquad y(0) = 0.1, \qquad \text{exact solution } y = \frac{1}{1+9e^{-x}},$$

and estimate the order from errors at $x = 4$ with h and $h/2$: $p \approx \log_2 \left(E(h)/E(h/2)\right)$.

```
logistic = lambda x, y: y * (1 - y)
log_exact = lambda x: 1 / (1 + 9 * np.exp(-x))

print(f"{'method':9s} {'err h=0.2':>11} {'err h=0.1':>11} {'err h=0.05':>11} {'observed order':>15}")
for name, t in TABLEAUX.items():
    e = [abs(log_exact(4) - explicit_rk(t, logistic, 0, 0.1, h, round(4 / h))[1][-1]) for h in (0.2, 0.1
    print(f"{name:9s} {e[0]:11.2e} {e[1]:11.2e} {e[2]:11.2e} {np.log2(e[1] / e[2]):15.2f}")
```

```
method      err h=0.2   err h=0.1  err h=0.05  observed order
Euler        4.22e-03    1.96e-03    9.45e-04            1.05
Heun         1.38e-03    3.46e-04    8.69e-05            1.99
Midpoint     4.59e-04    1.17e-04    2.94e-05            1.99
Ralston      7.64e-04    1.93e-04    4.86e-05            1.99
Kutta3       8.19e-06    1.11e-06    1.44e-07            2.94
RK4          1.54e-06    9.72e-08    6.11e-09            3.99
3/8 rule     1.12e-06    7.12e-08    4.49e-09            3.99
```

The observed orders match the order conditions: about 1 for Euler, 2 for the two-stage methods, 3 for Kutta's method and 4 for both four-stage methods.

3.8 Large steps on a smooth problem

High order pays off most when the solution is smooth and the interval is long. Take


$$y' = y\cos x, \qquad y(0) = 1, \qquad \text{exact solution } y = e^{\sin x}, \qquad 0 \le x \le 20,$$

with a **large** step $h = 0.5$ (only 40 steps over about three periods).

```
w = lambda x, y: y * np.cos(x)
w_exact = lambda x: np.exp(np.sin(x))
h, n = 0.5, 40
xf = np.linspace(0, 20, 600)
plt.figure(figsize=(10, 4.2))
plt.plot(xf, w_exact(xf), "k", lw=2, label="exact  e^(sin x)")
```

```
for name, method, style in [("Euler", euler, "o--"), ("Heun", heun, "s--"), ("RK4", rk4, "^")]:
    x, y = method(w, 0, 1, h, n)
    plt.plot(x, y, style, ms=5, label=name)
    print(f"{name:6s} max |error| on [0, 20] = {np.max(np.abs(y - w_exact(x))):.2e}   ({n * {'Euler': 1,
plt.xlabel("x"); plt.ylabel("y"); plt.title("y' = y cos x with h = 0.5"); plt.legend();
```

```
Euler  max |error| on [0, 20] = 2.32e+00   (40 f evaluations)
Heun   max |error| on [0, 20] = 2.22e-01   (80 f evaluations)
RK4    max |error| on [0, 20] = 2.20e-03   (160 f evaluations)
```



With only 40 steps, RK4 is visually indistinguishable from the exact solution, while Euler's solution has drifted far from it.

3.9 Summary

- RK4: four stages, order 4, Simpson's rule in disguise.
- A Butcher tableau $(A, \mathbf{b}, \mathbf{c})$ describes any Runge–Kutta method; one generic function can run them all.
- A method has order p when its tableau satisfies all order conditions up to order p. Their number grows rapidly, and beyond order 4 more stages than the order are needed.

Exercises

1. Take one RK4 step with $h = 0.1$ for $y' = y$, $y(0) = 1$. Show that the result is $1 + h + \frac{h^2}{2} + \frac{h^3}{6} + \frac{h^4}{24}$ and compare it with $e^{0.1}$.
2. Show by hand that Kutta's third-order method satisfies the four conditions for order 3.
3. Show that for $y' = \lambda y$ one RK4 step multiplies y_n by $R(z) = 1 + z + \frac{z^2}{2} + \frac{z^3}{6} + \frac{z^4}{24}$ with $z = h\lambda$. (This polynomial reappears in Section 5.)
4. The tableau with $c = (0, 1)$, $a_{21} = 1$ and $b = (1, 0)$ is consistent ($\sum b_i = 1$). What is its order, and why is it a poor method?

Answers

1. $k_1 = 1$, $k_2 = 1 + \frac{h}{2}$, $k_3 = 1 + \frac{h}{2} + \frac{h^2}{4}$, $k_4 = 1 + h + \frac{h^2}{2} + \frac{h^3}{4}$. Substituting gives the Taylor polynomial: $1.10517083\overline{3}$ against $e^{0.1} = 1.1051709181$.
2. $\mathbf{b} \cdot \mathbf{1} = \frac{1}{6} + \frac{2}{3} + \frac{1}{6} = 1$; $\mathbf{b} \cdot \mathbf{c} = \frac{2}{3} \cdot \frac{1}{2} + \frac{1}{6} = \frac{1}{2}$; $\mathbf{b} \cdot \mathbf{c}^2 = \frac{2}{3} \cdot \frac{1}{4} + \frac{1}{6} = \frac{1}{3}$; $A\mathbf{c} = (0, 0, 2 \cdot \frac{1}{2}) = (0, 0, 1)$, so $\mathbf{b} \cdot A\mathbf{c} = \frac{1}{6}$.

3. Same calculation as Exercise 1 with λy in place of y.
4. It is Euler's method with a wasted second stage: $\mathbf{b} \cdot \mathbf{c} = 0 \neq \frac{1}{2}$, so order 1, at twice Euler's cost.

```
_, y = rk4(lambda x, y: y, 0, 1, 0.1, 1)
h = 0.1
print(f"1.  RK4 step = {y[1]:.10f},  Taylor polynomial = {1 + h + h**2/2 + h**3/6 + h**4/24:.10f},  e^0.
print(f"2.  Kutta3 order from conditions: {order_of(TABLEAUX['Kutta3'])}")
waste = {k: np.array(v, dtype=float) for k, v in dict(c=[0, 1], A=[[0, 0], [1, 0]], b=[1, 0]).items()}
print(f"4.  order from conditions: {order_of(waste)}")
```

```
1.  RK4 step = 1.1051708333,  Taylor polynomial = 1.1051708333,  e^0.1 = 1.1051709181
2.  Kutta3 order from conditions: 3
4.  order from conditions: 1
```