A2 · Unit 4 · WCH14 Rates, Equilibria and Further Organic Chemistry

Carbonyls, carboxylic acids and chirality

The C=O group is the busiest functional group in organic chemistry. This topic covers aldehydes, ketones, carboxylic acids and the compounds made from them, the mirror-image molecules that some of their reactions produce, and the instrumental methods (NMR, mass spectrometry and chromatography) that chemists use to work out what they have made.

Naming the compounds in this topic

Family	Functional group	Name ending	Example
aldehyde	–CHO (C=O at the end of the chain)	-al	CH_3CH_2CHO, propanal
ketone	C=O within the chain	-one	$CH_3COCH_2CH_3$, butanone; $CH_3COCH_2CH_2CH_3$, pentan-2-one
carboxylic acid	–COOH	-oic acid	CH_3CH_2COOH, propanoic acid; HOOC–COOH, ethanedioic acid
ester	–COO–	alkyl …-oate	$CH_3COOCH_2CH_3$, ethyl ethanoate
acyl chloride	–COCl	-oyl chloride	CH_3COCl, ethanoyl chloride
amide	$–CONH_2$	-amide	CH_3CONH_2, ethanamide; $CH_3CONHCH_3$, N-methylethanamide
nitrile	–C≡N	-nitrile	CH_3CH_2CN, propanenitrile

- **Aldehydes and carboxylic acids** always have their functional carbon at the end of the chain, so it is carbon 1 and needs no number. In a nitrile, the carbon of the C≡N counts as part of the chain: CH_3CH_2CN has three carbons.
- **Esters** are named from two parts: the alkyl group from the alcohol comes first, then the part from the acid (including the C=O carbon) ending in -oate. $CH_3CH_2COOCH_3$ is methyl propanoate.
- **Hydroxynitriles** are named as nitriles, with the C≡N carbon as carbon 1: $CH_3CH(OH)CN$ is 2-hydroxypropanenitrile.

Chirality and optical isomerism

A carbon atom joined to **four different groups** is a **chiral centre** (an asymmetric carbon). The four groups are arranged tetrahedrally, and there are two ways of doing that which are mirror images of each other. No amount of turning makes one fit exactly on top of the other: they are **non-superimposable mirror images**, like your left and right hands. The two forms are **enantiomers**, or optical isomers.

- **Same chemistry:** enantiomers have identical physical properties and react in the same way with most reagents.
- **Different effect on light:** each enantiomer rotates the plane of **plane-polarised light**, by the same angle but in opposite directions. One turns it clockwise, the other anticlockwise.
- **Racemic mixture (racemate):** a 50 : 50 mixture of the two enantiomers. The rotations cancel, so it has no effect on plane-polarised light: it is optically inactive.
- Draw enantiomers in 3D: two bonds in the plane of the paper, one wedge (towards you) and one dashed (away from you), with the second molecule drawn as the mirror image of the first.

How reaction mechanisms decide optical activity

Whether a reaction gives one enantiomer or a racemic mixture depends on the shape of the particle that the attacking species meets.

- **Attack on a planar group gives a racemate.** A trigonal planar C=O group or carbocation can be attacked equally from above and below. If the new carbon is chiral, the two directions of attack give the two enantiomers in equal amounts. This happens in nucleophilic addition of HCN to aldehydes and most ketones, in the S_N1 substitution of a tertiary halogenoalkane, and in electrophilic addition of HBr to an unsymmetrical alkene such as but-1-ene, where the carbocation is planar.
- **S_N2 inverts the chiral centre.** In S_N2 the nucleophile attacks from the side opposite the leaving group in one step, through a transition state with five groups partly bonded to carbon. The other three groups flip over like an umbrella in the wind, so a single enantiomer of the reactant gives a single enantiomer of the product with the opposite arrangement. The product is optically active.
- **S_N1 goes through a planar carbocation**: the leaving group goes first, then the nucleophile can attack either face, so an optically active reactant gives an optically inactive racemic product. Measuring the optical activity of the product is therefore evidence for which mechanism happened.

Aldehydes and ketones

In the C=O group oxygen is much more electronegative than carbon, so the bond is polar, $C^{\delta+}=O^{\delta-}$. The electron-poor carbon is attacked by **nucleophiles**, unlike the C=C in alkenes, which attracts electrophiles. The C=O bond is also stronger than a C=C bond, so carbonyls do not take part in the electrophilic addition reactions of alkenes.

- **Boiling points:** carbonyl molecules cannot hydrogen bond to each other (no H on O), but they have permanent dipole–dipole forces as well as London forces. Their boiling points are higher than alkanes of similar size but lower than alcohols.

- **Solubility:** small carbonyls such as propanone mix with water, because water molecules hydrogen bond to the lone pairs on the carbonyl oxygen. Solubility falls as the hydrocarbon chain gets longer.

Oxidation: telling aldehydes from ketones

An aldehyde has a hydrogen on its carbonyl carbon and is easily oxidised to a carboxylic acid. A ketone has no such hydrogen, and mild oxidising agents leave it alone. Primary alcohols oxidise to aldehydes and then acids, secondary alcohols to ketones, and tertiary alcohols not at all.

Reagent and conditions	With an aldehyde	With a ketone
potassium dichromate(VI) and dilute sulfuric acid, heat under reflux	orange $Cr_2O_7^{2-}$ turns green (Cr^{3+})	stays orange
Fehling's or Benedict's solution (blue Cu^{2+} complex), warm	brick-red precipitate of copper(I) oxide, Cu_2O	stays blue
Tollens' reagent ($[Ag(NH_3)_2]^+$, made from silver nitrate and aqueous ammonia), warm in a water bath	silver mirror forms on the inside of the tube	no change

Full equations, with ethanal being oxidised to ethanoic acid in each case:

- Dichromate: $3CH_3CHO + Cr_2O_7^{2-} + 8H^+ \rightarrow 3CH_3COOH + 2Cr^{3+} + 4H_2O$ (or simply RCHO + [O] → RCOOH).
- Fehling's: $CH_3CHO + 2Cu^{2+} + 2H_2O \rightarrow CH_3COOH + Cu_2O + 4H^+$
- Tollens': $CH_3CHO + 2Ag^+ + H_2O \rightarrow CH_3COOH + 2Ag + 2H^+$

Reduction with lithium tetrahydridoaluminate

$LiAlH_4$ in dry ether, at room temperature, reduces aldehydes to **primary** alcohols and ketones to **secondary** alcohols. Write the reducing agent as [H]:


- $CH_3CH_2CHO + 2[H] \rightarrow CH_3CH_2CH_2OH$ (propanal → propan-1-ol)
- $CH_3COCH_3 + 2[H] \rightarrow CH_3CH(OH)CH_3$ (propanone → propan-2-ol)
- The ether must be dry because $LiAlH_4$ reacts violently with water.

Nucleophilic addition of hydrogen cyanide

HCN adds across the C=O bond to give a **hydroxynitrile**, adding one carbon to the chain. The reagent is HCN with some potassium cyanide, at room temperature; HCN itself barely dissociates, so the KCN supplies enough cyanide ions, the actual nucleophile, to start the reaction.

- $CH_3CHO + HCN \rightarrow CH_3CH(OH)CN$ (2-hydroxypropanenitrile)

- CH_3COCH_3 + HCN → $(CH_3)_2C(OH)CN$ (2-hydroxy-2-methylpropanenitrile)



ethanal — intermediate (alkoxide ion) — 2-hydroxypropanenitrile

Nucleophilic addition. The lone pair on the cyanide ion attacks the δ+ carbon while the C=O π electrons move onto the oxygen. The negative oxygen of the intermediate then takes a proton from HCN, which regenerates a cyanide ion.

KEY IDEA

The carbonyl group is planar, so CN^- can attack from either face. With ethanal, or any unsymmetrical ketone, the product carbon is chiral and the two faces give the two enantiomers equally: the product is a **racemic mixture** with no optical activity.

Two more tests

- **2,4-dinitrophenylhydrazine (2,4-DNPH, Brady's reagent)** gives an orange precipitate with any aldehyde or ketone, but not with other C=O compounds such as acids or esters. Purify the precipitate by recrystallisation and measure its melting point: comparing with tables of known values identifies the exact carbonyl compound.
- **The iodoform test**: warm with iodine and sodium hydroxide. A pale yellow precipitate of triiodomethane (iodoform, CHI_3), with an antiseptic smell, shows a **$CH_3CO–$** group: methyl ketones and ethanal (the only aldehyde that gives it). Alcohols with a $CH_3CH(OH)–$ group also give a positive result, because the alkaline iodine first oxidises them to $CH_3CO–$ compounds.
  $CH_3COCH_3 + 3I_2 + 4NaOH \rightarrow CHI_3 + CH_3COONa + 3NaI + 3H_2O$

Carboxylic acids

- **Hydrogen bonding**: the –COOH group hydrogen bonds strongly, so carboxylic acids have high boiling points. In the pure liquid, and in non-polar solvents, pairs of molecules hydrogen bond into **dimers**, which is why ethanoic acid can appear to have M_r 120 rather than 60.
- **Solubility**: acids with up to four carbons mix completely with water because they hydrogen bond to water molecules; longer chains make them less soluble.
- **Weak acids**: $CH_3COOH \rightleftharpoons CH_3COO^- + H^+$, only slightly dissociated, but still strong enough to release CO_2 from carbonates.

- **Why they are acidic at all**: in the carboxylate ion the negative charge is **delocalised** over both oxygen atoms (the two C–O bonds become the same length). Spreading the charge stabilises the ion, which encourages the O–H bond to break.
- **Effect of substituents**: alkyl groups push electrons towards the COO^- group, concentrating its charge and making the ion less stable, so propanoic acid is slightly weaker than ethanoic acid. Electronegative chlorine atoms pull electron density away, spreading the charge, so chloroethanoic acid is stronger.

Making carboxylic acids

- **Oxidising a primary alcohol or an aldehyde** with excess acidified potassium dichromate(VI), heating under reflux so the aldehyde formed on the way cannot escape: $CH_3CH_2CH_2OH + 2[O] \rightarrow CH_3CH_2COOH + H_2O$.
- **Hydrolysing a nitrile** by heating under reflux with dilute hydrochloric or sulfuric acid: $CH_3CH_2CN + 2H_2O + H^+ \rightarrow CH_3CH_2COOH + NH_4^+$. Together with HCN addition, this lets you lengthen a carbon chain and finish with an acid.

Reactions of carboxylic acids

Reagent	Product	Equation
$LiAlH_4$ in dry ether	primary alcohol	$CH_3CH_2COOH + 4[H] \rightarrow CH_3CH_2CH_2OH + H_2O$
reactive metal	salt + hydrogen	$2CH_3COOH + 2Na \rightarrow 2CH_3COONa + H_2$
alkali	salt + water	$CH_3COOH + NaOH \rightarrow CH_3COONa + H_2O$
carbonate or hydrogencarbonate	salt + water + CO_2	$2CH_3COOH + Na_2CO_3 \rightarrow 2CH_3COONa + H_2O + CO_2$
phosphorus(V) chloride, room temperature	acyl chloride	$CH_3COOH + PCl_5 \rightarrow CH_3COCl + POCl_3 + HCl$
alcohol + concentrated H_2SO_4 catalyst, heat	ester + water	$CH_3COOH + C_2H_5OH \rightleftharpoons CH_3COOC_2H_5 + H_2O$

- **Tests for –COOH**: fizzing with sodium carbonate or sodium hydrogencarbonate solution (CO_2 turns limewater milky), and misty fumes of HCl with PCl_5 (PCl_5 also reacts with alcohols, so it shows an –OH group of either kind).
- **Methanoic acid is the exception** that can be oxidised: HCOOH has a structure like an aldehyde (an H on the C=O carbon). It is oxidised to carbonic acid, which breaks down to CO_2 and water, so it gives a positive result with Tollens' and Fehling's.

- **Esterification** is slow and reversible, giving only a modest equilibrium yield; acyl chlorides (below) make esters much better.

Acyl chlorides

Acyl chlorides are far more reactive than carboxylic acids. The carbonyl carbon carries two electronegative atoms, O and Cl, making it strongly δ+, and Cl^- is a good leaving group. They react with nucleophiles at room temperature, quickly and completely, and every reaction gives off steamy fumes of HCl (or makes a chloride salt if a base is present).

Reacts with	Product	Equation
water	carboxylic acid	$CH_3COCl + H_2O \rightarrow CH_3COOH + HCl$
alcohol	ester	$CH_3COCl + CH_3CH_2OH \rightarrow CH_3COOCH_2CH_3 + HCl$
concentrated ammonia	primary amide	$CH_3COCl + 2NH_3 \rightarrow CH_3CONH_2 + NH_4Cl$
primary amine	N-substituted amide	$CH_3COCl + 2CH_3NH_2 \rightarrow CH_3CONHCH_3 + CH_3NH_3^+Cl^-$

With ammonia and amines, the HCl formed is neutralised by a second molecule of the base, so two are needed; with ammonia you see white smoke of ammonium chloride. The reaction with an alcohol is a better way of making esters than using the carboxylic acid, because it is fast and goes to completion.

Esters and polyesters

- **Uses**: many esters smell sweet and fruity, so they are used in perfumes and flavourings (a perfume ester must also be volatile, non-toxic, and unreactive with water). Ethyl ethanoate is a common solvent for glues and inks.
- **Physical properties**: esters are polar but cannot hydrogen bond to each other, so they boil at much lower temperatures than the acids they come from, and they are almost insoluble in water.
- **Acid hydrolysis**: heat under reflux with dilute acid. This is esterification in reverse, so it is reversible and incomplete: $CH_3CH_2COOCH_3 + H_2O \rightleftharpoons CH_3CH_2COOH + CH_3OH$.
- **Alkaline hydrolysis**: heat under reflux with aqueous sodium hydroxide. It goes to completion, giving the salt of the acid and the alcohol: $CH_3CH_2COOCH_3 + NaOH \rightarrow CH_3CH_2COONa + CH_3OH$. The carboxylate ion is not attacked by the alcohol, so the reaction cannot reverse.

Condensation polymers: polyesters

In **condensation polymerisation** monomers join and a small molecule, usually water or HCl, is lost at each link. To build a chain, each monomer needs a reactive group at both ends. A polyester forms from:

- a **dicarboxylic acid and a diol**, losing water at each ester link (slow, needs a catalyst);
- a **diacyl dichloride and a diol**, losing HCl at each link (fast, complete, but the HCl fumes are hazardous);
- a single **hydroxy acid**, whose –OH and –COOH react with neighbouring molecules: 2-hydroxypropanoic acid (lactic acid) gives poly(lactic acid), a biodegradable plastic.

Terylene (PET), used for clothing fibres and drinks bottles, is made from benzene-1,4-dicarboxylic acid and ethane-1,2-diol. Its repeat unit is –OC–C_6H_4–COO–CH_2CH_2–O–. To find the repeat unit of any polyester, take one of each monomer, remove the atoms lost as water or HCl, and join them through the ester link, with bonds left open at both ends.

EXAM TIP

Polyesters contain polar C=O and C–O bonds that can be attacked, so unlike poly(alkenes) they can be **hydrolysed**: by acid, giving back the dicarboxylic acid and the diol, or by alkali, giving the diol and the salt of the acid. This makes them biodegradable.

Spectroscopy: what radiation does to molecules

Radiation	Effect on molecules	Used for
radio waves (in a strong magnetic field)	change the spin state of certain nuclei, such as ^{1}H and ^{13}C	NMR spectroscopy and MRI body scanners
microwaves	make polar molecules rotate faster	heating food, which contains water
infrared	make bonds vibrate more strongly	IR spectroscopy to identify bonds
ultraviolet	can break bonds, for example C–Cl or Cl–Cl, into radicals	starting radical reactions

NMR spectroscopy

Nuclei with an odd number of protons or neutrons, such as ^{1}H and ^{13}C, behave like tiny magnets. In a strong magnetic field they can line up with or against the field, and radio waves of exactly the right frequency flip them between the two. The frequency needed depends on the **chemical environment** of each nucleus: electronegative atoms nearby pull electron density away and shift the signal. This shift is measured on the **δ scale** in parts per million (ppm), relative to a reference.

- **TMS, tetramethylsilane, $Si(CH_3)_4$**, is the reference, set at δ = 0. All twelve of its hydrogens are equivalent, so it gives one strong signal, well away from almost every other signal; it is inert, non-toxic, and has a low boiling point so it is easily removed afterwards.
- **Solvents** must not contain ^{1}H, or they would swamp the spectrum: $CDCl_3$ (D is deuterium, ^{2}H) or CCl_4.

^{13}C NMR

Each different carbon environment gives one peak, so the number of peaks is the number of carbon environments. Carbons near electronegative atoms or in C=O groups appear at higher δ. Symmetry matters: propanone has three carbons but only two environments, so two peaks.

Type of carbon	Approximate δ / ppm
C–C (alkyl)	5 to 40
C–O, C–Cl, C–N	40 to 90
C=C, aromatic carbons	90 to 150
C=O in acids and esters	160 to 185
C=O in aldehydes and ketones	190 to 220

High resolution ^{1}H NMR

A proton NMR spectrum gives four pieces of information:

1. **Number of peaks** = the number of different hydrogen environments. Hydrogens in the same environment are **equivalent**; ethanol, CH_3CH_2OH, has three.
2. **Chemical shift**, δ, tells you what is near each hydrogen (table below).
3. **Relative peak area** (the integration trace) is proportional to the number of hydrogens in each environment: 3 : 2 : 1 for ethanol.
4. **Splitting**: each peak is split by the hydrogens on the **neighbouring** carbon atoms. A peak with *n* neighbouring hydrogens splits into **n + 1** lines (the n + 1 rule). Equivalent hydrogens do not split each other, and O–H protons normally appear as a singlet that does not split its neighbours.

Type of hydrogen	Approximate δ / ppm
R–CH_3	0.7 to 1.2
R–CH_2–R	1.2 to 1.4
R–OH (alcohol)	0.5 to 5.0, variable
H–C next to C=O (H–C–C=O)	2.0 to 2.6
H–C next to O (H–C–O, alcohols, ethers, esters)	3.3 to 4.3
H–C=C (alkenes)	4.5 to 6.0
aldehyde, R–CHO	9.0 to 10.0
carboxylic acid, R–COOH	10.0 to 12.0

Neighbouring H atoms (n)	0	1	2	3	4
Lines in the peak (n + 1)	1, singlet	2, doublet	3, triplet	4, quartet	5, quintet
Relative line heights	1	1 : 1	1 : 2 : 1	1 : 3 : 3 : 1	1 : 4 : 6 : 4 : 1

EXAM TIP

The ethyl group, CH_3CH_2–, is the pattern to recognise: a **triplet** of area 3 (the CH_3, next to a CH_2) and a **quartet** of area 2 (the CH_2, next to a CH_3). A singlet means no hydrogens on the neighbouring carbons, for example a CH_3 joined to a C=O or an O.

Mass spectrometry

In the mass spectrometer an electron is knocked off each molecule, forming the **molecular ion**, M^+, which is both a positive ion and a radical:

$$M \rightarrow [M]^{+\bullet} + e^-$$

The molecular ion peak is the one with the highest m/z (ignoring the small M+1 peak from ^{13}C), and it gives the relative molecular mass. Many molecular ions break up, or **fragment**: a covalent bond breaks to give a positive ion, which is detected, and a radical, which is not: $[M]^{+\bullet} \rightarrow X^+ + Y^\bullet$. The more stable the ion, the taller its peak; acylium ions, $R–C{\equiv}O^+$, and secondary or tertiary carbocations are especially common.

m/z	Common fragment ion
15	CH_3^+
29	$C_2H_5^+$ or CHO^+
43	CH_3CO^+ or $C_3H_7^+$
57	$C_2H_5CO^+$ or $C_4H_9^+$

Telling pentan-2-one from pentan-3-one

1. Both are $C_5H_{10}O$, so both give the molecular ion at m/z 86.

2. Pentan-3-one, $CH_3CH_2COCH_2CH_3$, breaks next to the C=O to give $[CH_3CH_2CO]^+$ at m/z 57 (and $C_2H_5^+$ at 29): $[CH_3CH_2COCH_2CH_3]^{+\bullet} \rightarrow CH_3CH_2CO^+ + \bullet CH_2CH_3$.

3. Pentan-2-one, $CH_3COCH_2CH_2CH_3$, gives a strong peak at m/z 43 from CH_3CO^+ and one at 71 from $CH_3CH_2CH_2CO^+$, but no peak at 57.

The same molecular ion but different fragments: fragmentation patterns tell isomers apart.

Chromatography

Chromatography separates the components of a mixture between a **mobile phase**, which moves, and a **stationary phase**, which does not. Each component is shared between the two phases to a different extent: one that dissolves better in the mobile phase, or is held less strongly by the stationary phase, travels faster. A solid stationary phase separates by **adsorption**; a liquid stationary phase separates by **relative solubility**.

- **Thin-layer chromatography (TLC)**: the stationary phase is a thin layer of silica or alumina on a plate; the mobile phase is a solvent. Draw the start line in **pencil** (ink would dissolve and run), spot tiny amounts of each sample, and stand the plate in a covered tank with the solvent below the start line. Remove it before the solvent reaches the top, mark the solvent front, and show colourless spots with UV light (or ninhydrin spray for amino acids). Wear gloves, so skin oils do not contaminate the plate, and work in a fume cupboard.
- $R_f = \dfrac{\text{distance moved by the spot}}{\text{distance moved by the solvent front}}$. Compare with known substances run on the same plate. Similar compounds may have similar R_f values and not separate.
- **Column chromatography** uses the same principles in a vertical column, so larger amounts can be separated and collected.
- **Gas chromatography (GC)**: the mobile phase is an unreactive carrier gas such as helium or nitrogen; the stationary phase is a high-boiling liquid coated on a solid support inside a long coiled column in an oven. Each component has a characteristic **retention time**, and the area under its peak shows how much is present. It suits volatile liquids.
- **HPLC** (high-performance liquid chromatography): the mobile phase is a liquid forced under high pressure through a column packed with very small solid particles, giving fast, efficient separations.
- **GC–MS**: each component leaving a gas chromatograph is fed straight into a mass spectrometer, and its spectrum is matched against a database. It is used in forensic science, airport security, drug testing and environmental monitoring.

Putting the techniques together

Identifying an unknown compound

1. A compound contains 54.5% carbon, 9.1% hydrogen and 36.4% oxygen by mass. Divide by A_r: C 54.5 ÷ 12 = 4.54, H 9.1 ÷ 1 = 9.1, O 36.4 ÷ 16 = 2.28. Divide by the smallest: C : H : O = 2 : 4 : 1, so the empirical formula is C_2H_4O (mass 44).

2. The mass spectrum has its molecular ion at m/z 88, so the molecular formula is $C_4H_8O_2$.

3. The IR spectrum has a strong absorption at 1740 cm^{-1} (C=O) but no broad absorption at 2500 to 3300 cm^{-1}, so there is no O–H: the compound is an ester, not a carboxylic acid.

4. The ^1H NMR spectrum shows three peaks: δ 4.1, quartet, area 2; δ 2.0, singlet, area 3; δ 1.3, triplet, area 3.

5. The quartet and triplet together are an ethyl group, and the CH_2 at δ 4.1 must be joined to oxygen (O–CH_2CH_3). The singlet of area 3 at δ 2.0 is a CH_3 next to C=O with no neighbouring hydrogens (CH_3CO–).

6. The compound is **ethyl ethanoate**, $CH_3COOCH_2CH_3$. Check: its ^{13}C NMR spectrum would show four peaks, including one near 171 ppm for the ester C=O.

Quick checks

QUICK CHECK

Which of these molecules has a chiral centre?

- propan-2-ol, $CH_3CH(OH)CH_3$
- butan-2-ol, $CH_3CH(OH)CH_2CH_3$
- propanone, CH_3COCH_3
- 2-methylpropan-2-ol, $(CH_3)_3COH$

In butan-2-ol, carbon 2 carries H, OH, CH_3 and C_2H_5: four different groups. In propan-2-ol the two CH_3 groups are the same.

QUICK CHECK

A single enantiomer of 2-bromobutane reacts with OH^- by an S_N2 mechanism. What is the product?

- a racemic mixture of butan-2-ol
- an optically inactive alcohol, because the carbocation is planar
- one enantiomer of butan-2-ol, with the arrangement inverted
- but-2-ene

S_N2 attack comes from the side opposite the bromine in one step, inverting the centre. There is no carbocation; a racemate would point to S_N1.

QUICK CHECK

Which reagent gives a positive result with propanal but not with propanone?

- 2,4-DNPH
- Tollens' reagent
- iodine in sodium hydroxide
- $LiAlH_4$

Only the aldehyde is oxidised, reducing Ag^+ to a silver mirror. 2,4-DNPH reacts with both; the iodoform test is positive for propanone (a methyl ketone) but not propanal.

QUICK CHECK

Why is the product of HCN addition to ethanal optically inactive?

- The product has no chiral centre
- HCN is a weak acid
- The reaction goes by S_N1
- CN^- attacks the planar C=O equally from both sides, forming a racemic mixture

2-hydroxypropanenitrile does have a chiral centre, but both enantiomers form in equal amounts, so their rotations cancel.

QUICK CHECK

Which reaction is the best way to make ethyl ethanoate quickly and completely?

- ethanoic acid + ethanol with a little concentrated H_2SO_4
- ethanoyl chloride + ethanol
- ethanoic acid + ethanal
- ethanoyl chloride + water

Acyl chlorides react with alcohols at room temperature, quickly and irreversibly. Acid-catalysed esterification with the acid is slow and reaches equilibrium.

QUICK CHECK

How many peaks are there in the ^{1}H NMR spectrum of methyl propanoate, $CH_3CH_2COOCH_3$, and what is the splitting of the CH_3 joined to oxygen?

- 2 peaks; triplet
- 3 peaks; quartet
- **3 peaks; singlet**
- 4 peaks; singlet

There are three environments: CH_3 of the ethyl group (triplet), CH_2 (quartet) and O–CH_3. The O–CH_3 is a singlet because its neighbour across the oxygen is the C=O carbon, which carries no hydrogen.

QUICK CHECK

Which is NOT a reason for using TMS as the reference in NMR?

- it gives a single strong peak
- it is inert and has a low boiling point
- **it dissolves the sample without giving any peaks**
- its peak lies away from most other peaks

TMS is added in small amounts as a reference and gives a peak at δ 0. Dissolving the sample without adding peaks is the job of a solvent such as $CDCl_3$.

QUICK CHECK

On a TLC plate the solvent front moved 8.0 cm and a spot moved 2.6 cm. What is its R_f value?

- 3.1
- 0.26
- 0.33
- 5.4

$R_f = 2.6 \div 8.0 = 0.325 \approx 0.33$. R_f is always less than 1; 3.1 is the ratio upside down.