

# Radiative-convective equilibrium in a grey atmosphere

Marco Casari

University of Turin

Complex systems in climate physics, 3 October 2023

- Average vertical temperature profile  $T(t, z)$  of atmosphere.

- Average vertical temperature profile  $T(t, z)$  of atmosphere.
- Radiative Transfer Equation (RTE).

- Average vertical temperature profile  $T(t, z)$  of atmosphere.
- Radiative Transfer Equation (RTE).
- Fluid dynamics equations.

- Thermodynamic energy equation in Local Thermodynamic Equilibrium (LTE):

$$\frac{\partial T}{\partial t} = -\frac{1}{\rho c_P} \frac{\partial q}{\partial z} \quad . \quad (1)$$

- Thermodynamic energy equation in Local Thermodynamic Equilibrium (LTE):

$$\frac{\partial T}{\partial t} = -\frac{1}{\rho c_P} \frac{\partial q}{\partial z} \quad . \quad (1)$$

- Radiative-convective equilibrium.

- Thermodynamic energy equation in Local Thermodynamic Equilibrium (LTE):

$$\frac{\partial T}{\partial t} = -\frac{1}{\rho c_P} \frac{\partial q}{\partial z} \quad . \quad (1)$$

- Radiative-convective equilibrium.
- Grey atmosphere.

- Hypotheses on the planet.



- Hypotheses on the planet.
- Hypotheses on the composition of atmosphere.

- Hypotheses on the planet.
- Hypotheses on the composition of atmosphere.
- Hypotheses on total heat flux.
- Resulting thermodynamic energy equation:

$$\frac{\partial T}{\partial t} = -\frac{1}{\rho c_P} \frac{\partial}{\partial z} (E_U - E_D) \quad . \quad (2)$$

- Relation between pressure and altitude:

$$P(z) = P_g \exp \left( - \frac{z - z_g}{z_0} \right) \quad . \quad (3)$$

- Relation between optical depth and pressure:

$$\delta(P) = \frac{\mu_m}{g} (P - P_{\text{TOA}}) \quad . \quad (4)$$

- Relation between optical depth and altitude:

$$\delta(z) = \frac{\mu_m}{g} \left( P_g \exp \left( - \frac{z - z_g}{z_0} \right) - P_{\text{TOA}} \right) \quad . \quad (5)$$

- RTE for non-scattering medium in LTE:

$$\frac{1}{\mu} \frac{\partial L}{\partial z} = B_\nu - L \quad . \quad (6)$$

## Equations in radiative equilibrium

- RTE for non-scattering medium in LTE:

$$\frac{1}{\mu} \frac{\partial L}{\partial z} = B_\nu - L \quad . \quad (6)$$

- Integration over frequency and solid angle.
- Diffusion approximation:  $\delta' = D\delta$ .

## Equations in radiative equilibrium

- RTE for non-scattering medium in LTE:

$$\frac{1}{\mu} \frac{\partial L}{\partial z} = B_\nu - L \quad . \quad (6)$$

- Integration over frequency and solid angle.
- Diffusion approximation:  $\delta' = D\delta$ .
- Equations for irradiances:

$$-\frac{\partial}{\partial \delta'} E_U(t, \delta') = \sigma T(t, \delta')^4 - E_U(t, \delta') \quad , \quad (7)$$

$$\frac{\partial}{\partial \delta'} E_D(t, \delta') = \sigma T(t, \delta')^4 - E_D(t, \delta') \quad . \quad (8)$$

- Steady state.

- Steady state.
- Initial conditions on irradiances:

$$E_D(0) = 0 \quad , \quad (9)$$

$$E_U(0) = S_t \quad . \quad (10)$$



- Steady state.
- Initial conditions on irradiances:

$$E_D(0) = 0 \quad , \quad (9)$$

$$E_U(0) = S_t \quad . \quad (10)$$

- Additional relations:

$$\frac{d}{d\delta'} (E_U(\delta') - E_D(\delta')) = 0 \quad , \quad (11)$$

$$E_U(\delta') - E_D(\delta') = S_t \quad . \quad (12)$$

- Steady state.
- Initial conditions on irradiances:

$$E_D(0) = 0 \quad , \quad (9)$$

$$E_U(0) = S_t \quad . \quad (10)$$

- Additional relations:

$$\frac{d}{d\delta'} (E_U(\delta') - E_D(\delta')) = 0 \quad , \quad (11)$$

$$E_U(\delta') - E_D(\delta') = S_t \quad . \quad (12)$$

- Initial condition on temperature:

$$T(0) = \left( \frac{S_t}{2\sigma} \right)^{\frac{1}{4}} \quad . \quad (13)$$

- Temperature:

$$T(\delta) = \left( \frac{S_t}{2\sigma} (1 + D\delta) \right)^{\frac{1}{4}} . \quad (14)$$

- Irradiances:

$$E_U(\delta) = \frac{S_t}{2} (2 + D\delta) \quad , \quad (15)$$

$$E_D(\delta) = \frac{S_t}{2} D\delta \quad . \quad (16)$$


- Normalisation:


$$Y_0 = \frac{T^4}{T_0^4} \quad , \quad Y_1 = \frac{E_U}{S_t} \quad , \quad Y_2 = \frac{E_U}{S_t} \quad (17)$$

- Resulting system of ODEs:

$$\begin{cases} \frac{dY_0}{d\delta} = \frac{D}{2} \\ \frac{dY_1}{d\delta} = D(Y_1 - Y_0) \\ \frac{dY_2}{d\delta} = D(Y_0 - Y_2) \end{cases} . \quad (18)$$

- Runge-Kutta method of order 4.
- Non-uniform step size.





- Iterative procedure.

- Iterative procedure.
- Additional normalisation:

$$Y_3 = \frac{T}{T_0} \quad . \quad (19)$$



- Iterative procedure.
- Additional normalisation:

$$Y_3 = \frac{T}{T_0} \quad . \quad (19)$$

- Resulting system of PDEs:

$$\begin{cases} \frac{\partial}{\partial t} Y_3(t, \delta) = \frac{\mu_m S_t}{c_P T_0} \frac{\partial}{\partial \delta} (Y_1(t, \delta) - Y_2(t, \delta)) \\ \frac{\partial}{\partial \delta} Y_1(t, \delta) = D(Y_1(t, \delta) - Y_3(t, \delta)^4) \\ \frac{\partial}{\partial \delta} Y_2(t, \delta) = D(Y_3(t, \delta)^4 - Y_2(t, \delta)) \end{cases} \quad . \quad (20)$$

- Euler method for temporal integration.
- Constant time step.

# Errors of time integration at steady state

Radiative-convective equilibrium in a grey atmosphere


Marco Casari

Introduction

Radiative equilibrium

Radiative-convective equilibrium

Conclusion



- Lapse rate at steady state.

- Lapse rate at steady state.
- Convection faster than radiative processes.

- Lapse rate at steady state.
- Convection faster than radiative processes.
- Convective adjustment if  $-\frac{\partial T}{\partial z} > \Gamma_0$ .

# Temperature plot

Radiative-convective equilibrium in a grey atmosphere


Marco Casari


Introduction

Radiative equilibrium

Radiative-convective equilibrium

Conclusion





- Vertical temperature profile of a grey atmosphere described in general by a system of PDEs and at the steady state by a system of ODEs.
- Negligible errors for ODEs numerical solutions.
- Possible improvements on description of physical phenomena.
- Possible improvements on precision of PDEs numerical solutions.