

Study Nuclear Winter with a radiative-convective climate model

Marco Casari

July 30, 2023

Abstract

Reflections from atmospheric layers have negligible intensities compared to the first one.[1, p. 470]

1 Introduction

2 Methods

The definition of wavenumber $\nu = \frac{1}{\lambda}$ is used in this work, where λ is the wavelength.

2.1 Longwave radiation

At wavelengths $\lambda \geq 4\mu\text{m}$, solar radiation has lower intensities than radiation emitted by Earth's surface and atmosphere at the same wavelengths. Moreover, it presents negligible scattering in atmosphere with respect to absorption. For these reasons longwave radiation is considered to be emitted only by Earth's surface and atmosphere.[1, p. 468]

2.2 Shortwave radiation

At wavelengths $\lambda < 4\mu\text{m}$, solar radiation has much greater intensity than radiation emitted from Earth's surface and atmosphere. Both scattering and absorption by gases, aerosols and clouds of atmosphere dissipates solar radiation.[1, p. 469]

Specific intensity of solar radiation can be expressed by a differential equation whose resolution is complex even applying approximations and numerical methods.[1, p. 469]

A lower complexity parametrisation is adopted instead, where the atmosphere is divided in a given number of layers and radiation is absorbed, scattered and reflected between each layer. Multiple reflections can occur from each layer but only one is considered in this model because successive re-

2.3 Numerical approach

Euler's method is used to solve the Ordinary Differential Equation (ODE) for the time-dependant temperature function.[1, p. 472] A solution for each atmospheric layer is evaluated, hence the resulting values are triplets of temperature, altitude (i.e. proxy for the atmospheric layer) and time (i.e. simulation time). Further information on storage and plotting of data are presented in Section B.1.

3 Results

3.1 Stability analysis

4 Discussion

A Source code

In this section the C++ code used to obtain the results presented in this work is shown and commented.

A.1 Classes

B Supplementary information

B.1 Plotting

Software Gnuplot is used to generate plots shown in this work. Output values from the simulation are stored in a DAT file with the following structure, line spacing between data blocks is important:

```
t[0]      P[0]      T[0][0]
...      ...      ...
t[0]      P[N_P - 1] T[0][N_P - 1]

t[1]      P[0]      T[1][0]
...      ...      ...
t[1]      P[N_P - 1] T[1][N_P - 1]

...

t[N_t - 1] P[0]      T[N_t - 1][0]
...      ...      ...
t[N_t - 1] P[N_P - 1] T[N_t - 1][N_P - 1]
```

Value N_t is not fixed a priori since it is the number of temporal steps needed to reach convergence, instead N_P is the number of atmospheric layers.

References

- [1] V. Ramanathan and J. A. Coakley Jr., “Climate modeling through radiative-convective models,” *Reviews of Geophysics*, vol. 16, no. 4, pp. 465–489, 1978. DOI: <https://doi.org/10.1029/RG016i004p00465>. eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/RG016i004p00465>. [Online]. Available: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/RG016i004p00465>.