

7. Lagrange interpolation in Newton's divided differences form

 $f: [0, 1] \rightarrow \mathbb{R}, f(x) = \cos(\pi x)$

x	0	$1/3$	$1/2$	1
$f(x)$	1	$1/2$	0	-1

$$m = 3$$

$$\Downarrow$$

$$\boxed{m=3}$$

(a) Lagrange polynomial

$$L_3 f(x) = N_3 f(x) = f(x_0) + f[x_0, x_1](x-x_0) \\ + f[x_0, x_1, x_2](x-x_0)(x-x_1) \\ + f[x_0, x_1, x_2, x_3](x-x_0)(x-x_1)(x-x_2)$$

x	$f(x)$	$D^1 f$	$D^2 f$	$D^3 f$
0	1	$-3/2$	-3	$9/2$
$1/3$	$1/2$	-3	$3/2$	
$1/2$	0	-2		
1	-1			

$$D^i f(x_0) = f[x_0, \dots, x_i]$$

Hence, $L_3 f(x) = 1 - \frac{3}{2}(x-0) - 3(x-0)(x-\frac{1}{3}) + \frac{9}{2}(x-0)(x-\frac{1}{3})(x-\frac{1}{2})$

$$= 1 - \frac{3}{2}x - \frac{3x(3x-1)}{3} + \frac{9x(3x-1)(2x-1)}{12}$$

$$\Rightarrow L_3 f(x) = 1 - \frac{3}{2}x - x(3x-1) + \frac{3x(3x-1)(2x-1)}{4}$$

$$= \frac{9}{2}x^3 - \frac{27}{4}x^2 + \frac{1}{4}x + 1$$

b) Approx. for $\cos(\frac{\pi}{5})$

$$\cos(\frac{\pi}{5}) = f(\frac{1}{5}) \approx L_3 f(\frac{1}{5}) ,$$

$$\text{where } L_3 f(\frac{1}{5}) = \frac{9}{2}(\frac{1}{5})^3 - \frac{27}{4}(\frac{1}{5})^2 + \frac{1}{4} \cdot \frac{1}{5} + 1 = \dots = 0,816$$

c) The error $R_3 f$

$$R_3 f(x) = \frac{u(x)}{(3+1)!} f^{(4)}(\xi) , \text{ where } \xi \in (0, 1)$$

$$f(x) = \cos(\pi x) \Rightarrow f'(x) = -\pi \sin(\pi x)$$

$$f''(x) = -\pi^2 \cos(\pi x)$$

$$f'''(x) = \pi^3 \sin(\pi x)$$

$$\boxed{f^{(4)}(x) = \pi^4 \cos(\pi x)}$$

$$\text{and } u(x) = (x-x_0)(x-x_1)(x-x_2)(x-x_3)$$

$$= (x-0)(x-\frac{1}{3})(x-\frac{1}{2})(x-1)$$

$$\text{Then } R_3 f(x) = \frac{x(3x-1)(2x-1)(x-1)}{6 \cdot 4!} \pi^4 \cos(\pi \cdot \xi) ,$$

$$\xi \in (0, 1).$$

d) A bound for the error

$$\begin{aligned}
 0 \leq |R_3 f(x)| &= \left| \frac{x(3x-1)(2x-1)(x-1)}{6 \cdot 4!} \pi^4 \cos(\pi \cdot x) \right| \\
 &\leq \frac{|x(3x-1)(2x-1)(x-1)|}{6 \cdot 24} \cdot \pi^4 \cdot \underbrace{|\cos(\pi \cdot x)|}_{\leq 1} \\
 &\leq \frac{\pi^4}{144} \cdot \underbrace{\left| \frac{v(x)}{x(3x-1)(2x-1)(x-1)} \right|}_{\leq \max_{x \in [0,1]} v(x)}, \quad x \in [0,1]
 \end{aligned}$$

• In this case, v is of degree 4 and is difficult to determine by hand the maximum value (see Lecture 5 - pag. 7 for an example)

e) A bound for the error of approx. $f(\frac{1}{5})$

$$0 \leq |R_3 f(\frac{1}{5})| \leq \frac{\pi^4}{6 \cdot 24} \cdot \frac{1}{5} \cdot \frac{2}{5} \cdot \frac{3}{5} \cdot \frac{4}{5} = \frac{\pi^4}{5^4 \cdot 6} = \frac{1}{6} \left(\frac{\pi}{5} \right)^4$$

$$\Rightarrow 0 \leq |R_3 f(\frac{1}{5})| \leq \frac{1}{6} \cdot \left(\frac{\pi}{5} \right)^4 \approx 0.0259$$

$$\text{Indeed, } |L_3 f(\frac{1}{5}) - f(\frac{1}{5})| = 0.00698 < 0.0259$$

□