

Week 4 Quiz

TOTAL POINTS 9

1. Person 1 has hazard $h_1(t) = 1$, and Person 2 has hazard $h_2(t) = 2$. What is the probability of dying within the first year for each patient?

1 point

Hint:

The survival function $S(t)$ in terms of the hazard function is:

$$S(t) = e^{-\int_0^t h(s)ds}$$

- ☐ 0.63, 0.86
- ☐ 0.6, 0.6
- ☒ 0.37, 0.14

2. Let $T > 0$.

1 point

For patient 1, let the survival function be $S_1(t)$ and the hazard function be $h_1(t)$.

For patient 2, let the survival function be $S_2(t)$ and the hazard function be $h_2(t)$

You see that $S_1(T) > S_2(T)$. The survival probability of patient 1 at time T is higher than the survival probability of patient 2 at time T.

Which of the following is true about the hazard of patient 1 and 2 at time T?

Hint:

$$S(t) = e^{-\int_0^t h(s)ds}$$

- ☐ $h_1(T) > h_2(T)$
- ☒ $h_1(T) < h_2(T)$
- ☐ $h_1(T) = h_2(T)$
- ☐ None of the above

3. Now assume that the hazards for patient 1, h_1 and for patient 2, h_2 are proportional to each other. Also assume that $S_1(T) > S_2(T)$ for some $T > 0$.

1 point

Then which of the following is true about the hazards?

- ☐ $h_1(T) < h_2(T)$
- ☒ $h_1(T) = h_2(T)$
- ☐ $h_1(T) > h_2(T)$

4. You've fit a Cox model on 2 features: age and smoking status.

1 point

The coefficients of these features are:

$$\beta_{age} = 0.9 \text{ and } \beta_{smoker} = 10.0.$$

What is the hazard ratio between Person 1, a 40 year old non-smoker, and Person 2, a 30 year old smoker?

Recall that Cox Proportional Hazards assumes a model of the form:

$$h(t) = \lambda_0(t)e^{(\beta_{age} \times Age + \beta_{smoker} \times Smoker)}$$

We're asking you to find the ratio:

$$\frac{h_1(t)}{h_2(t)}$$

☐ 2.7

☒ 0.36

☐ 2.64

5. You've fit a cox model and have the following coefficients:

1 point

$$\beta_{female} = -1.0$$

$$\beta_{age} = 1.0,$$

$$\beta_{BP} = 0.6$$

$$h(t) = \lambda_0(t)e^{((\beta_{female} \times female) + (\beta_{age} \times Age) + (\beta_{BP} \times BP))}$$

Which of the following interpretations is most correct?

- ☐ All other things held equal, having higher BP decreases your risk
- ☐ All other things held equal, having lower age increases your risk
- ☒ All other things held equal, being a female decreases your risk

6. Assume $h_1(t) = t$, and $h_2(t) = 1.0$. At which time $T > 0$ does $S_1(T) = S_2(T)$?

1 point

- ☐ 2
- ☒ 1
- ☐ None of the above
- ☐ 0.5

7. Using the Nelson-Aalen estimator estimate $H(7)$, the value of the cumulative hazard at $t=7$ for this dataset.

1 point

ID	Outcome
1	3
2	4
3	8
4	6+

The Nelson-Aalen estimator is:

$$H(t) = \sum_{i=0}^t \frac{d_i}{n_i}$$

- ☐ 5/9
- ☐ 8/11
- ☒ 7/12

8. Which risk assignments would make this pair **concordant**?

1 point

Patient 1



T

10

Patient 2



5+

- ☐ 0.3, 0.8
- ☐ 0.5, 0.3
- ☐ 0.5, 0.5
- ☒ The pair is not permissible

9. Compute the Harrell C-index for the following dataset and risk scores:

1 point

ID	Outcome	Score
1	4	1.6
2	6+	1.2
3	5	0.8
4	7	0.1

Step 1: Find all the permissible pairs

Step 2: of the permissible pairs, determine which ones are concordant.

Step 3: of the permissible pairs, determine which ones are risk ties.

Harrell's c-index = $\frac{\text{concordant} + 0.5 \times \text{riskties}}{\text{permissible}}$

☐ 1.0

☒ 0.8

☐ 0.7

☐ I, **Mirko Jerber Rodriguez Mallma**, understand that submitting work that isn't my own may result in permanent failure of this course or deactivation of my Coursera account.

[Learn more about Coursera's Honor Code](#)

