

STAKEHOLDER 1: Hospital Management

Q1: Which age groups have the highest disease prevalence?

Patients aged 50–59 have the highest disease prevalence, while the 30–39 age group has the lowest. Disease occurrence increases with age, showing that older adults are more affected.

Q2: What proportion of patients suffer from cardiovascular disease?

About 49% of patients have cardiovascular disease, while 51% do not. This means nearly half of the population is affected by the disease.

Q3: Which risk factors are most common among patients?

High cholesterol and physical inactivity are the most common risk factors. Elevated glucose levels are also present in some patients, while smoking and alcohol consumption are relatively low.

Q4: How does BMI relate to cardiovascular disease?

Patients with normal BMI have the lowest disease rates. Disease prevalence increases among overweight patients and is highest among obese patients. Higher BMI is associated with greater cardiovascular disease risk.

STAKEHOLDER 2: Cardiologists

Q1: How do blood pressure levels vary among diseased patients?

Many diseased patients have elevated systolic and diastolic blood pressure levels. High blood pressure is common and appears to be a major warning sign for cardiovascular disease.

Q2: What is the relationship between cholesterol and cardiovascular disease?

As cholesterol levels increase, the proportion of cardiovascular disease cases also increases. Patients with well-above-normal cholesterol have the highest disease risk.

Q3: Does glucose level affect cardiovascular disease risk?

Higher glucose levels are linked with higher cardiovascular disease risk. Although the relationship is weaker than blood pressure and cholesterol, it is still clinically important.

Q4: Which patient groups require early intervention?

Patients with systolic blood pressure above 140 mmHg and diastolic blood pressure above 90 mmHg are at the greatest risk and should receive early medical attention.

STAKEHOLDER 3: Public Health Department

Q1: Which lifestyle factors contribute most to disease?

Physical inactivity is the most common lifestyle-related risk factor. Smoking and alcohol consumption have a much smaller presence in this dataset.

Q2: Which demographic groups are at highest risk?

Adults aged 60 and above have the highest disease risk, followed by those aged 50–59. Disease risk increases steadily with age in both genders.

Q3: How does obesity influence disease prevalence?

Overweight and obese individuals experience higher rates of cardiovascular disease. Obese patients show the highest disease prevalence, indicating a strong relationship between obesity and heart disease.

Q4: What preventive programs should be prioritized?

The most important programs include blood pressure control, obesity and weight management, cholesterol screening, and regular cardiovascular checkups for adults aged 50 years and above.

Conclusion

Age, blood pressure, cholesterol, and BMI are the strongest predictors of cardiovascular disease. Focusing on blood pressure control, weight management, cholesterol monitoring, and screening of adults aged 50+ can significantly reduce disease risk.