

# Computational Photography

Matthias Zwicker, Daniel Donatsch  
University of Bern  
Fall 2012

# Project 3

- Topics
  - Linear filtering
  - Convolution
  - Discrete Fourier transform
  - Frequency domain filtering
  - Inverse filtering/deconvolution

# Linear Filter

$$\mathcal{F}(f(x)) = e^{f(x)}$$

# Linear Filter

$$\mathcal{F}(f(x)) = e^{f(x)}$$

$$\mathcal{F}(f_1(x)) + \mathcal{F}(f_2(x))$$

# Linear Filter

$$\mathcal{F}(f(x)) = e^{f(x)}$$

$$\mathcal{F}(f_1(x)) + \mathcal{F}(f_2(x)) = e^{f_1(x)} + e^{f_2(x)}$$

# Linear Filter

$$\mathcal{F}(f(x)) = e^{f(x)}$$

$$\mathcal{F}(f_1(x)) + \mathcal{F}(f_2(x)) = e^{f_1(x)} + e^{f_2(x)}$$

$$\mathcal{F}(f_1(x) + f_2(x))$$

# Linear Filter

$$\mathcal{F}(f(x)) = e^{f(x)}$$

$$\mathcal{F}(f_1(x)) + \mathcal{F}(f_2(x)) = e^{f_1(x)} + e^{f_2(x)}$$

$$\mathcal{F}(f_1(x) + f_2(x)) = e^{f_1(x) + f_2(x)}$$

# Linear Filter

$$\mathcal{F}(f(x)) = e^{f(x)}$$

$$\mathcal{F}(f_1(x)) + \mathcal{F}(f_2(x)) = e^{f_1(x)} + e^{f_2(x)}$$

$$\mathcal{F}(f_1(x) + f_2(x)) = e^{f_1(x) + f_2(x)} = e^{f_1(x)} e^{f_2(x)}$$



# Linear Filter

$$\mathcal{F}(f(x)) = e^{f(x)}$$

$$\mathcal{F}(f_1(x)) + \mathcal{F}(f_2(x)) = e^{f_1(x)} + e^{f_2(x)}$$

$$\mathcal{F}(f_1(x) + f_2(x)) = e^{f_1(x) + f_2(x)} = e^{f_1(x)} e^{f_2(x)}$$

$\Rightarrow$  **non-linear**

# Linear Filter

$$\mathcal{F}(f(x)) = e^{f(x)}$$

$$\mathcal{F}(f_1(x)) + \mathcal{F}(f_2(x)) = e^{f_1(x)} + e^{f_2(x)}$$

$$\mathcal{F}(f_1(x) + f_2(x)) = e^{f_1(x) + f_2(x)} = e^{f_1(x)} e^{f_2(x)}$$

$\Rightarrow$  **non-linear**

# Shift Invariance

$$\mathcal{F}(f(x)) = e^{f(x)}$$

# Shift Invariance

$$\mathcal{F}(f(x)) = e^{f(x)}$$

$$\mathcal{F}(S_\delta(f(x))) =$$

# Shift Invariance

$$\mathcal{F}(f(x)) = e^{f(x)}$$

$$\mathcal{F}(S_\delta(f(x))) = \mathcal{F}((f(x + \delta))) =$$

# Shift Invariance

$$\mathcal{F}(f(x)) = e^{f(x)}$$

$$\mathcal{F}(S_\delta(f(x))) = \mathcal{F}((f(x + \delta))) = e^{f(x+\delta)}$$

# Shift Invariance

$$\mathcal{F}(f(x)) = e^{f(x)}$$

$$\mathcal{F}(S_\delta(f(x))) = \mathcal{F}((f(x + \delta))) = e^{f(x+\delta)}$$

$$S_\delta(\mathcal{F}(f(x))) =$$

# Shift Invariance

$$\mathcal{F}(f(x)) = e^{f(x)}$$

$$\mathcal{F}(S_\delta(f(x))) = \mathcal{F}((f(x + \delta))) = e^{f(x+\delta)}$$

$$S_\delta(\mathcal{F}(f(x))) = S_\delta(e^{f(x)}) =$$



# Shift Invariance

$$\mathcal{F}(f(x)) = e^{f(x)}$$

$$\mathcal{F}(S_\delta(f(x))) = \mathcal{F}((f(x + \delta))) = e^{f(x+\delta)}$$

$$S_\delta(\mathcal{F}(f(x))) = S_\delta(e^{f(x)}) = e^{f(x+\delta)}$$

# Shift Invariance

$$\mathcal{F}(f(x)) = e^{f(x)}$$

$$\mathcal{F}(S_\delta(f(x))) = \mathcal{F}((f(x + \delta))) = e^{f(x+\delta)}$$

$$S_\delta(\mathcal{F}(f(x))) = S_\delta(e^{f(x)}) = e^{f(x+\delta)}$$

$\Rightarrow$  **shift invariant**

# Fourier Transform of Gaussian

$$F(w) = \int_{-\infty}^{\infty} f(x) e^{-i2\pi wx} dx$$

# Fourier Transform of Gaussian

$$\begin{aligned} F(w) &= \int_{-\infty}^{\infty} f(x) e^{-i2\pi wx} dx \\ &= \int_{-\infty}^{\infty} A e^{-ax^2} e^{-i2\pi wx} dx \end{aligned}$$

# Fourier Transform of Gaussian

$$\begin{aligned} F(w) &= \int_{-\infty}^{\infty} f(x) e^{-i2\pi wx} dx \\ &= \int_{-\infty}^{\infty} A e^{-ax^2} e^{-i2\pi wx} dx \\ &= A \int_{-\infty}^{\infty} e^{-(ax^2 + i2\pi wx)} dx \end{aligned}$$

# Fourier Transform of Gaussian

$$\begin{aligned} F(w) &= \int_{-\infty}^{\infty} f(x) e^{-i2\pi wx} dx \\ &= \int_{-\infty}^{\infty} A e^{-ax^2} e^{-i2\pi wx} dx \\ &= A \int_{-\infty}^{\infty} e^{-(ax^2 + i2\pi wx)} dx \end{aligned}$$

completing the square:

$$ax^2 + i2\pi wx = \left( \sqrt{a}x + \frac{iw\pi}{\sqrt{a}} \right)^2 + \frac{w^2\pi^2}{a}$$

# Fourier Transform of Gaussian

$$\begin{aligned} F(w) &= \int_{-\infty}^{\infty} f(x) e^{-i2\pi wx} dx \\ &= \int_{-\infty}^{\infty} A e^{-ax^2} e^{-i2\pi wx} dx \\ &= A \int_{-\infty}^{\infty} e^{-(ax^2 + i2\pi wx)} dx \\ &= A \int_{-\infty}^{\infty} e^{-\left(\sqrt{a}x + \frac{iw\pi}{\sqrt{a}}\right)^2 - \frac{w^2\pi^2}{a}} dx \end{aligned}$$

completing the square:

$$ax^2 + i2\pi wx = \left(\sqrt{a}x + \frac{iw\pi}{\sqrt{a}}\right)^2 + \frac{w^2\pi^2}{a}$$

# Fourier Transform of Gaussian

$$\begin{aligned}F(w) &= \int_{-\infty}^{\infty} f(x) e^{-i2\pi wx} dx \\&= \int_{-\infty}^{\infty} A e^{-ax^2} e^{-i2\pi wx} dx \\&= A \int_{-\infty}^{\infty} e^{-(ax^2 + i2\pi wx)} dx \\&= A \int_{-\infty}^{\infty} e^{-\left(\sqrt{a}x + \frac{iw\pi}{\sqrt{a}}\right)^2 - \frac{w^2\pi^2}{a}} dx \\&= A e^{-w^2\pi^2/a} \int_{-\infty}^{\infty} e^{-a\left(x + \frac{iw\pi}{a}\right)^2} dx\end{aligned}$$

completing the square:

$$ax^2 + i2\pi wx = \left(\sqrt{a}x + \frac{iw\pi}{\sqrt{a}}\right)^2 + \frac{w^2\pi^2}{a}$$



# Fourier Transform of Gaussian

$$\begin{aligned}F(w) &= \int_{-\infty}^{\infty} f(x) e^{-i2\pi wx} dx \\&= \int_{-\infty}^{\infty} A e^{-ax^2} e^{-i2\pi wx} dx \\&= A \int_{-\infty}^{\infty} e^{-(ax^2 + i2\pi wx)} dx \\&= A \int_{-\infty}^{\infty} e^{-\left(\sqrt{a}x + \frac{iw\pi}{\sqrt{a}}\right)^2 - \frac{w^2\pi^2}{a}} dx \\&= A e^{-w^2\pi^2/a} \int_{-\infty}^{\infty} e^{-a\left(x + \frac{iw\pi}{a}\right)^2} dx\end{aligned}$$

completing the square:

$$ax^2 + i2\pi wx = \left(\sqrt{a}x + \frac{iw\pi}{\sqrt{a}}\right)^2 + \frac{w^2\pi^2}{a}$$

substitute

$$y = x + \frac{iw\pi}{a}$$

# Fourier Transform of Gaussian

$$\begin{aligned}F(w) &= \int_{-\infty}^{\infty} f(x) e^{-i2\pi wx} dx \\&= \int_{-\infty}^{\infty} A e^{-ax^2} e^{-i2\pi wx} dx \\&= A \int_{-\infty}^{\infty} e^{-(ax^2 + i2\pi wx)} dx \\&= A \int_{-\infty}^{\infty} e^{-\left(\sqrt{a}x + \frac{iw\pi}{\sqrt{a}}\right)^2 - \frac{w^2\pi^2}{a}} dx \\&= A e^{-w^2\pi^2/a} \int_{-\infty}^{\infty} e^{-a\left(x + \frac{iw\pi}{a}\right)^2} dx \\&= A e^{-w^2\pi^2/a} \int_{-\infty}^{\infty} e^{-ay^2} dy\end{aligned}$$

completing the square:

$$ax^2 + i2\pi wx = \left(\sqrt{a}x + \frac{iw\pi}{\sqrt{a}}\right)^2 + \frac{w^2\pi^2}{a}$$

substitute

$$y = x + \frac{iw\pi}{a}$$

# Fourier Transform of Gaussian

$$\begin{aligned}F(w) &= \int_{-\infty}^{\infty} f(x) e^{-i2\pi wx} dx \\&= \int_{-\infty}^{\infty} A e^{-ax^2} e^{-i2\pi wx} dx \\&= A \int_{-\infty}^{\infty} e^{-(ax^2 + i2\pi wx)} dx \\&= A \int_{-\infty}^{\infty} e^{-\left(\sqrt{a}x + \frac{iw\pi}{\sqrt{a}}\right)^2 - \frac{w^2\pi^2}{a}} dx \\&= A e^{-w^2\pi^2/a} \int_{-\infty}^{\infty} e^{-a\left(x + \frac{iw\pi}{a}\right)^2} dx \\&= A e^{-w^2\pi^2/a} \int_{-\infty}^{\infty} e^{-ay^2} dy \\&= A e^{-w^2\pi^2/a} \sqrt{\frac{\pi}{a}}\end{aligned}$$

completing the square:

$$ax^2 + i2\pi wx = \left(\sqrt{a}x + \frac{iw\pi}{\sqrt{a}}\right)^2 + \frac{w^2\pi^2}{a}$$

substitute

$$y = x + \frac{iw\pi}{a}$$

# Integral of a Gaussian

$$\int_{-\infty}^{\infty} e^{-ay^2} dy$$

# Integral of a Gaussian

$$\int_{-\infty}^{\infty} e^{-ay^2} dy$$

**Trick No 1:** Squaring and taking the root doesn't change anything!

# Integral of a Gaussian

$$\int_{-\infty}^{\infty} e^{-ay^2} dy = \sqrt{\int_{-\infty}^{\infty} e^{-ay^2} dy \int_{-\infty}^{\infty} e^{-ay^2} dy}$$

**Trick No 1:** Squaring and taking the root doesn't change anything!

# Integral of a Gaussian

$$\begin{aligned}\int_{-\infty}^{\infty} e^{-ay^2} dy &= \sqrt{\int_{-\infty}^{\infty} e^{-ay^2} dy \int_{-\infty}^{\infty} e^{-ay^2} dy} \\ &= \sqrt{\int_{-\infty}^{\infty} e^{-ax^2} dx \int_{-\infty}^{\infty} e^{-ay^2} dy}\end{aligned}$$

**Trick No 1:** Squaring and taking the root doesn't change anything!

# Integral of a Gaussian

$$\begin{aligned}\int_{-\infty}^{\infty} e^{-ay^2} dy &= \sqrt{\int_{-\infty}^{\infty} e^{-ay^2} dy \int_{-\infty}^{\infty} e^{-ay^2} dy} \\ &= \sqrt{\int_{-\infty}^{\infty} e^{-ax^2} dx \int_{-\infty}^{\infty} e^{-ay^2} dy} \\ &= \sqrt{\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-a(x^2+y^2)} dx dy}\end{aligned}$$

**Trick No 1:** Squaring and taking the root doesn't change anything!



# Integral of a Gaussian

$$\begin{aligned}\int_{-\infty}^{\infty} e^{-ay^2} dy &= \sqrt{\int_{-\infty}^{\infty} e^{-ay^2} dy \int_{-\infty}^{\infty} e^{-ay^2} dy} \\ &= \sqrt{\int_{-\infty}^{\infty} e^{-ax^2} dx \int_{-\infty}^{\infty} e^{-ay^2} dy} \\ &= \sqrt{\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-a(x^2+y^2)} dx dy}\end{aligned}$$

**Trick No 1:** Squaring and taking the root doesn't change anything!

**Trick No 2:**  
use polar coordinates!

# Integral of a Gaussian

$$\begin{aligned}\int_{-\infty}^{\infty} e^{-ay^2} dy &= \sqrt{\int_{-\infty}^{\infty} e^{-ay^2} dy \int_{-\infty}^{\infty} e^{-ay^2} dy} \\&= \sqrt{\int_{-\infty}^{\infty} e^{-ax^2} dx \int_{-\infty}^{\infty} e^{-ay^2} dy} \\&= \sqrt{\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-a(x^2+y^2)} dx dy} \\&= \sqrt{\int_0^{2\pi} \int_0^{\infty} e^{-ar^2} r dr d\theta}\end{aligned}$$

**Trick No 1:** Squaring and taking the root doesn't change anything!

**Trick No 2:**  
use polar coordinates!

# Integral of a Gaussian

$$\begin{aligned}\int_{-\infty}^{\infty} e^{-ay^2} dy &= \sqrt{\int_{-\infty}^{\infty} e^{-ay^2} dy \int_{-\infty}^{\infty} e^{-ay^2} dy} \\&= \sqrt{\int_{-\infty}^{\infty} e^{-ax^2} dx \int_{-\infty}^{\infty} e^{-ay^2} dy} \\&= \sqrt{\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-a(x^2+y^2)} dx dy} \\&= \sqrt{\int_0^{2\pi} \int_0^{\infty} e^{-ar^2} r dr d\theta} \\&= \sqrt{2\pi \int_0^{\infty} e^{-ar^2} r dr}\end{aligned}$$

**Trick No 1:** Squaring and taking the root doesn't change anything!

**Trick No 2:**  
use polar coordinates!

# Integral of a Gaussian

$$\begin{aligned}\int_{-\infty}^{\infty} e^{-ay^2} dy &= \sqrt{\int_{-\infty}^{\infty} e^{-ay^2} dy \int_{-\infty}^{\infty} e^{-ay^2} dy} \\&= \sqrt{\int_{-\infty}^{\infty} e^{-ax^2} dx \int_{-\infty}^{\infty} e^{-ay^2} dy} \\&= \sqrt{\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-a(x^2+y^2)} dx dy} \\&= \sqrt{\int_0^{2\pi} \int_0^{\infty} e^{-ar^2} r dr d\theta} \\&= \sqrt{2\pi \int_0^{\infty} e^{-ar^2} r dr}\end{aligned}$$

**Trick No 1:** Squaring and taking the root doesn't change anything!

**Trick No 2:**  
use polar coordinates!

$$\frac{d}{dr} \frac{-1}{2a} e^{-ar^2} = \frac{-1}{2a} e^{-ar^2} (-2ar) = r e^{-ar^2}$$

# Integral of a Gaussian

$$\begin{aligned}\int_{-\infty}^{\infty} e^{-ay^2} dy &= \sqrt{\int_{-\infty}^{\infty} e^{-ay^2} dy \int_{-\infty}^{\infty} e^{-ay^2} dy} \\&= \sqrt{\int_{-\infty}^{\infty} e^{-ax^2} dx \int_{-\infty}^{\infty} e^{-ay^2} dy} \\&= \sqrt{\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-a(x^2+y^2)} dx dy} \\&= \sqrt{\int_0^{2\pi} \int_0^{\infty} e^{-ar^2} r dr d\theta} \\&= \sqrt{2\pi \int_0^{\infty} e^{-ar^2} r dr} \\&= \sqrt{2\pi \left[ -\frac{1}{2a} e^{-ar^2} \right]_0^{\infty}}\end{aligned}$$

**Trick No 1:** Squaring and taking the root doesn't change anything!

**Trick No 2:**  
use polar coordinates!

$$\frac{d}{dr} \frac{-1}{2a} e^{-ar^2} = \frac{-1}{2a} e^{-ar^2} (-2ar) = r e^{-ar^2}$$

# Integral of a Gaussian

$$\begin{aligned}\int_{-\infty}^{\infty} e^{-ay^2} dy &= \sqrt{\int_{-\infty}^{\infty} e^{-ay^2} dy \int_{-\infty}^{\infty} e^{-ay^2} dy} \\&= \sqrt{\int_{-\infty}^{\infty} e^{-ax^2} dx \int_{-\infty}^{\infty} e^{-ay^2} dy} \\&= \sqrt{\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-a(x^2+y^2)} dx dy} \\&= \sqrt{\int_0^{2\pi} \int_0^{\infty} e^{-ar^2} r dr d\theta} \\&= \sqrt{2\pi \int_0^{\infty} e^{-ar^2} r dr} \\&= \sqrt{2\pi \left[ -\frac{1}{2a} e^{-ar^2} \right]_0^{\infty}} \\&= \sqrt{\frac{\pi}{a}}\end{aligned}$$

**Trick No 1:** Squaring and taking the root doesn't change anything!

**Trick No 2:**  
use polar coordinates!

$$\frac{d}{dr} \frac{-1}{2a} e^{-ar^2} = \frac{-1}{2a} e^{-ar^2} (-2ar) = r e^{-ar^2}$$

# Naive DFT implementation

$$F_m = \sum_{n=0}^{M-1} f_n e^{-i2\pi \frac{mn}{M}}$$

# Naive DFT implementation

$$F_m = \sum_{n=0}^{M-1} f_n e^{-i2\pi \frac{mn}{M}}$$

$$F_m = e^{-i2\pi \frac{m0}{M}} f_0 + e^{-i2\pi \frac{m1}{M}} f_1 + \dots e^{-i2\pi \frac{m(M-1)}{M}} f_{M-1}$$



# Naive DFT implementation

$$F_m = \sum_{n=0}^{M-1} f_n e^{-i2\pi \frac{mn}{M}}$$

$$F_m = e^{-i2\pi \frac{m0}{M}} f_0 + e^{-i2\pi \frac{m1}{M}} f_1 + \dots e^{-i2\pi \frac{m(M-1)}{M}} f_{M-1}$$

$$\begin{pmatrix} F_0 \\ F_1 \\ \vdots \\ F_{M-1} \end{pmatrix} = \begin{pmatrix} e^{-i2\pi \frac{0*0}{M}} & e^{-i2\pi \frac{0*1}{M}} & \dots & e^{-i2\pi \frac{0(M-1)}{M}} \\ e^{-i2\pi \frac{1*0}{M}} & e^{-i2\pi \frac{1*1}{M}} & \dots & e^{-i2\pi \frac{1(M-1)}{M}} \\ \vdots & \vdots & \vdots & \vdots \\ e^{-i2\pi \frac{(M-1)0}{M}} & e^{-i2\pi \frac{(M-1)1}{M}} & \dots & e^{-i2\pi \frac{(M-1)(M-1)}{M}} \end{pmatrix} \begin{pmatrix} f_0 \\ f_1 \\ \vdots \\ f_{M-1} \end{pmatrix}$$

# DFT in Matlab

- Commands `fft2`, `ifft2`, `fftshift`, `ifftshift`
- Result of `fft2` has center of frequency domain (i.e., zero-frequency, or DC component) at top left
- `fftshift` shifts center of frequency domain to center of matrix
  - `ifftshift` performs inverse shift
- Use `fftshift` for visualization

# DFT filtering

1. DFT with zero-padding using  $\text{fft2}$ 
  - DFT filtering corresponds to circular convolution
  - Padding to avoid boundary artifacts
  - Padding to twice the image size is always safe
2. Generate filter function
3. Multiply Fourier transforms
4. Apply inverse DFT using  $\text{ifft2}$
5. Crop to original size

# Linear filtering

- Matlab functions
  - `fspecial` creates a 2-d filter of a specified type and size.
  - `imfilter` filters an image with a filter created using `fspecial`.

# Generating filter functions

- Generate a frequency domain filter from a spatial filter using `freqz2`
- Directly construct filter in frequency domain

# Meshgrid

- Use `meshgrid` to generate distance matrices

```
>> [x y] = meshgrid(-1:1,-1:1)
```

```
x =
```

-1	0	1
-1	0	1
-1	0	1

```
y =
```

-1	-1	-1
0	0	0
1	1	1

```
>> d = x.^2+y.^2
```

```
d =
```

2	1	2
1	0	1
2	1	2

```
>> 1./(1+d)
```

```
ans =
```

0.3333	0.5000	0.3333
0.5000	1.0000	0.5000
0.3333	0.5000	0.3333