

On a new method towards proof of Riemann's Hypothesis

Akhila Raman

Email: akhila.raman@berkeley.edu.

Abstract

We consider the analytic continuation of Riemann's Zeta Function derived from **Riemann's Xi function** $\xi(s)$ which is evaluated at $s = \frac{1}{2} + \sigma + i\omega$, given by $\xi(\frac{1}{2} + \sigma + i\omega) = E_{p\omega}(\omega)$, where σ, ω are real and $-\infty \leq \omega \leq \infty$ and compute its inverse Fourier transform given by $E_p(t)$.

We use a new method and show that the Fourier Transform of $E_p(t)$ given by $E_{p\omega}(\omega) = \xi(\frac{1}{2} + \sigma + i\omega)$ **does not have zeros** for finite and real ω when $0 < |\sigma| < \frac{1}{2}$, corresponding to the critical strip **excluding** the critical line and prove Riemann's hypothesis.

More importantly, the new method **does not** contradict the existence of non-trivial zeros on the critical line with real part of $s = \frac{1}{2}$ and **does not** contradict Riemann Hypothesis. It is shown that the new method is **not** applicable to Hurwitz zeta function and related functions and **does not** contradict the existence of their non-trivial zeros away from the critical line.

If the specific solution presented in this paper is incorrect, it is **hoped** that the new method discussed in this paper will lead to a correct solution by other researchers.

Keywords: Riemann, Hypothesis, Zeta, Xi, exponential functions

1. Introduction

It is well known that Riemann's Zeta function given by $\zeta(s) = \sum_{m=1}^{\infty} \frac{1}{m^s}$ converges in the half-plane where the real part of s is greater than 1. Riemann proved that $\zeta(s)$ has an analytic continuation to the whole s -plane apart from a simple pole at $s = 1$ and that $\zeta(s)$ satisfies a symmetric functional equation given by $\xi(s) = \xi(1-s) = \frac{1}{2}s(s-1)\pi^{-\frac{s}{2}}\Gamma(\frac{s}{2})\zeta(s)$ where $\Gamma(s) = \int_0^{\infty} e^{-u}u^{s-1}du$ is the Gamma function.^{[4] [5]} We can see that if Riemann's Xi function has a zero in the critical strip, then Riemann's Zeta function also has a zero at the same location. Riemann made his conjecture in his 1859 paper, that all of the non-trivial zeros of $\zeta(s)$ lie on the critical line with real part of $s = \frac{1}{2}$, which is called the Riemann Hypothesis.^[1]

Hardy and Littlewood later proved that infinitely many of the zeros of $\zeta(s)$ are on the critical line with real part of $s = \frac{1}{2}$.^[2] It is well known that $\zeta(s)$ does not have non-trivial zeros when real part of $s = \frac{1}{2} + \sigma + i\omega$, given by $\frac{1}{2} + \sigma \geq 1$ and $\frac{1}{2} + \sigma \leq 0$. In this paper, **critical strip** $0 < \text{Re}[s] < 1$ corresponds to $0 \leq |\sigma| < \frac{1}{2}$.

In this paper, a **new method** is discussed and a specific solution is presented to prove Riemann's Hypothesis. If the specific solution presented in this paper is incorrect, it is **hoped** that the new

method discussed in this paper will lead to a correct solution by other researchers.

In Section 2 to Section 6, we prove Riemann's hypothesis by taking the analytic continuation of Riemann's Zeta Function derived from Riemann's Xi function $\xi(\frac{1}{2} + \sigma + i\omega) = E_{p\omega}(\omega)$ and compute inverse Fourier transform of $E_{p\omega}(\omega)$ given by $E_p(t)$ and show that its Fourier transform $E_{p\omega}(\omega)$ does not have zeros for finite and real ω when $0 < |\sigma| < \frac{1}{2}$, corresponding to the critical strip **excluding** the critical line.

In Section 7, it is shown that the new method is **not** applicable to Hurwitz zeta function and related functions and **does not** contradict the existence of their non-trivial zeros away from the critical line with real part of $s = \frac{1}{2}$.

We present an **outline** of the new method below.

1.1. Step 1: Inverse Fourier Transform of $\xi(\frac{1}{2} + i\omega)$

Let us start with Riemann's Xi Function $\xi(s)$ evaluated at $s = \frac{1}{2} + i\omega$ given by $\xi(\frac{1}{2} + i\omega) = \Xi(\omega) = E_{0\omega}(\omega)$, where $-\infty \leq \omega \leq \infty$. Its inverse Fourier Transform is given by $E_0(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} E_{0\omega}(\omega) e^{i\omega t} d\omega$, where ω, t are real, as follows (link).^[3]

$$E_0(t) = \Phi(t) = 2 \sum_{n=1}^{\infty} [2n^4 \pi^2 e^{\frac{9t}{2}} - 3n^2 \pi e^{\frac{5t}{2}}] e^{-\pi n^2 e^{2t}} = \sum_{n=1}^{\infty} [4\pi^2 n^4 e^{4t} - 6\pi n^2 e^{2t}] e^{-\pi n^2 e^{2t}} e^{\frac{t}{2}} \quad (1)$$

We see that $E_0(t) = E_0(-t)$ is a real and **even** function of t , given that $E_{0\omega}(\omega) = E_{0\omega}(-\omega)$ because $\xi(s) = \xi(1-s)$ and hence $\xi(\frac{1}{2} + i\omega) = \xi(\frac{1}{2} - i\omega)$ when evaluated at $s = \frac{1}{2} + i\omega$.

The inverse Fourier Transform of $\xi(\frac{1}{2} + \sigma + i\omega) = E_{p\omega}(\omega)$ is given by the real function $E_p(t)$. We can write $E_p(t)$ as follows for $0 < |\sigma| < \frac{1}{2}$ and this is shown in detail in Appendix A.

$$E_p(t) = E_0(t) e^{-\sigma t} = \sum_{n=1}^{\infty} [4\pi^2 n^4 e^{4t} - 6\pi n^2 e^{2t}] e^{-\pi n^2 e^{2t}} e^{\frac{t}{2}} e^{-\sigma t} \quad (2)$$

We can see that $E_p(t)$ is an analytic function in the interval $|t| \leq \infty$, given that the sum and product of exponential functions are analytic in the same interval and hence infinitely differentiable in that interval.

1.2. Step 2: On the zeros of a related function $G(\omega, t_2, t_0)$

Statement 1: Let us assume that Riemann's Xi function $\xi(\frac{1}{2} + \sigma + i\omega) = E_{p\omega}(\omega)$ has a zero at $\omega = \omega_0$ where ω_0 is real and finite and $0 < |\sigma| < \frac{1}{2}$, corresponding to the critical strip excluding the critical line. We will prove that this assumption leads to a **contradiction**.

Let us consider $0 < \sigma < \frac{1}{2}$ at first. Let us consider a new function $g(t, t_2, t_0) = f(t, t_2, t_0) e^{-\sigma t} u(-t) + f(t, t_2, t_0) e^{\sigma t} u(t)$, where $f(t, t_2, t_0) = e^{-2\sigma t_0} f_1(t, t_2, t_0) + e^{2\sigma t_0} f_2(t, t_2, t_0)$ and $f_1(t, t_2, t_0) = e^{\sigma t_0} E'_p(t + t_0, t_2)$ and $f_2(t, t_2, t_0) = e^{-\sigma t_0} E'_p(t - t_0, t_2)$ and $E'_p(t, t_2) = e^{-\sigma t_2} E_p(t - t_2) - e^{\sigma t_2} E_p(t + t_2)$ and t_0, t_2 are real and $g(t, t_2, t_0)$ is a real function of variable t and $u(t)$ is Heaviside unit step function. We can

73 see that $g(t, t_2, t_0)h(t) = f(t, t_2, t_0)$ where $h(t) = [e^{\sigma t}u(-t) + e^{-\sigma t}u(t)]$.

74
75 In Section 2.1, we will show that the Fourier transform of the **even function** $g_{even}(t, t_2, t_0) =$
76 $\frac{1}{2}[g(t, t_2, t_0) + g(-t, t_2, t_0)]$ given by $G_R(\omega, t_0, t_2)$ must have **at least one zero** at $\omega = \omega_z(t_2, t_0) \neq 0$,
77 for every value of t_0 , for a given value of t_2 , where $G_R(\omega, t_0, t_2)$ crosses the zero line to the opposite
78 sign, to satisfy Statement 1, where $\omega_z(t_2, t_0)$ is real and finite.

79 1.3. *Step 3: On the zeros of the function $G_R(\omega, t_0, t_2)$*

80
81 In Section 2.2, we compute the Fourier transform of the function $g(t, t_2, t_0)$ and compute its real
82 part given by $G_R(\omega, t_2, t_0)$ and we can write as follows.

$$\begin{aligned} G_R(\omega, t_2, t_0) = & e^{-2\sigma t_0} \int_{-\infty}^0 [E'_0(\tau + t_0, t_2)e^{-2\sigma\tau} + E'_{0n}(\tau - t_0, t_2)] \cos(\omega\tau) d\tau \\ & + e^{2\sigma t_0} \int_{-\infty}^0 [E'_0(\tau - t_0, t_2)e^{-2\sigma\tau} + E'_{0n}(\tau + t_0, t_2)] \cos(\omega\tau) d\tau \end{aligned}$$

(3)

84 We require $G_R(\omega, t_2, t_0) = 0$ for $\omega = \omega_z(t_2, t_0)$ for every value of t_0 , for **each fixed value**
85 of t_2 , to satisfy **Statement 1**. In general $\omega_z(t_2, t_0) \neq \omega_0$. Hence we can see that $P(t_2, t_0) =$
86 $G_R(\omega_z(t_2, t_0), t_2, t_0) = 0$.

87 1.4. *Step 4: Zero Crossing function $\omega_z(t_2, t_0)$ is an even function of variable t_0*

88
89 In Section 2.3, we show the result in Eq. 4 and that $\omega_z(t_2, t_0) = \omega_z(t_2, -t_0)$. It is shown that
90 $P(t_2, t_0) = G_R(\omega_z(t_2, t_0), t_2, t_0) = P_{odd}(t_2, t_0) + P_{odd}(t_2, -t_0) = 0$ and that $P_{odd}(t_2, t_0)$ is an **odd**
91 function of t_0 , for a given value of t_2 as follows.

$$\begin{aligned} P_{odd}(t_2, t_0) = & [\cos(\omega_z(t_2, t_0)t_0) \int_{-\infty}^{t_0} E'_0(\tau, t_2)e^{-2\sigma\tau} \cos(\omega_z(t_2, t_0)\tau) d\tau \\ & + \sin(\omega_z(t_2, t_0)t_0) \int_{-\infty}^{t_0} E'_0(\tau, t_2)e^{-2\sigma\tau} \sin(\omega_z(t_2, t_0)\tau) d\tau] \\ & + e^{2\sigma t_0} [\cos(\omega_z(t_2, t_0)t_0) \int_{-\infty}^{t_0} E'_{0n}(\tau, t_2) \cos(\omega_z(t_2, t_0)\tau) d\tau + \sin(\omega_z(t_2, t_0)t_0) \int_{-\infty}^{t_0} E'_{0n}(\tau, t_2) \sin(\omega_z(t_2, t_0)\tau) d\tau] \end{aligned}$$

(4)

93 1.5. *Step 5: Final Step*

94
95 In Section 4, it is shown that $\omega_z(t_2, t_0)$ is a **continuous** function of variable t_0 and t_2 , for all
96 $|t_0| < \infty$ and $|t_2| < \infty$. In Section 6, it is shown that $E_0(t)$ is **strictly decreasing** for $t > 0$.

97
98 In Section 3, we set $t_0 = t_{0c}$ and $t_2 = t_{2c} = 2t_{0c}$, such that $\omega_z(t_{2c}, t_{0c})t_{0c} = \frac{\pi}{2}$ and substitute
99 in the equation for $P_{odd}(t_2, t_0)$ in Eq. 4 and show that this leads to the result in Eq. 5. We use
100 $E'_0(t, t_2) = E_0(t - t_2) - E_0(t + t_2)$ and $E'_{0n}(t, t_2) = E'_0(-t, t_2)$.

$$\int_0^{t_{0c}} (E_0(\tau - t_{2c}) - E_0(\tau + t_{2c}))(\cosh(2\sigma t_{0c}) - \cosh(2\sigma\tau)) \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau = 0$$

(5)

We show that the **each** of the terms in the integrand in Eq. 5 are **greater than zero**, in the interval $0 < \tau < t_{0c}$ and the integrand is zero at $\tau = 0$ and $\tau = t_{0c}$, where $t_{0c} > 0$.

Hence the result in Eq. 5 leads to a **contradiction** for $0 < \sigma < \frac{1}{2}$.

We have shown this result for $0 < \sigma < \frac{1}{2}$ and then use the property $\xi(\frac{1}{2} + \sigma + i\omega) = \xi(\frac{1}{2} - \sigma - i\omega)$ to show the result for $-\frac{1}{2} < \sigma < 0$. Hence we have produced a **contradiction** of **Statement 1** that the Fourier Transform of the function $E_p(t) = E_0(t)e^{-\sigma t}$ has a zero at $\omega = \omega_0$ for $0 < |\sigma| < \frac{1}{2}$.

2. An Approach towards Riemann's Hypothesis

Theorem 1: Riemann's Xi function $\xi(\frac{1}{2} + \sigma + i\omega) = E_{p\omega}(\omega)$ does not have zeros for any real value of $-\infty < \omega < \infty$, for $0 < |\sigma| < \frac{1}{2}$, corresponding to the critical strip excluding the critical line, given that $E_0(t) = E_0(-t)$ is an even function of variable t , where $E_p(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} E_{p\omega}(\omega) e^{i\omega t} d\omega$, $E_p(t) = E_0(t)e^{-\sigma t}$ and $E_0(t) = \sum_{n=1}^{\infty} [4\pi^2 n^4 e^{4t} - 6\pi n^2 e^{2t}] e^{-\pi n^2 e^{2t}} e^{\frac{t}{2}}$.

Proof: We assume that Riemann Hypothesis is false and prove its truth using proof by contradiction.

Statement 1: Let us assume that Riemann's Xi function $\xi(\frac{1}{2} + \sigma + i\omega) = E_{p\omega}(\omega)$ has a zero at $\omega = \omega_0$ where ω_0 is real and finite and $0 < |\sigma| < \frac{1}{2}$, corresponding to the critical strip excluding the critical line. We will prove that this assumption leads to a **contradiction**.

We will prove it for $0 < \sigma < \frac{1}{2}$ first and then use the property $\xi(\frac{1}{2} + \sigma + i\omega) = \xi(\frac{1}{2} - \sigma - i\omega)$ to show the result for $-\frac{1}{2} < \sigma < 0$ and hence show the result for $0 < |\sigma| < \frac{1}{2}$. We use the fact that $E_p(t)$ is real and its Fourier transform $E_{p\omega}(\omega)$ has symmetry property. (link).

We know that $\omega_0 \neq 0$, because $\zeta(s)$ has no zeros on the real axis between 0 and 1, when $s = \frac{1}{2} + \sigma + i\omega$ is real, $\omega = 0$ and $0 < |\sigma| < \frac{1}{2}$. [3] This is shown in detail in first two paragraphs in Appendix B.1.

2.1. New function $g(t, t_2, t_0)$

Let us consider the function $E'_p(t, t_2) = e^{-\sigma t_2} E_p(t - t_2) - e^{\sigma t_2} E_p(t + t_2) = (E_0(t - t_2) - E_0(t + t_2))e^{-\sigma t} = E'_0(t, t_2)e^{-\sigma t}$, where t_2 is finite and real, and $E'_0(t, t_2) = E_0(t - t_2) - E_0(t + t_2)$. Its Fourier transform is given by $E'_{p\omega}(\omega, t_2) = E_{p\omega}(\omega)(e^{-\sigma t_2} e^{-i\omega t_2} - e^{\sigma t_2} e^{i\omega t_2})$ which has a zero at the **same** $\omega = \omega_0$.

Let us consider the function $f(t, t_2, t_0) = e^{-2\sigma t_0} f_1(t, t_2, t_0) + e^{2\sigma t_0} f_2(t, t_2, t_0)$ where $f_1(t, t_2, t_0) = e^{\sigma t_0} E'_p(t + t_0, t_2)$ and $f_2(t, t_2, t_0) = e^{-\sigma t_0} E'_p(t - t_0, t_2)$ where t_0 is finite and real and we can see that the Fourier Transform of this function $F(\omega, t_2, t_0) = E'_{p\omega}(\omega, t_2)(e^{-\sigma t_0} e^{i\omega t_0} + e^{\sigma t_0} e^{-i\omega t_0})$ also has a zero

at the **same** $\omega = \omega_0$.

Let us consider a new function $g(t, t_2, t_0) = g_-(t, t_2, t_0)u(-t) + g_+(t, t_2, t_0)u(t)$ where $g(t, t_2, t_0)$ is a real function of variable t and $u(t)$ is Heaviside unit step function and $g_-(t, t_2, t_0) = f(t, t_2, t_0)e^{-\sigma t}$ and $g_+(t, t_2, t_0) = f(t, t_2, t_0)e^{\sigma t}$. We can see that $g(t, t_2, t_0)h(t) = f(t, t_2, t_0)$ where $h(t) = [e^{\sigma t}u(-t) + e^{-\sigma t}u(t)]$.

We can show that $E_p(t), E'_p(t, t_2), h(t)$ are real absolutely integrable functions and go to zero as $t \rightarrow \pm\infty$. Hence their respective Fourier transforms given by $E_{p\omega}(\omega), E'_{p\omega}(\omega, t_2), H(\omega)$ are finite for $|\omega| \leq \infty$ and go to zero as $|\omega| \rightarrow \infty$, as per Riemann Lebesgue Lemma (link). We can show that $E_0(t)$ and $E_0(t)e^{-2\sigma t}$ are absolutely **integrable** functions. This is shown in detail in Appendix B.1. We see that $g(t, t_2, t_0)$ is a Fourier transformable function. (Section 2.2 and Section 2.3)

If we take the Fourier transform of the equation $g(t, t_2, t_0)h(t) = f(t, t_2, t_0)$ where $h(t) = [e^{\sigma t}u(-t) + e^{-\sigma t}u(t)]$, we get $\frac{1}{2\pi}[G(\omega, t_2, t_0)*H(\omega)] = F(\omega, t_2, t_0) = E'_{p\omega}(\omega, t_2)(e^{-\sigma t_0}e^{i\omega t_0} + e^{\sigma t_0}e^{-i\omega t_0}) = F_R(\omega, t_2, t_0) + iF_I(\omega, t_2, t_0)$ as per convolution theorem (link), where $*$ denotes convolution operation given by $F(\omega, t_2, t_0) = \frac{1}{2\pi} \int_{-\infty}^{\infty} G(\omega', t_2, t_0)H(\omega - \omega')d\omega'$ and $H(\omega) = H_R(\omega) = [\frac{1}{\sigma - i\omega} + \frac{1}{\sigma + i\omega}] = \frac{2\sigma}{(\sigma^2 + \omega^2)}$ is real and is the Fourier transform of the function $h(t)$ and $G(\omega, t_2, t_0) = G_R(\omega, t_0, t_2) + iG_I(\omega, t_2, t_0)$ is the Fourier transform of the function $g(t, t_2, t_0)$. We can write $g(t, t_2, t_0) = g_{\text{even}}(t, t_2, t_0) + g_{\text{odd}}(t, t_2, t_0)$ where $g_{\text{even}}(t, t_2, t_0)$ is an even function and $g_{\text{odd}}(t, t_2, t_0)$ is an odd function of variable t .

If Statement 1 is true, then we require the Fourier transform of the function $f(t, t_2, t_0)$ given by $F(\omega, t_2, t_0)$ to have a zero at $\omega = \omega_0$ for **every value** of t_0 , for a given fixed value of t_2 . This implies that the **real** part of the Fourier transform of the **even function** $g_{\text{even}}(t, t_2, t_0) = \frac{1}{2}[g(t, t_2, t_0) + g(-t, t_2, t_0)]$ given by $G_R(\omega, t_0, t_2)$ must have **at least one zero** at $\omega = \omega_z(t_2, t_0) \neq 0$ where $\omega_z(t_2, t_0)$ is real and finite, where $G_R(\omega, t_0, t_2)$ crosses the zero line to the opposite sign. We call this **Statement 2**. We note that $\omega_z(t_2, t_0)$ can be different from ω_0 in general.

Because $H(\omega) = \frac{2\sigma}{(\sigma^2 + \omega^2)}$ is real and does not have zeros for any finite value of ω , **if** $G_R(\omega, t_0, t_2)$ does not have at least one zero for some $\omega = \omega_z(t_2, t_0) \neq 0$, where $G_R(\omega, t_0, t_2)$ crosses the zero line to the opposite sign, **then** the **real part** of $F(\omega, t_2, t_0)$ given by $F_R(\omega, t_2, t_0) = \frac{1}{2\pi}[G_R(\omega, t_0, t_2) * H(\omega)]$, obtained by the convolution of $H(\omega)$ and $G_R(\omega, t_0, t_2)$, **cannot** possibly have zeros for any non-zero finite value of ω , which goes against **Statement 1**. This is shown in detail in Lemma 1.

Lemma 1: If Riemann's Xi function $\xi(\frac{1}{2} + \sigma + i\omega) = E_{p\omega}(\omega)$ has a zero at $\omega = \omega_0 \neq 0$ where ω_0 is real and finite, then the **real** part of the Fourier transform of the **even function** $g_{\text{even}}(t, t_2, t_0) = \frac{1}{2}[g(t, t_2, t_0) + g(-t, t_2, t_0)]$ given by $G_R(\omega, t_0, t_2)$ must have **at least one zero** at $\omega = \omega_z(t_2, t_0) \neq 0$ for **every value** of t_0 , for a given finite value of t_2 , where $G_R(\omega, t_0, t_2)$ crosses the zero line to the opposite sign and $\omega_z(t_2, t_0)$ is real and finite, where $g(t, t_2, t_0)h(t) = f(t, t_2, t_0) = e^{-2\sigma t_0}f_1(t, t_2, t_0) + e^{2\sigma t_0}f_2(t, t_2, t_0)$ where $f_1(t, t_2, t_0) = e^{\sigma t_0}E'_p(t + t_0, t_2)$ and $f_2(t, t_2, t_0) = e^{-\sigma t_0}E'_p(t - t_0, t_2)$, $E'_p(t, t_2) = e^{-\sigma t_2}E_p(t - t_2) - e^{\sigma t_2}E_p(t + t_2)$, and $h(t) = e^{\sigma t}u(-t) + e^{-\sigma t}u(t)$ and $0 < \sigma < \frac{1}{2}$.

Proof: If $E_{p\omega}(\omega)$ has a zero at finite $\omega = \omega_0 \neq 0$ to satisfy Statement 1, then $F(\omega, t_2, t_0) = E'_{p\omega}(\omega, t_2)(e^{-\sigma t_0}e^{i\omega t_0} + e^{\sigma t_0}e^{-i\omega t_0}) = E_{p\omega}(\omega)(e^{-\sigma t_2}e^{-i\omega t_2} - e^{\sigma t_2}e^{i\omega t_2})(e^{-\sigma t_0}e^{i\omega t_0} + e^{\sigma t_0}e^{-i\omega t_0})$ also has a zero at $\omega = \omega_0$ and its real part given by $F_R(\omega, t_2, t_0)$ also has a zero at the same location $\omega = \omega_0 \neq 0$.

Let us consider the case where $G_R(\omega, t_2, t_0)$ **does not** have at least one zero for finite $\omega = \omega_z(t_2, t_0) \neq 0$, where $G_R(\omega, t_0, t_2)$ crosses the zero line to the opposite sign and show that $F_R(\omega, t_2, t_0)$ does not have at least one zero at finite $\omega \neq 0$ for this case, which **contradicts** Statement 1. Given that $H(\omega)$ is real, we can write the convolution theorem only for the real parts as follows.

$$F_R(\omega, t_2, t_0) = \frac{1}{2\pi} \int_{-\infty}^{\infty} G_R(\omega', t_2, t_0) H(\omega - \omega') d\omega' \quad (6)$$

We can show that the above integral converges for all $|\omega| \leq \infty$, given that $G(\omega, t_2, t_0)$ and $H(\omega)$ have fall-off rate of $\frac{1}{\omega^2}$ as $|\omega| \rightarrow \infty$ because the first derivatives of $g(t, t_2, t_0)$ and $h(t)$ are discontinuous at $t = 0$. (Appendix B.2)

We substitute $H(\omega) = \frac{2\sigma}{(\sigma^2 + \omega^2)}$ in Eq. 6 and we get

$$F_R(\omega, t_2, t_0) = \frac{\sigma}{\pi} \int_{-\infty}^{\infty} G_R(\omega', t_2, t_0) \frac{1}{(\sigma^2 + (\omega - \omega')^2)} d\omega' \quad (7)$$

We can split the integral in Eq. 7 as follows.

$$F_R(\omega, t_2, t_0) = \frac{\sigma}{\pi} \left[\int_{-\infty}^0 G_R(\omega', t_2, t_0) \frac{1}{(\sigma^2 + (\omega - \omega')^2)} d\omega' + \int_0^{\infty} G_R(\omega', t_2, t_0) \frac{1}{(\sigma^2 + (\omega - \omega')^2)} d\omega' \right] \quad (8)$$

We see that $G_R(-\omega, t_2, t_0) = G_R(\omega, t_2, t_0)$ because $g(t, t_2, t_0)$ is a real function of variable t . (link) We can substitute $\omega' = -\omega''$ in the first integral in Eq. 8 and substituting $\omega'' = \omega'$ in the result, we can write as follows.

$$F_R(\omega, t_2, t_0) = \frac{\sigma}{\pi} \int_0^{\infty} G_R(\omega', t_2, t_0) \left[\frac{1}{(\sigma^2 + (\omega - \omega')^2)} + \frac{1}{(\sigma^2 + (\omega + \omega')^2)} \right] d\omega' \quad (9)$$

In Appendix B.2, it is shown that $G(\omega, t_2, t_0)$ is finite for $|\omega| \leq \infty$ and goes to zero as $|\omega| \rightarrow \infty$. We can see that for $\omega' \rightarrow \infty$, the integrand in Eq. 9 is zero. For finite $\omega \geq 0$, and $0 \leq \omega' < \infty$, we can see that the term $\frac{1}{(\sigma^2 + (\omega - \omega')^2)} + \frac{1}{(\sigma^2 + (\omega + \omega')^2)} > 0$. We see that $G_R(\omega', t_0, t_2)$ is **not** an all zero function and that $G_R(\omega', t_2, t_0)$ is a continuous function of ω' , for a fixed t_0 and t_2 (Section 4.1).

• **Case 1:** $G_R(\omega', t_2, t_0) \geq 0$ for all finite $\omega' \geq 0$

We see that $F_R(\omega, t_2, t_0) > 0$ for all finite $\omega \geq 0$. We see that $F_R(-\omega, t_2, t_0) = F_R(\omega, t_2, t_0)$ because $f(t, t_2, t_0)$ is a real function (link). Hence $F_R(\omega, t_2, t_0) > 0$ for all finite $\omega \leq 0$.

This **contradicts** Statement 1 which requires $F_R(\omega, t_2, t_0)$ to have at least one zero at finite $\omega \neq 0$ because we showed that $\omega_0 \neq 0$ in **Section 2** paragraph 5. Therefore $G_R(\omega', t_2, t_0)$ must have **at least one zero** at $\omega' = \omega_z(t_2, t_0) \neq 0$ where it crosses the zero line and becomes negative, where $\omega_z(t_2, t_0)$ is real and finite.

• **Case 2:** $G_R(\omega', t_2, t_0) \leq 0$ for all finite $\omega' \geq 0$

220 We see that $F_R(\omega, t_2, t_0) < 0$ for all finite $\omega \geq 0$. We see that $F_R(-\omega, t_2, t_0) = F_R(\omega, t_2, t_0)$ because
 221 $f(t, t_2, t_0)$ is a real function (link). Hence $F_R(\omega, t_2, t_0) < 0$ for all finite $\omega \leq 0$.

222
 223 This **contradicts** Statement 1 which requires $F_R(\omega, t_2, t_0)$ to have at least one zero at finite $\omega \neq 0$.
 224 Therefore $G_R(\omega', t_2, t_0)$ must have **at least one zero** at $\omega' = \omega_z(t_2, t_0) \neq 0$, where it crosses the
 225 zero line and becomes positive, where $\omega_z(t_2, t_0)$ is real and finite.

226
 227 We have shown that, $G_R(\omega, t_2, t_0)$ must have **at least one zero** at finite $\omega = \omega_z(t_2, t_0) \neq 0$ where
 228 it crosses the zero line to the opposite sign, to satisfy **Statement 1**. We call this **Statement 2**. In
 229 the rest of the sections, we consider only the **first** zero crossing away from origin, where $G_R(\omega, t_0, t_2)$
 230 crosses the zero line to the opposite sign. Hence $0 < \omega_z(t_2, t_0) < \infty$, for all $|t_0| < \infty$, for every finite
 231 value of t_2 .

2.2. On the zeros of a related function $G(\omega, t_2, t_0)$

232
 233 We can compute the fourier transform of the function $g_{\text{even}}(t, t_2, t_0) = \frac{1}{2}[g(t, t_2, t_0) + g(-t, t_2, t_0)]$
 234 given by $G_R(\omega, t_2, t_0)$. We require $G_R(\omega, t_2, t_0) = 0$ for $\omega = \omega_z(t_2, t_0)$ for **every value** of t_0 , for a
 236 given value of t_2 , to satisfy **Statement 1**. In general, $\omega_z(t_2, t_0) \neq \omega_0$.

237
 238 First we compute the Fourier transform of the function $g_1(t, t_2, t_0)$ given by $G_1(\omega, t_2, t_0) = G_{1R}(\omega, t_2, t_0) +$
 239 $iG_{1I}(\omega, t_2, t_0)$. We use $g_1(t, t_2, t_0) = f_1(t, t_2, t_0)e^{-\sigma t}u(-t) + f_1(t, t_2, t_0)e^{\sigma t}u(t) = e^{\sigma t_0}E'_p(t+t_0, t_2)e^{-\sigma t}u(-t) +$
 240 $e^{\sigma t_0}E'_p(t+t_0, t_2)e^{\sigma t}u(t)$.

241

$$\begin{aligned} G_1(\omega, t_2, t_0) &= \int_{-\infty}^{\infty} g_1(t, t_2, t_0)e^{-i\omega t}dt = \int_{-\infty}^0 g_1(t, t_2, t_0)e^{-i\omega t}dt + \int_0^{\infty} g_1(t, t_2, t_0)e^{-i\omega t}dt \\ G_1(\omega, t_2, t_0) &= \int_{-\infty}^0 e^{\sigma t_0}E'_p(t+t_0, t_2)e^{-\sigma t}e^{-i\omega t}dt + \int_0^{\infty} e^{\sigma t_0}E'_p(t+t_0, t_2)e^{\sigma t}e^{-i\omega t}dt \end{aligned}$$

242

(10)

243 We use $E'_p(t, t_2) = E'_0(t, t_2)e^{-\sigma t}$ where $E'_0(t, t_2) = E_0(t - t_2) - E_0(t + t_2)$ and $E'_p(t + t_0, t_2) =$
 244 $E'_0(t + t_0, t_2)e^{-\sigma t}e^{-\sigma t_0}$. Substituting $t = -t$ in the second integral in Eq. 10, we have

$$\begin{aligned} G_1(\omega, t_2, t_0) &= \int_{-\infty}^0 E'_0(t+t_0, t_2)e^{-2\sigma t}e^{-i\omega t}dt + \int_0^{\infty} E'_0(t+t_0, t_2)e^{-i\omega t}dt \\ G_1(\omega, t_2, t_0) &= \int_{-\infty}^0 E'_0(t+t_0, t_2)e^{-2\sigma t}e^{-i\omega t}dt + \int_{-\infty}^0 E'_0(-t+t_0, t_2)e^{i\omega t}dt \end{aligned}$$

245

(11)

246 We define $E'_{0n}(t, t_2) = E'_0(-t, t_2)$ and get $E'_0(-t+t_0, t_2) = E'_{0n}(t-t_0, t_2)$ and write Eq. 11 as
 247 follows.

$$G_1(\omega, t_2, t_0) = \int_{-\infty}^0 E'_0(t+t_0, t_2)e^{-2\sigma t}e^{-i\omega t}dt + \int_{-\infty}^0 E'_{0n}(t-t_0, t_2)e^{i\omega t}dt = G_{1R}(\omega, t_2, t_0) + iG_{1I}(\omega, t_2, t_0)$$

$$(12)$$

The above equations can be expanded as follows using the identity $e^{i\omega t} = \cos(\omega t) + i \sin(\omega t)$.
Comparing the **real parts** of $G_1(\omega, t_2, t_0)$, we have

$$G_{1R}(\omega, t_2, t_0) = \int_{-\infty}^0 E'_0(t + t_0, t_2) e^{-2\sigma t} \cos(\omega t) dt + \int_{-\infty}^0 E'_{0n}(t - t_0, t_2) \cos(\omega t) dt$$

$$(13)$$

2.3. Zero crossing function $\omega_z(t_2, t_0)$ is an even function of variable t_0 , for a fixed t_2

Now we consider the function $f(t, t_2, t_0) = e^{-2\sigma t_0} f_1(t, t_2, t_0) + e^{2\sigma t_0} f_2(t, t_2, t_0) = e^{-\sigma t_0} E'_p(t + t_0, t_2) + e^{\sigma t_0} E'_p(t - t_0, t_2)$ where $f_1(t, t_2, t_0) = e^{\sigma t_0} E'_p(t + t_0, t_2)$ and $f_2(t, t_2, t_0) = f_1(t, t_2, -t_0) = e^{-\sigma t_0} E'_p(t - t_0, t_2)$ and $g(t, t_2, t_0)h(t) = f(t, t_2, t_0)$ where $g(t, t_2, t_0) = f(t, t_2, t_0)e^{-\sigma t}u(-t) + f(t, t_2, t_0)e^{\sigma t}u(t)$ and $h(t) = [e^{\sigma t}u(-t) + e^{-\sigma t}u(t)]$ and compute the Fourier transform of the function $g(t, t_2, t_0)$ and compute its real part $G_R(\omega, t_2, t_0)$ using the procedure in above section, similar to Eq. 13 and we can write as follows. We substitute $t = \tau$.

$$\begin{aligned} G_R(\omega, t_2, t_0) &= e^{-2\sigma t_0} G_{1R}(\omega, t_2, t_0) + e^{2\sigma t_0} G_{1R}(\omega, t_2, -t_0) \\ G_{1R}(\omega, t_2, t_0) &= \int_{-\infty}^0 [E'_0(\tau + t_0, t_2) e^{-2\sigma \tau} + E'_{0n}(\tau - t_0, t_2)] \cos(\omega \tau) d\tau \\ G_R(\omega, t_2, t_0) &= e^{-2\sigma t_0} \int_{-\infty}^0 [E'_0(\tau + t_0, t_2) e^{-2\sigma \tau} + E'_{0n}(\tau - t_0, t_2)] \cos(\omega \tau) d\tau \\ &\quad + e^{2\sigma t_0} \int_{-\infty}^0 [E'_0(\tau - t_0, t_2) e^{-2\sigma \tau} + E'_{0n}(\tau + t_0, t_2)] \cos(\omega \tau) d\tau \end{aligned}$$

$$(14)$$

We require $G_R(\omega, t_2, t_0) = 0$ for $\omega = \omega_z(t_2, t_0)$ for every value of t_0 , for **every finite value** of t_2 , to satisfy **Statement 1**. In general $\omega_z(t_2, t_0) \neq \omega_0$. Hence we can see that $P(t_2, t_0) = G_R(\omega_z(t_2, t_0), t_2, t_0) = 0$ and we can rearrange the terms as follows.

$$\begin{aligned} P(t_2, t_0) &= \int_{-\infty}^0 [e^{-2\sigma t_0} E'_0(\tau + t_0, t_2) e^{-2\sigma \tau} + e^{2\sigma t_0} E'_{0n}(\tau + t_0, t_2)] \cos(\omega_z(t_2, t_0) \tau) d\tau \\ &\quad + \int_{-\infty}^0 [e^{2\sigma t_0} E'_0(\tau - t_0, t_2) e^{-2\sigma \tau} + e^{-2\sigma t_0} E'_{0n}(\tau - t_0, t_2)] \cos(\omega_z(t_2, t_0) \tau) d\tau = 0 \end{aligned}$$

$$(15)$$

We can write as follows, where $P_{odd}(t_2, t_0)$ is an **odd** function of variable t_0 , for every finite value of t_2 .

$$\begin{aligned} P(t_2, t_0) &= P_{odd}(t_2, t_0) + P_{odd}(t_2, -t_0) = 0 \\ P_{odd}(t_2, t_0) &= \int_{-\infty}^0 [e^{-2\sigma t_0} E'_0(\tau + t_0, t_2) e^{-2\sigma \tau} + e^{2\sigma t_0} E'_{0n}(\tau + t_0, t_2)] \cos(\omega_z(t_2, t_0) \tau) d\tau \end{aligned}$$

We see that $f(t, t_2, t_0) = e^{-\sigma t_0} E'_p(t + t_0, t_2) + e^{\sigma t_0} E'_p(t - t_0, t_2) = f(t, t_2, -t_0)$ is **unchanged** by the substitution $t_0 = -t_0$ and hence $\omega_z(t_2, t_0)$ is an **even** function of variable t_0 , for **every finite value** of t_2 .

271

3. Final Step

273

We expand $P_{odd}(t_2, t_0)$ in Eq. 16 as follows, using the substitution $\tau + t_0 = \tau'$ and substituting back $\tau' = \tau$. We use $E'_{0n}(\tau, t_2) = E'_0(-\tau, t_2)$ and $E'_0(\tau, t_2) = E_0(\tau - t_2) - E_0(\tau + t_2)$.

275

$$\begin{aligned}
 P_{odd}(t_2, t_0) &= [\cos(\omega_z(t_2, t_0)t_0) \int_{-\infty}^{t_0} E'_0(\tau, t_2) e^{-2\sigma\tau} \cos(\omega_z(t_2, t_0)\tau) d\tau \\
 &\quad + \sin(\omega_z(t_2, t_0)t_0) \int_{-\infty}^{t_0} E'_0(\tau, t_2) e^{-2\sigma\tau} \sin(\omega_z(t_2, t_0)\tau) d\tau] \\
 &+ e^{2\sigma t_0} [\cos(\omega_z(t_2, t_0)t_0) \int_{-\infty}^{t_0} E'_{0n}(\tau, t_2) \cos(\omega_z(t_2, t_0)\tau) d\tau + \sin(\omega_z(t_2, t_0)t_0) \int_{-\infty}^{t_0} E'_{0n}(\tau, t_2) \sin(\omega_z(t_2, t_0)\tau) d\tau]
 \end{aligned}$$

276

(17)

In Section 2.1, it is shown that $0 < \omega_z(t_2, t_0) < \infty$, for all $|t_0| < \infty$, for a given finite value of t_2 .

277

In Section 4, it is shown that $\omega_z(t_2, t_0)$ is a **continuous** function of variable t_0 and t_2 , for all $|t_0| < \infty$ and $|t_2| < \infty$.

281

In Section 6, it is shown that $E_0(t)$ is **strictly decreasing** for $t > 0$.

282

Given that $\omega_z(t_2, t_0)$ is a continuous function of both t_0 and t_2 , we can find a suitable value of $t_0 = t_{0c}$ and $t_2 = t_{2c} = 2t_{0c}$ such that $\omega_z(t_{2c}, t_{0c})t_{0c} = \frac{\pi}{2}$. Given that $\omega_z(t_2, t_0)$ is a continuous function of t_0 and t_2 and given that t_0 is a continuous function, we see that the **product** of two continuous functions $\omega_z(t_2, t_0)t_0$ is a **continuous** function and is positive for $t_0 > 0$ because $0 < \omega_z(t_2, t_0) < \infty$.

288

We see that $\omega_z(t_2, t_0) > 0$ and is a **continuous** function of variable t_0 and t_2 , as t_0 and t_2 increase to a larger and larger finite value without bounds and that the order of $\omega_z(t_2, t_0)t_0$ is greater than 1 (Section 5). As t_0 and t_2 increase from zero to a larger and larger finite value without bounds, the continuous function $\omega_z(t_2, t_0)t_0$ starts from zero and increases with order greater than $O[1]$ and will pass through $\frac{\pi}{2}$.

294

We use $P_{odd}(t_2, t_0) + P_{odd}(t_2, -t_0) = 0$ as follows. We set $t_0 = t_{0c} > 0$ and $t_2 = t_{2c} = 2t_{0c}$ such that $\omega_z(t_{2c}, t_{0c})t_{0c} = \frac{\pi}{2}$ in Eq. 17 as follows. We use the fact that $\cos(\omega_z(t_{2c}, t_{0c})t_{0c}) = 0$, $\sin(\omega_z(t_{2c}, t_{0c})t_{0c}) = 1$ and $\omega_z(t_{2c}, -t_{0c}) = \omega_z(t_{2c}, t_{0c})$ shown in Section 2.3.

296

297

$$\begin{aligned}
 &\int_{-\infty}^{t_{0c}} E'_0(\tau, t_{2c}) e^{-2\sigma\tau} \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau + e^{2\sigma t_{0c}} \int_{-\infty}^{t_{0c}} E'_{0n}(\tau, t_{2c}) \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau \\
 &- \int_{-\infty}^{-t_{0c}} E'_0(\tau, t_{2c}) e^{-2\sigma\tau} \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau - e^{-2\sigma t_{0c}} \int_{-\infty}^{-t_{0c}} E'_{0n}(\tau, t_{2c}) \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau = 0
 \end{aligned}$$

299

We split the first two integrals in the left hand side of Eq. 18 and write as follows.

$$\begin{aligned}
& \left[\int_{-\infty}^{-t_{0c}} E'_0(\tau, t_{2c}) e^{-2\sigma\tau} \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau + \int_{-t_{0c}}^{t_{0c}} E'_0(\tau, t_{2c}) e^{-2\sigma\tau} \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau \right] \\
& + e^{2\sigma t_{0c}} \left[\int_{-\infty}^{-t_{0c}} E'_{0n}(\tau, t_{2c}) \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau + \int_{-t_{0c}}^{t_{0c}} E'_{0n}(\tau, t_{2c}) \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau \right] \\
& - \int_{-\infty}^{-t_{0c}} E'_0(\tau, t_{2c}) e^{-2\sigma\tau} \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau - e^{-2\sigma t_{0c}} \int_{-\infty}^{-t_{0c}} E'_{0n}(\tau, t_{2c}) \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau = 0
\end{aligned}
\tag{19}$$

300

301

We combine the terms with common integrals and cancel common terms in Eq. 19 as follows.

$$\begin{aligned}
& \int_{-t_{0c}}^{t_{0c}} E'_0(\tau, t_{2c}) e^{-2\sigma\tau} \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau + e^{2\sigma t_{0c}} \int_{-t_{0c}}^{t_{0c}} E'_{0n}(\tau, t_{2c}) \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau \\
& = -2 \sinh(2\sigma t_{0c}) \int_{-\infty}^{-t_{0c}} E'_{0n}(\tau, t_{2c}) \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau
\end{aligned}
\tag{20}$$

302

303

We can rearrange the terms in Eq. 20 as follows.

$$\begin{aligned}
& \int_{-t_{0c}}^{t_{0c}} [E'_0(\tau, t_{2c}) e^{-2\sigma\tau} + E'_{0n}(\tau, t_{2c}) e^{2\sigma t_{0c}}] \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau \\
& = -2 \sinh(2\sigma t_{0c}) \int_{-\infty}^{-t_{0c}} E'_{0n}(\tau, t_{2c}) \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau
\end{aligned}
\tag{21}$$

304

305

306

We denote the right hand side of Eq. 21 as *RHS*. We can split the integral in Eq. 21 using $\int_{-t_{0c}}^{t_{0c}} = \int_{-t_{0c}}^0 + \int_0^{t_{0c}}$ as follows.

$$\begin{aligned}
& \int_{-t_{0c}}^0 [E'_0(\tau, t_{2c}) e^{-2\sigma\tau} + E'_{0n}(\tau, t_{2c}) e^{2\sigma t_{0c}}] \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau \\
& + \int_0^{t_{0c}} [E'_0(\tau, t_{2c}) e^{-2\sigma\tau} + E'_{0n}(\tau, t_{2c}) e^{2\sigma t_{0c}}] \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau = RHS
\end{aligned}
\tag{22}$$

307

308

309

We substitute $\tau = -\tau$ in the first integral in Eq. 22 as follows. We use $E'_0(-\tau, t_{2c}) = E'_{0n}(\tau, t_{2c})$ and $E'_{0n}(-\tau, t_{2c}) = E'_0(\tau, t_{2c})$.

$$\begin{aligned}
& \int_{t_{0c}}^0 [E'_{0n}(\tau, t_{2c}) e^{2\sigma\tau} + E'_0(\tau, t_{2c}) e^{2\sigma t_{0c}}] \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau \\
& + \int_0^{t_{0c}} [E'_0(\tau, t_{2c}) e^{-2\sigma\tau} + E'_{0n}(\tau, t_{2c}) e^{2\sigma t_{0c}}] \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau = RHS
\end{aligned}$$

311 Given that $\int_{t_{0c}}^0 = -\int_0^{t_{0c}}$, we can simplify as follows.

$$\int_0^{t_{0c}} [E'_0(\tau, t_{2c})(e^{-2\sigma\tau} - e^{2\sigma t_{0c}}) + E'_{0n}(\tau, t_{2c})(-e^{2\sigma\tau} + e^{2\sigma t_{0c}})] \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau = RHS$$

312

313 We substitute $\tau = -\tau$ in the right hand side of Eq. 21 as follows. We use $E'_{0n}(-\tau, t_{2c}) = E'_0(\tau, t_{2c})$.

$$RHS = 2 \sinh(2\sigma t_{0c}) \int_{t_{0c}}^{\infty} E'_0(\tau, t_{2c}) \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau$$

314

315 We split the integral on the right hand side in Eq. 25 as follows.

$$RHS = 2 \sinh(2\sigma t_{0c}) \left[\int_0^{\infty} E'_0(\tau, t_{2c}) \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau - \int_0^{t_{0c}} E'_0(\tau, t_{2c}) \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau \right]$$

316

317 We consolidate the integrals with the term $\int_0^{t_{0c}} E'_0(\tau, t_{2c})$ in Eq. 24 and Eq. 26 as follows. We use
 318 $2 \sinh(2\sigma t_{0c}) = e^{2\sigma t_{0c}} - e^{-2\sigma t_{0c}}$.

$$\begin{aligned} \int_0^{t_{0c}} [E'_0(\tau, t_{2c})(e^{-2\sigma\tau} - e^{2\sigma t_{0c}} + e^{2\sigma t_{0c}} - e^{-2\sigma t_{0c}}) + E'_{0n}(\tau, t_{2c})(-e^{2\sigma\tau} + e^{2\sigma t_{0c}})] \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau \\ = 2 \sinh(2\sigma t_{0c}) \int_0^{\infty} E'_0(\tau, t_{2c}) \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau \end{aligned}$$

319

320 We cancel common terms in Eq. 27 as follows.

$$\begin{aligned} \int_0^{t_{0c}} [E'_0(\tau, t_{2c})(e^{-2\sigma\tau} - e^{2\sigma t_{0c}}) + E'_{0n}(\tau, t_{2c})(-e^{2\sigma\tau} + e^{2\sigma t_{0c}})] \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau \\ = 2 \sinh(2\sigma t_{0c}) \int_0^{\infty} E'_0(\tau, t_{2c}) \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau \end{aligned}$$

321

322 We substitute $E'_0(\tau, t_{2c}) = E_0(\tau - t_{2c}) - E_0(\tau + t_{2c})$ and $E'_{0n}(\tau, t_{2c}) = E'_0(-\tau, t_{2c}) = E_0(-\tau - t_{2c}) -$
 323 $E_0(-\tau + t_{2c})$. We see that $E_0(-\tau - t_{2c}) = E_0(\tau + t_{2c})$ and $E_0(-\tau + t_{2c}) = E_0(\tau - t_{2c})$ given that
 324 $E_0(\tau) = E_0(-\tau)$. Hence we see that $E'_{0n}(\tau, t_{2c}) = E_0(\tau + t_{2c}) - E_0(\tau - t_{2c}) = -E'_0(\tau, t_{2c})$. We can
 325 write Eq. 28 as follows.

$$\begin{aligned}
& \int_0^{t_{0c}} (E_0(\tau - t_{2c}) - E_0(\tau + t_{2c}))(e^{-2\sigma\tau} - e^{-2\sigma t_{0c}} + e^{2\sigma\tau} - e^{2\sigma t_{0c}}) \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau \\
& = 2 \sinh(2\sigma t_{0c}) \int_0^\infty (E_0(\tau - t_{2c}) - E_0(\tau + t_{2c})) \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau
\end{aligned}$$

(29)

We substitute $2 \cosh(2\sigma\tau) = e^{2\sigma\tau} + e^{-2\sigma\tau}$ and $2 \cosh(2\sigma t_{0c}) = e^{2\sigma t_{0c}} + e^{-2\sigma t_{0c}}$ and cancel the common factor of 2 in Eq. 29 as follows.

$$\begin{aligned}
& \int_0^{t_{0c}} (E_0(\tau - t_{2c}) - E_0(\tau + t_{2c}))(\cosh(2\sigma\tau) - \cosh(2\sigma t_{0c})) \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau \\
& = \sinh(2\sigma t_{0c}) \int_0^\infty (E_0(\tau - t_{2c}) - E_0(\tau + t_{2c})) \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau
\end{aligned}$$

(30)

Next Step:

We denote the right hand side of Eq. 30 as *RHS*. We substitute $\tau + t_{2c} = \tau'$ in the right hand side of Eq. 30 and then substitute $\tau' = \tau$. Similarly we substitute $\tau - t_{2c} = \tau'$ as follows.

$$\begin{aligned}
RHS = & \sinh(2\sigma t_{0c}) [\cos(\omega_z(t_{2c}, t_{0c}))t_{2c} \int_{-t_{2c}}^\infty E_0(\tau) \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau \\
& + \sin(\omega_z(t_{2c}, t_{0c}))t_{2c} \int_{-t_{2c}}^\infty E_0(\tau) \cos(\omega_z(t_{2c}, t_{0c})\tau) d\tau \\
& - \cos(\omega_z(t_{2c}, t_{0c}))t_{2c} \int_{t_{2c}}^\infty E_0(\tau) \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau + \sin(\omega_z(t_{2c}, t_{0c}))t_{2c} \int_{t_{2c}}^\infty E_0(\tau) \cos(\omega_z(t_{2c}, t_{0c})\tau) d\tau]
\end{aligned}$$

(31)

In Eq. 31, given that $\omega_z(t_{2c}, t_{0c})t_{0c} = \frac{\pi}{2}$ and $t_{2c} = 2t_{0c}$ and hence $\omega_z(t_{2c}, t_{0c})t_{2c} = 2\frac{\pi}{2} = \pi$ and $\sin(\omega_z(t_{2c}, t_{0c})t_{2c}) = 0$ and $\cos(\omega_z(t_{2c}, t_{0c})t_{2c}) = -1$. Hence we cancel common terms and write Eq. 31 and Eq. 30 as follows.

$$\begin{aligned}
& \int_0^{t_{0c}} (E_0(\tau - t_{2c}) - E_0(\tau + t_{2c}))(\cosh(2\sigma\tau) - \cosh(2\sigma t_{0c})) \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau \\
& = -\sinh(2\sigma t_{0c}) [\int_{-t_{2c}}^\infty E_0(\tau) \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau - \int_{t_{2c}}^\infty E_0(\tau) \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau]
\end{aligned}$$

(32)

We use $\int_{-t_{2c}}^\infty E_0(\tau) \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau = \int_{-t_{2c}}^{t_{2c}} E_0(\tau) \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau + \int_{t_{2c}}^\infty E_0(\tau) \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau$ and cancel the common term $\int_{t_{2c}}^\infty E_0(\tau) \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau$ in Eq. 32 as follows. Given that $E_0(\tau) \sin(\omega_z(t_{2c}, t_{0c})\tau)$ is an **odd** function of variable τ , we get $\int_{-t_{2c}}^{t_{2c}} E_0(\tau) \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau = 0$.

$$\int_0^{t_{0c}} (E_0(\tau - t_{2c}) - E_0(\tau + t_{2c}))(\cosh(2\sigma\tau) - \cosh(2\sigma t_{0c}) \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau = 0$$

(33)

We can multiply Eq. 33 by a factor of -1 as follows.

$$\int_0^{t_{0c}} [E_0(\tau - t_{2c}) - E_0(\tau + t_{2c})](\cosh 2\sigma t_{0c} - \cosh(2\sigma\tau) \sin(\omega_z(t_{2c}, t_{0c})\tau) d\tau = 0$$

(34)

In Eq. 34, given that $\omega_z(t_{2c}, t_{0c})t_{0c} = \frac{\pi}{2}$, as τ varies over the interval $[0, t_{0c}]$, $\omega_z(t_{2c}, t_{0c})\tau = \frac{\pi\tau}{2t_{0c}}$ varies from $[0, \frac{\pi}{2}]$ and the sinusoidal function is > 0 , in the interval $0 < \tau < t_{0c}$, for $t_{0c} > 0$.

In Eq. 34, we see that in the interval $0 < \tau < t_{0c}$, the integral on the left hand side is > 0 for $t_{0c} > 0$, because each of the terms in the integrand are > 0 , in the interval $0 < \tau < t_{0c}$ as follows. Given that $E_0(t)$ is a **strictly decreasing** function for $t > 0$ (Section 6), we see that $E_0(\tau - t_{2c}) - E_0(\tau + t_{2c})$ is > 0 (Section 6.3) in the interval $0 < \tau < t_{0c}$. The term $(\cosh(2\sigma t_{0c}) - \cosh(2\sigma\tau))$ is > 0 in the interval $0 < \tau < t_{0c}$ and the integrand is zero at $\tau = 0$ and $\tau = t_{0c}$ and hence the integral **cannot** equal zero, as required by the right hand side of Eq. 34. Hence this leads to a **contradiction** for $0 < \sigma < \frac{1}{2}$.

For $\sigma = 0$, both sides of Eq. 34 is zero and **does not** lead to a contradiction.

We have shown this result for $0 < \sigma < \frac{1}{2}$ and then use the property $\xi(\frac{1}{2} + \sigma + i\omega) = \xi(\frac{1}{2} - \sigma - i\omega)$ to show the result for $-\frac{1}{2} < \sigma < 0$. Hence we have produced a **contradiction** of **Statement 1** that the Fourier Transform of the function $E_p(t) = E_0(t)e^{-\sigma t}$ has a zero at $\omega = \omega_0$ for $0 < |\sigma| < \frac{1}{2}$.

Therefore, the assumption in **Statement 1** that Riemann's Xi Function given by $\xi(\frac{1}{2} + \sigma + i\omega) = E_{p\omega}(\omega)$ has a zero at $\omega = \omega_0$, where ω_0 is real and finite, leads to a **contradiction** for the region $0 < |\sigma| < \frac{1}{2}$ which corresponds to the critical strip excluding the critical line. This means $\zeta(s)$ does not have non-trivial zeros in the critical strip excluding the critical line and we have proved Riemann's Hypothesis.

4. $\omega_z(t_2, t_0)$ is a continuous function of t_0 and t_2

We see from Section 2.1 that $\omega_z(t_2, t_0)$ is shown to be **finite and non-zero** for all $|t_0| < \infty$ and $|t_2| < \infty$ and that $\omega_z(t_2, t_0)$ is an even function of variable t_0 , for a given value of t_2 . For a given t_2 and t_0 , $\omega_z(t_2, t_0)$ can have more than one value, but we consider only the first zero crossing away from origin in the section below, where $G_R(\omega, t_2, t_0)$ crosses the zero line to the opposite sign, as detailed in **Lemma 1** in Section 2.1 and $\frac{\partial G_R(\omega, t_2, t_0)}{\partial \omega} \neq 0$ at $\omega = \omega_z(t_2, t_0)$. (example plot)

We consider the Fourier transform of the even part of $g(t, t_2, t_0)$ given by $G_R(\omega, t_2, t_0)$ in the section below and show that, under this Fourier transformation, as we change t_0 , the zero crossing in $G_R(\omega, t_2, t_0)$ given by $\omega_z(t_2, t_0)$ is a continuous function of t_0 , for all $|t_0| < \infty$, for **each** finite value of t_2 . This is shown in the steps below. For a given **finite** value of t_2 , $G_R(\omega, t_2, t_0)$ is a function of two

variables ω and t_0 , and we use Implicit Function Theorem in R^2 .

• It is shown in Section 4.1 that $G_R(\omega, t_2, t_0)$ is partially differentiable at least twice with respect to ω , as shown in Eq. 35.

• It is shown in Section 4.2 that $G_R(\omega, t_2, t_0)$ is partially differentiable at least twice with respect to t_0 , as shown in Eq. 37 and Eq. 42.

• It is shown in Section 4.3 that the zero crossing in $G_R(\omega, t_2, t_0)$ given by $\omega_z(t_2, t_0)$, is a **continuous** function of t_0 , for a given t_2 , using **Implicit Function Theorem** in R^2 .

• It is shown in Section 4.4 that $\omega_z(t_2, t_0)$ is a **continuous** function of t_0 and t_2 , for all $|t_0| < \infty$ and $|t_2| < \infty$, using **Implicit Function Theorem** in R^3 .

4.1. $G_R(\omega, t_2, t_0)$ is *partially differentiable twice as a function of ω*

$G_R(\omega, t_2, t_0)$ in Eq. 14 is copied below and we can expand $G_R(\omega, t_2, t_0)$ in Eq. 35 by substituting $\tau + t_0 = t$ and expanding it, similar to Eq. 17.

$$\begin{aligned}
G_R(\omega, t_2, t_0) &= e^{-2\sigma t_0} \int_{-\infty}^0 [E'_0(\tau + t_0, t_2)e^{-2\sigma\tau} + E'_{0n}(\tau - t_0, t_2)] \cos(\omega\tau) d\tau \\
&+ e^{2\sigma t_0} \int_{-\infty}^0 [E'_0(\tau - t_0, t_2)e^{-2\sigma\tau} + E'_{0n}(\tau + t_0, t_2)] \cos(\omega\tau) d\tau = G'_{1R}(\omega, t_2, t_0) + G'_{1R}(\omega, t_2, -t_0) \\
G'_{1R}(\omega, t_2, t_0) &= [\cos(\omega t_0) \int_{-\infty}^{t_0} E'_0(\tau, t_2)e^{-2\sigma\tau} \cos(\omega\tau) d\tau + \sin(\omega t_0) \int_{-\infty}^{t_0} E'_0(\tau, t_2)e^{-2\sigma\tau} \sin(\omega\tau) d\tau] \\
&+ e^{-2\sigma t_0} [\cos(\omega t_0) \int_{-\infty}^{-t_0} E'_{0n}(\tau, t_2) \cos(\omega\tau) d\tau - \sin(\omega t_0) \int_{-\infty}^{-t_0} E'_{0n}(\tau, t_2) \sin(\omega\tau) d\tau]
\end{aligned} \tag{35}$$

We could then use $E'_0(t, t_2) = (E_0(t - t_2) - E_0(t + t_2))$ and $E'_{0n}(t, t_2) = E'_0(-t, t_2)$ and substitute $t + t_2 = t$ and $t - t_2 = t'$ and expanding it using the procedure used in Eq. 35. The integrands are absolutely integrable and we could then use theorem of dominated convergence as follows.

$G_R(\omega, t_2, t_0)$ is partially differentiable at least twice with respect to ω and the integrals converge in Eq. 35 for $0 < \sigma < \frac{1}{2}$, because the term $\tau^r E'_0(\tau \pm t_0, t_2)e^{-2\sigma\tau}$ has exponential asymptotic fall-off rate as $|\tau| \rightarrow \infty$, for $r = 0, 1, 2$ (Appendix B.4). The integrands are absolutely integrable and the integrands are analytic functions of variables ω and t_0 , for a given t_2 . We can interchange the order of partial differentiation and integration in Eq. 36 using theorem of dominated convergence, recursively as follows.(link) (We could also use theorem 3 in link and link.)

$$\begin{aligned}
\frac{\partial G_R(\omega, t_2, t_0)}{\partial \omega} &= -[e^{-2\sigma t_0} \int_{-\infty}^0 \tau [E'_0(\tau + t_0, t_2)e^{-2\sigma\tau} + E'_{0n}(\tau - t_0, t_2)] \sin(\omega\tau) d\tau \\
&\quad + e^{2\sigma t_0} \int_{-\infty}^0 \tau [E'_0(\tau - t_0, t_2)e^{-2\sigma\tau} + E'_{0n}(\tau + t_0, t_2)] \sin(\omega\tau) d\tau] \\
\frac{\partial^2 G_R(\omega, t_2, t_0)}{\partial \omega^2} &= -[e^{-2\sigma t_0} \int_{-\infty}^0 \tau^2 [E'_0(\tau + t_0, t_2)e^{-2\sigma\tau} + E'_{0n}(\tau - t_0, t_2)] \cos(\omega\tau) d\tau \\
&\quad + e^{2\sigma t_0} \int_{-\infty}^0 \tau^2 [E'_0(\tau - t_0, t_2)e^{-2\sigma\tau} + E'_{0n}(\tau + t_0, t_2)] \cos(\omega\tau) d\tau]
\end{aligned}$$

(36)

406

407 *4.2. $G_R(\omega, t_2, t_0)$ is partially differentiable twice as a function of t_0*

408

409 $G_R(\omega, t_2, t_0)$ is partially differentiable at least twice as a function of t_0 and the integrals converge
410 in Eq. 37 and Eq. 42 shown as follows. The integrands are absolutely integrable because the term
411 $E'_0(\tau \pm t_0, t_2)e^{-2\sigma\tau}$ has exponential asymptotic fall-off rate as $|\tau| \rightarrow \infty$ (Appendix B.4). The
412 integrands are analytic functions of variables ω and t_0 , for a given t_2 and we can expand $G_R(\omega, t_2, t_0)$
413 in Eq. 37 by substituting $\tau + t_0 = t$ and expanding it, similar to Eq. 35. We can interchange the
414 order of partial differentiation and integration in Eq. 37 and Eq. 42 using theorem of dominated
415 convergence as follows. (link) (We could also use theorem 3 in link and link)

$$\begin{aligned}
G_R(\omega, t_2, t_0) &= e^{-2\sigma t_0} \int_{-\infty}^0 [E'_0(\tau + t_0, t_2)e^{-2\sigma\tau} + E'_{0n}(\tau - t_0, t_2)] \cos(\omega\tau) d\tau \\
&\quad + e^{2\sigma t_0} \int_{-\infty}^0 [E'_0(\tau - t_0, t_2)e^{-2\sigma\tau} + E'_{0n}(\tau + t_0, t_2)] \cos(\omega\tau) d\tau \\
\frac{\partial G_R(\omega, t_2, t_0)}{\partial t_0} &= -2\sigma e^{-2\sigma t_0} \int_{-\infty}^0 [E'_0(\tau + t_0, t_2)e^{-2\sigma\tau} + E'_{0n}(\tau - t_0, t_2)] \cos(\omega\tau) d\tau \\
&\quad + e^{-2\sigma t_0} \int_{-\infty}^0 \frac{\partial(E'_0(\tau + t_0, t_2)e^{-2\sigma\tau} + E'_{0n}(\tau - t_0, t_2))}{\partial t_0} \cos(\omega\tau) d\tau \\
&\quad + 2\sigma e^{2\sigma t_0} \int_{-\infty}^0 [E'_0(\tau - t_0, t_2)e^{-2\sigma\tau} + E'_{0n}(\tau + t_0, t_2)] \cos(\omega\tau) d\tau \\
&\quad + e^{2\sigma t_0} \int_{-\infty}^0 \frac{\partial(E'_0(\tau - t_0, t_2)e^{-2\sigma\tau} + E'_{0n}(\tau + t_0, t_2))}{\partial t_0} \cos(\omega\tau) d\tau
\end{aligned}$$

(37)

416

417 We can show that the integrals in Eq. 37 converge, as follows. We see that $E'_0(\tau + t_0, t_2) =$
418 $E_0(\tau + t_0 - t_2) - E_0(\tau + t_0 + t_2)$.

419

420 We see that $E'_0(\tau, t_2) = E_0(\tau - t_2) - E_0(\tau + t_2)$ and $E'_{0n}(\tau, t_2) = E'_0(-\tau, t_2) = -E'_0(\tau, t_2)$ because
421 $E_0(-\tau) = E_0(\tau)$. We get $E'_{0n}(\tau - t_0, t_2) = -E'_0(\tau - t_0, t_2) = E_0(\tau - t_0 + t_2) - E_0(\tau - t_0 - t_2)$
422 given that $E_0(\tau) = E_0(-\tau)$. We see that the third integral in Eq. 37 converges because the term
423 $E'_0(\tau \pm t_0, t_2)e^{-2\sigma\tau}$ has exponential asymptotic fall-off rate as $|\tau| \rightarrow \infty$ (Appendix B.4).

424

425 We consider the integrand in the fourth integral in Eq. 37 first and use the results in the above
 426 paragraph.

$$\begin{aligned} \frac{\partial(E'_0(\tau + t_0, t_2)e^{-2\sigma\tau} + E'_{0n}(\tau - t_0, t_2))}{\partial t_0} &= \frac{\partial(E_0(\tau + t_0 - t_2)e^{-2\sigma\tau} - E_0(\tau + t_0 + t_2)e^{-2\sigma\tau})}{\partial t_0} \\ &\quad + \frac{\partial(E_0(\tau - t_0 + t_2) - E_0(\tau - t_0 - t_2))}{\partial t_0} \end{aligned} \quad (38)$$

428 We consider the term $E_0(\tau + t_0 + t_2)$ first in Eq. 38 and can show that the integrals converge in
 429 Eq. 37, as follows.

$$\begin{aligned} E_0(\tau) &= 2 \sum_{n=1}^{\infty} [2\pi^2 n^4 e^{4\tau} - 3\pi n^2 e^{2\tau}] e^{-\pi n^2 e^{2\tau}} e^{\frac{\tau}{2}} \\ E_0(\tau + t_2 + t_0) &= 2 \sum_{n=1}^{\infty} [2\pi^2 n^4 e^{4\tau} e^{4(t_2+t_0)} - 3\pi n^2 e^{2\tau} e^{2(t_2+t_0)}] e^{-\pi n^2 e^{2\tau} e^{2(t_2+t_0)}} e^{\frac{\tau}{2}} e^{\frac{(t_2+t_0)}{2}} \end{aligned} \quad (39)$$

431 We can show that $\frac{\partial}{\partial t_0} E_0(\tau + t_2 + t_0) = \frac{\partial}{\partial \tau} E_0(\tau + t_2 + t_0)$ as follows. (**Result A**)

$$\begin{aligned} \frac{\partial}{\partial t_0} E_0(\tau + t_2 + t_0) &= 2 \sum_{n=1}^{\infty} e^{-\pi n^2 e^{2\tau} e^{2(t_2+t_0)}} e^{\frac{\tau}{2}} e^{\frac{(t_2+t_0)}{2}} [8\pi^2 n^4 e^{4\tau} e^{4(t_2+t_0)} - 6\pi n^2 e^{2\tau} e^{2(t_2+t_0)}] \\ &\quad + \left(\frac{1}{2} - 2\pi n^2 e^{2\tau} e^{2(t_2+t_0)}\right) (2\pi^2 n^4 e^{4\tau} e^{4(t_2+t_0)} - 3\pi n^2 e^{2\tau} e^{2(t_2+t_0)})] \\ \frac{\partial}{\partial \tau} E_0(\tau + t_2 + t_0) &= 2 \sum_{n=1}^{\infty} e^{-\pi n^2 e^{2\tau} e^{2(t_2+t_0)}} e^{\frac{\tau}{2}} e^{\frac{(t_2+t_0)}{2}} [8\pi^2 n^4 e^{4\tau} e^{4(t_2+t_0)} - 6\pi n^2 e^{2\tau} e^{2(t_2+t_0)}] \\ &\quad + \left(\frac{1}{2} - 2\pi n^2 e^{2\tau} e^{2(t_2+t_0)}\right) (2\pi^2 n^4 e^{4\tau} e^{4(t_2+t_0)} - 3\pi n^2 e^{2\tau} e^{2(t_2+t_0)})] \end{aligned} \quad (40)$$

433 We can replace t_0 by $-t_0$ in Eq. 40 and show that $\frac{\partial}{\partial t_0} E_0(\tau + t_2 - t_0) = -\frac{\partial}{\partial \tau} E_0(\tau + t_2 - t_0)$. (**Result B**)

434
 435 We can write the term $E_0(\tau + t_0 + t_2)e^{-2\sigma\tau}$ in Eq. 38, corresponding to the term in the fourth
 436 integral in Eq. 37, using Result A, as follows.

$$\begin{aligned} \int_{-\infty}^0 \frac{\partial(E_0(\tau + t_2 + t_0)e^{-2\sigma\tau})}{\partial t_0} \cos(\omega\tau) d\tau &= \int_{-\infty}^0 \frac{\partial(E_0(\tau + t_2 + t_0))}{\partial \tau} e^{-2\sigma\tau} \cos(\omega\tau) d\tau \\ &= \int_{-\infty}^0 \frac{\partial(E_0(\tau + t_2 + t_0)e^{-2\sigma\tau} \cos(\omega\tau))}{\partial \tau} d\tau - \int_{-\infty}^0 E_0(\tau + t_2 + t_0) \frac{\partial(e^{-2\sigma\tau} \cos(\omega\tau))}{\partial \tau} d\tau \\ &= [E_0(\tau + t_2 + t_0)e^{-2\sigma\tau} \cos(\omega\tau)]_{-\infty}^0 + \omega \int_{-\infty}^0 E_0(\tau + t_2 + t_0) e^{-2\sigma\tau} \sin(\omega\tau) d\tau \\ &\quad + 2\sigma \int_{-\infty}^0 E_0(\tau + t_2 + t_0) e^{-2\sigma\tau} \cos(\omega\tau) d\tau \end{aligned}$$

We see that the integrals in Eq. 41 converge and hence the integral $\int_{-\infty}^0 \frac{\partial(E_0(\tau+t_2+t_0)e^{-2\sigma\tau})}{\partial t_0} \cos(\omega\tau) d\tau$ in Eq. 41 also converges. We set $\sigma = 0$ and $t_0 = -t_0$ and see that the integral $\int_{-\infty}^0 \frac{\partial(E_0(\tau+t_2-t_0))}{\partial t_0} \cos(\omega\tau) d\tau$ in Eq. 38 also converges, using Result B.

We set $t_2 = -t_2$ in Eq. 39 to Eq. 41 and see that the integral $\int_{-\infty}^0 \frac{\partial(E_0(\tau+t_0-t_2)e^{-2\sigma\tau})}{\partial t_0} \cos(\omega\tau) d\tau$ in Eq. 38 also converges. We set $\sigma = 0$ and $t_0 = -t_0$ and see that the integral $\int_{-\infty}^0 \frac{\partial(E_0(\tau-t_0-t_2))}{\partial t_2} \cos(\omega\tau) d\tau$ in Eq. 38 also converges, using Result B. Hence the fourth integral in Eq. 37 corresponding to the terms in Eq. 38, also converges.

We can see that the last two integrals in Eq. 37 converge, by setting $t_0 = -t_0$ and using Result B. Hence all the integrals in Eq. 37 converge.

The second partial derivative of $G_R(\omega, t_2, t_0)$ with respect to t_0 is given by $\frac{\partial^2 G_R(\omega, t_2, t_0)}{\partial t_0^2} = \frac{\partial}{\partial t_0} \frac{\partial G_R(\omega, t_2, t_0)}{\partial t_0}$ as follows. We use the result in Eq. 41 and we can interchange the order of partial differentiation and integration in Eq. 42 using theorem of dominated convergence as follows.

$$\begin{aligned} \frac{\partial^2 G_R(\omega, t_2, t_0)}{\partial t_0^2} &= 4\sigma^2 e^{-2\sigma t_0} \int_{-\infty}^0 [E'_0(\tau + t_0, t_2)e^{-2\sigma\tau} + E'_{0n}(\tau - t_0, t_2)] \cos(\omega\tau) d\tau \\ &\quad - 4\sigma e^{-2\sigma t_0} \int_{-\infty}^0 \frac{\partial(E'_0(\tau + t_0, t_2)e^{-2\sigma\tau} + E'_{0n}(\tau - t_0, t_2))}{\partial t_0} \cos(\omega\tau) d\tau \\ &\quad + e^{-2\sigma t_0} \int_{-\infty}^0 \frac{\partial^2(E'_0(\tau + t_0, t_2)e^{-2\sigma\tau} + E'_{0n}(\tau - t_0, t_2))}{\partial t_0^2} \cos(\omega\tau) d\tau \\ &\quad + 4\sigma^2 e^{2\sigma t_0} \int_{-\infty}^0 [E'_0(\tau - t_0, t_2)e^{-2\sigma\tau} + E'_{0n}(\tau + t_0, t_2)] \cos(\omega\tau) d\tau \\ &\quad + 4\sigma e^{2\sigma t_0} \int_{-\infty}^0 \frac{\partial(E'_0(\tau - t_0, t_2)e^{-2\sigma\tau} + E'_{0n}(\tau + t_0, t_2))}{\partial t_0} \cos(\omega\tau) d\tau \\ &\quad + e^{2\sigma t_0} \int_{-\infty}^0 \frac{\partial^2(E'_0(\tau - t_0, t_2)e^{-2\sigma\tau} + E'_{0n}(\tau + t_0, t_2))}{\partial t_0^2} \cos(\omega\tau) d\tau \end{aligned}$$

We can use the above procedure in Eq. 39 to Eq. 41 for the term $\frac{\partial^2(E'_0(\tau+t_0,t_2)e^{-2\sigma\tau}+E'_{0n}(\tau-t_0,t_2))}{\partial t_0^2} = \frac{\partial I(\tau, t_0, t_2)}{\partial t_0}$ where $I(\tau, t_0, t_2) = \frac{\partial(E'_0(\tau+t_0,t_2)e^{-2\sigma\tau}+E'_{0n}(\tau-t_0,t_2))}{\partial t_0}$ in the third integral in Eq. 42 and we can show that it converges, using the procedure used in Eq. 41 twice.

We can see that the last three integrals in Eq. 42 converge, by setting $t_0 = -t_0$ and using Result B. Hence all the integrals in Eq. 37 and Eq. 42 converge.

4.3. Zero Crossings in $G_R(\omega, t_2, t_0)$ move continuously as a function of t_0 , for a given t_2 .

We use **Implicit Function Theorem** for the two dimensional case (link and link). Given that $G_R(\omega, t_2, t_0)$ is partially differentiable with respect to ω and t_0 , for a given fixed value of t_2 ,

with continuous partial derivatives (Section 4.1 and Section 4.2) and given that $G_R(\omega, t_2, t_0) = 0$ at $\omega = \omega_z(t_2, t_0)$ and $\frac{\partial G_R(\omega, t_2, t_0)}{\partial \omega} \neq 0$ at $\omega = \omega_z(t_2, t_0)$ (using Lemma 1), we see that $\omega_z(t_2, t_0)$ is differentiable function of t_0 , for all $|t_0| < \infty$.

Hence $\omega_z(t_2, t_0)$ is a **continuous** function of t_0 for all $|t_0| < \infty$, for each finite value of t_2 .

• It is shown in Section 4.5 that $G_R(\omega, t_2, t_0)$ is partially differentiable at least twice with respect to t_2 . We can use the procedure in previous subsections and Implicit Function Theorem and show that $\omega_z(t_2, t_0)$ is a **continuous** function of t_2 , for all $|t_2| < \infty$, for **each** finite value of t_0 .

4.4. Zero Crossings in $G_R(\omega, t_2, t_0)$ move continuously as a function of t_0 and t_2

We can use the procedure in previous subsections and show that $\omega_z(t_2, t_0)$ is a **continuous** function of t_2 and t_0 , for all $|t_0| < \infty$ and $|t_2| < \infty$, using Implicit Function Theorem in R^3 .

We use **Implicit Function Theorem** for the three dimensional case (link). Given that $G_R(\omega, t_2, t_0)$ is partially differentiable with respect to ω and t_0 and t_2 , with continuous partial derivatives (Section 4.1, Section 4.2 and Section 4.5) and given that $G_R(\omega, t_2, t_0) = 0$ at $\omega = \omega_z(t_2, t_0)$ and $\frac{\partial G_R(\omega, t_2, t_0)}{\partial \omega} \neq 0$ at $\omega = \omega_z(t_2, t_0)$ (using Lemma 1), we see that $\omega_z(t_2, t_0)$ is differentiable function of t_0 and t_2 , for all $|t_0| < \infty$ and $|t_2| < \infty$.

Hence $\omega_z(t_2, t_0)$ is a **continuous** function of t_0 and t_2 , for all $|t_0| < \infty$ and $|t_2| < \infty$ (**Result E**).

4.5. $G_R(\omega, t_2, t_0)$ is partially differentiable twice as a function of t_2

$G_R(\omega, t_2, t_0)$ is partially differentiable at least twice as a function of t_2 and the integrals converge in Eq. 43 and Eq. 47 shown as follows. The integrands are analytic functions of variables ω and t_2 , for a given t_0 and we can expand $G_R(\omega, t_2, t_0)$ in Eq. 43 by substituting $\tau + t_0 = t$ and expanding it, similar to Eq. 35. We can interchange the order of partial differentiation and integration in Eq. 43 using theorem of dominated convergence as follows. (link) (We could also use theorem 3 in link and link)

$$\begin{aligned}
G_R(\omega, t_2, t_0) &= e^{-2\sigma t_0} \int_{-\infty}^0 [E'_0(\tau + t_0, t_2)e^{-2\sigma\tau} + E'_{0n}(\tau - t_0, t_2)] \cos(\omega\tau) d\tau \\
&\quad + e^{2\sigma t_0} \int_{-\infty}^0 [E'_0(\tau - t_0, t_2)e^{-2\sigma\tau} + E'_{0n}(\tau + t_0, t_2)] \cos(\omega\tau) d\tau \\
\frac{\partial G_R(\omega, t_2, t_0)}{\partial t_2} &= e^{-2\sigma t_0} \int_{-\infty}^0 \frac{\partial(E'_0(\tau + t_0, t_2)e^{-2\sigma\tau} + E'_{0n}(\tau - t_0, t_2))}{\partial t_2} \cos(\omega\tau) d\tau \\
&\quad + e^{2\sigma t_0} \int_{-\infty}^0 \frac{\partial(E'_0(\tau - t_0, t_2)e^{-2\sigma\tau} + E'_{0n}(\tau + t_0, t_2))}{\partial t_2} \cos(\omega\tau) d\tau
\end{aligned} \tag{43}$$

We use the procedure outlined in Eq. 38 to Eq. 41, with t_0 replaced by t_2 and show that all the integrals in Eq. 43 converge, as follows.

498 We see that $E'_0(\tau + t_0, t_2) = E_0(\tau + t_0 - t_2) - E_0(\tau + t_0 + t_2)$ and $E'_{0n}(\tau - t_0, t_2) = E_0(\tau - t_0 + t_2) -$
 499 $E_0(\tau - t_0 - t_2)$ (the paras above Eq. 38). We consider the integrand in the third integral in Eq. 43
 500 first.

$$\begin{aligned} \frac{\partial(E'_0(\tau + t_0, t_2)e^{-2\sigma\tau} + E'_{0n}(\tau - t_0, t_2))}{\partial t_2} &= \frac{\partial(E_0(\tau + t_0 - t_2)e^{-2\sigma\tau} - E_0(\tau + t_0 + t_2)e^{-2\sigma\tau})}{\partial t_2} \\ &\quad + \frac{\partial(E_0(\tau - t_0 + t_2) - E_0(\tau - t_0 - t_2))}{\partial t_2} \end{aligned} \quad (44)$$

501
 502 We consider the term $E_0(\tau + t_0 + t_2)$ first and can show that the integrals converge in Eq. 43, as
 503 follows. We consider Eq. 39 and show that $\frac{\partial}{\partial t_2}E_0(\tau + t_2 + t_0) = \frac{\partial}{\partial \tau}E_0(\tau + t_2 + t_0)$ as follows. (**Result**
 504 **C**)

$$\begin{aligned} \frac{\partial}{\partial t_2}E_0(\tau + t_2 + t_0) &= 2 \sum_{n=1}^{\infty} e^{-\pi n^2 e^{2\tau} e^{2(t_2+t_0)}} e^{\frac{\tau}{2}} e^{\frac{(t_2+t_0)}{2}} [8\pi^2 n^4 e^{4\tau} e^{4(t_2+t_0)} - 6\pi n^2 e^{2\tau} e^{2(t_2+t_0)}] \\ &\quad + \left(\frac{1}{2} - 2\pi n^2 e^{2\tau} e^{2(t_2+t_0)}\right) (2\pi^2 n^4 e^{4\tau} e^{4(t_2+t_0)} - 3\pi n^2 e^{2\tau} e^{2(t_2+t_0)})] \\ \frac{\partial}{\partial \tau}E_0(\tau + t_2 + t_0) &= 2 \sum_{n=1}^{\infty} e^{-\pi n^2 e^{2\tau} e^{2(t_2+t_0)}} e^{\frac{\tau}{2}} e^{\frac{(t_2+t_0)}{2}} [8\pi^2 n^4 e^{4\tau} e^{4(t_2+t_0)} - 6\pi n^2 e^{2\tau} e^{2(t_2+t_0)}] \\ &\quad + \left(\frac{1}{2} - 2\pi n^2 e^{2\tau} e^{2(t_2+t_0)}\right) (2\pi^2 n^4 e^{4\tau} e^{4(t_2+t_0)} - 3\pi n^2 e^{2\tau} e^{2(t_2+t_0)})] \end{aligned} \quad (45)$$

505
 506 We can replace t_2 by $-t_2$ in Eq. 45 and show that $\frac{\partial}{\partial t_2}E_0(\tau + t_0 - t_2) = -\frac{\partial}{\partial \tau}E_0(\tau + t_0 - t_2)$ (**Result**
 507 **D**). We can write the term $E_0(\tau + t_0 + t_2)e^{-2\sigma\tau}$ in Eq. 44, corresponding to the term in the third
 508 integral in Eq. 43 as follows.

$$\begin{aligned} \int_{-\infty}^0 \frac{\partial(E_0(\tau + t_2 + t_0)e^{-2\sigma\tau})}{\partial t_2} \cos(\omega\tau) d\tau &= \int_{-\infty}^0 \frac{\partial(E_0(\tau + t_2 + t_0))}{\partial \tau} e^{-2\sigma\tau} \cos(\omega\tau) d\tau \\ &= \int_{-\infty}^0 \frac{\partial(E_0(\tau + t_2 + t_0)e^{-2\sigma\tau} \cos(\omega\tau))}{\partial \tau} d\tau - \int_{-\infty}^0 E_0(\tau + t_2 + t_0) \frac{\partial(e^{-2\sigma\tau} \cos(\omega\tau))}{\partial \tau} d\tau \\ &= [E_0(\tau + t_2 + t_0)e^{-2\sigma\tau} \cos(\omega\tau)]_{-\infty}^0 + \omega \int_{-\infty}^0 E_0(\tau + t_2 + t_0) e^{-2\sigma\tau} \sin(\omega\tau) d\tau \\ &\quad + 2\sigma \int_{-\infty}^0 E_0(\tau + t_2 + t_0) e^{-2\sigma\tau} \cos(\omega\tau) d\tau \end{aligned} \quad (46)$$

509
 510 We see that the integrals in Eq. 46 converge and hence the integral $\int_{-\infty}^0 \frac{\partial(E_0(\tau + t_2 + t_0)e^{-2\sigma\tau})}{\partial t_2} \cos(\omega\tau) d\tau$
 511 in Eq. 46 also converges. We set $\sigma = 0$ and $t_0 = -t_0$ and see that the integral $\int_{-\infty}^0 \frac{\partial(E_0(\tau + t_2 - t_0))}{\partial t_2} \cos(\omega\tau) d\tau$
 512 in Eq. 44 also converges.

513
 514 We set $t_2 = -t_2$ in Eq. 45 to Eq. 46 and see that the integral $\int_{-\infty}^0 \frac{\partial(E_0(\tau + t_0 - t_2)e^{-2\sigma\tau})}{\partial t_2} \cos(\omega\tau) d\tau$
 515 in Eq. 44 also converges, using Result D. We set $\sigma = 0$ and $t_0 = -t_0$ and see that the integral

516 $\int_{-\infty}^0 \frac{\partial(E_0(\tau-t_2-t_0))}{\partial t_2} \cos(\omega\tau) d\tau$ in Eq. 44 also converges. Hence the third integral in Eq. 43 correspond-
 517 ing to the terms in Eq. 44, also converges. We set $t_0 = -t_0$ and see that the fourth integral in Eq. 43
 518 also converges.

519

520 The second partial derivative of $G_R(\omega, t_2, t_0)$ with respect to t_2 is given by $\frac{\partial^2 G_R(\omega, t_2, t_0)}{\partial t_2^2} = \frac{\partial}{\partial t_2} \frac{\partial G_R(\omega, t_2, t_0)}{\partial t_2}$
 521 as follows. We use the result in Eq. 46 and we can interchange the order of partial differentiation and
 522 integration in Eq. 47 using theorem of dominated convergence as follows.

$$\begin{aligned} \frac{\partial^2 G_R(\omega, t_2, t_0)}{\partial t_2^2} &= e^{-2\sigma t_0} \int_{-\infty}^0 \frac{\partial^2 (E'_0(\tau + t_0, t_2) e^{-2\sigma\tau} + E'_{0n}(\tau - t_0, t_2))}{\partial t_2^2} \cos(\omega\tau) d\tau \\ &\quad + e^{2\sigma t_0} \int_{-\infty}^0 \frac{\partial^2 (E'_0(\tau - t_0, t_2) e^{-2\sigma\tau} + E'_{0n}(\tau + t_0, t_2))}{\partial t_2^2} \cos(\omega\tau) d\tau \end{aligned}$$

523

(47)

524 We can use the above procedure in Eq. 45 to Eq. 46 for the term $\frac{\partial^2 (E'_0(\tau+t_0, t_2) e^{-2\sigma\tau} + E'_{0n}(\tau-t_0, t_2))}{\partial t_2^2} =$
 525 $\frac{\partial I(\tau, t_0, t_2)}{\partial t_2}$ where $I(\tau, t_0, t_2) = \frac{\partial(E'_0(\tau+t_0, t_2) e^{-2\sigma\tau} + E'_{0n}(\tau-t_0, t_2))}{\partial t_2}$ in the first integral in Eq. 47 and we can
 526 show that it converges, using the procedure used in Eq. 46 twice.

527

528 We can see that the second integral in Eq. 47 converge, by setting $t_0 = -t_0$ and using the proce-
 529 dure in this section. Hence all the integrals in Eq. 43 and Eq. 47 converge.

530

531 5. Order of $\omega_z(t_2, t_0)t_0$ is greater than $O[1]$

532

533 It is noted that we **do not** use $\lim_{t_0 \rightarrow \infty}$ in this section. Instead we consider real $t_0 > 0$ which
 534 increases to a larger and larger finite value without bounds.

535

536 We write $P_{odd}(t_2, t_0)$ in Eq. 17 concisely as follows.

$$\begin{aligned} P_{odd}(t_2, t_0) &= \int_{-\infty}^{t_0} E'_0(\tau, t_2) e^{-2\sigma\tau} \cos(\omega_z(t_2, t_0)(\tau - t_0)) d\tau + e^{2\sigma t_0} \int_{-\infty}^{t_0} E'_{0n}(\tau, t_2) \cos(\omega_z(t_2, t_0)(\tau - t_0)) d\tau \\ &\quad P_{odd}(t_2, t_0) + P_{odd}(t_2, -t_0) = 0 \end{aligned}$$

537

(48)

538 We note that $E'_{0n}(\tau, t_2) = E'_0(-\tau, t_2) = -E'_0(\tau, t_2) = E_0(\tau + t_2) - E_0(\tau - t_2)$. We choose $t_2 = 2t_0$
 539 and we choose t_1 such that $E_0(t)$ approximates zero for $|t| > t_1$ and we choose $t_0 \gg t_1$ and hence
 540 $E_0(\tau - t_2) = E_0(\tau - 2t_0)$ approximates zero in the interval $[-\infty, t_0]$. Hence in the interval $[-\infty, t_0]$,
 541 we see that $E'_0(\tau, t_2) \approx -E_0(\tau + t_2)$ and $E'_{0n}(\tau, t_2) \approx E_0(\tau + t_2)$, for sufficiently large t_0 .

542

543 We see that the term $P_{odd}(t_2, -t_0)$ approaches a value very close to zero, as real t_0 increases to a
 544 larger and larger finite value without bounds, due to the terms $e^{-2\sigma t_0}$ and the integrals $\int_{-\infty}^{-t_0}$. Hence
 545 we can write as follows.

$$Q(t_0) = P_{odd}(t_2, t_0) + P_{odd}(t_2, -t_0) \approx - \int_{-\infty}^{t_0} E_0(\tau + 2t_0) e^{-2\sigma\tau} \cos(\omega_z(t_2, t_0)(\tau - t_0)) d\tau \\ + e^{2\sigma t_0} \int_{-\infty}^{t_0} E_0(\tau + 2t_0) \cos(\omega_z(t_2, t_0)(\tau - t_0)) d\tau \approx 0$$

(49)

We substitute $\tau + 2t_0 = t$ and write as follows.

$$Q(t_0) \approx -e^{4\sigma t_0} \int_{-\infty}^{3t_0} E_0(t) e^{-2\sigma t} \cos(\omega_z(t_2, t_0)(t - 3t_0)) dt \\ + e^{2\sigma t_0} \int_{-\infty}^{3t_0} E_0(t) \cos(\omega_z(t_2, t_0)(t - 3t_0)) dt \approx 0$$

(50)

We multiply above equation by $e^{-3\sigma t_0}$ and ignore the last integral for sufficiently large t_0 , given that $|\int_{-\infty}^{3t_0} E_0(t) \cos(\omega_z(t_2, t_0)(t - 3t_0)) dt| \leq \int_{-\infty}^{3t_0} |E_0(t)| dt$ is finite.

$$Q(t_0) \approx -e^{\sigma t_0} \int_{-\infty}^{3t_0} E_0(t) e^{-2\sigma t} \cos(\omega_z(t_2, t_0)(t - 3t_0)) dt = -e^{\sigma t_0} R(t_0) \approx 0 \\ R(t_0) = \cos(\omega_z(t_2, t_0)3t_0) \int_{-\infty}^{3t_0} E_0(t) e^{-2\sigma t} \cos(\omega_z(t_2, t_0)t) dt + \sin(\omega_z(t_2, t_0)3t_0) \int_{-\infty}^{3t_0} E_0(t) e^{-2\sigma t} \sin(\omega_z(t_2, t_0)t) dt$$

(51)

Case 1: Order of $\omega_z(t_2, t_0)t_0$ less than 1

Let us assume that the order of $\omega_z(t_2, t_0)t_0$ is less than 1 and $\omega_z(t_2, t_0)t_0$ decreases to a very small finite value close to zero, as real t_0 increases to a larger and larger finite value without bounds. **(Statement B)** We see that t_0 is a real number and as it increases to a larger and larger finite value without bounds, we can use the approximations $\cos(\omega_z(t_2, t_0)t_0) \approx 1$, $\sin(\omega_z(t_2, t_0)t_0) \approx \omega_z(t_2, t_0)t_0 \approx 0$. We see that $\cos(\omega_z(t_2, t_0)t)$ and $\sin(\omega_z(t_2, t_0)t)$ are finite and the integrals in the expression for $Q(t_0)$ in Eq. 51 converge to a finite value.

We choose t_3 such that $E_0(t)e^{-2\sigma t}$ approximates zero for $|t| > t_3$. As t_0 increase without bounds, we see that $t_3 \ll t_0$ and in the interval $[-t_3, t_3]$, we see that the term $\cos(\omega_z(t_2, t_0)t) \approx 1$. Hence we can write Eq. 51 as follows.

$$R(t_0) \approx \int_{-\infty}^{3t_0} E_0(t) e^{-2\sigma t} dt$$

(52)

For sufficiently large t_0 , the integral $R(t_0) \approx \int_{-\infty}^{3t_0} E_0(t) e^{-2\sigma t} dt$ remains finite and non-zero and **does not** approach zero exponentially, as real t_0 increases to a larger and larger finite value without

567 bounds, given that $\int_{-\infty}^{\infty} E_0(t)e^{-2\sigma t}dt > 0$. (Appendix B.1) This is explained in detail in Section 5.1.

568

569 The term $e^{\sigma t_0}$ in $Q(t_0)$ in Eq. 51 increases to a larger and larger finite value **exponentially** and
 570 hence the term $Q(t_0)$ approaches a larger and larger finite value exponentially and hence $Q(t_0)$ and
 571 $P_{odd}(t_2, t_0) + P_{odd}(t_2, -t_0)$ **cannot** equal zero in this case.

572

573 Hence **Statement B** is **false** and $\omega_z(t_2, t_0)t_0$ **does not** decrease towards zero, as finite t_0 increases
 574 without bounds. Given that $\omega_z(t_2, t_0)$ is a **continuous** function of variable t_0 and t_2 , for all $|t_0| < \infty$
 575 and $|t_2| < \infty$ (Section 4), we see that the the order of $\omega_z(t_2, t_0)t_0$ is greater than or equal to 1, as
 576 finite t_0 increases without bounds.

577

578 **Case 2: Order of $\omega_z(t_2, t_0)t_0$ is 1**

579

580 Let us assume that the order of $\omega_z(t_2, t_0)t_0$ is 1, as real t_0 increases to a larger and larger finite
 581 value without bounds. (**Statement C**). In this case, the order of $\omega_z(t_2, t_0)$ is $O[\frac{1}{t_0}]$ and we consider
 582 $\omega_z(t_2, t_0) = \frac{K}{t_0}$ where $K < \frac{\pi}{2}$.

583

584 We choose t_3 such that $Kt_3 < t_0$ and $E_0(t)e^{-2\sigma t}$ is vanishingly small and approximates zero for
 585 $|t| > t_3$. As t_0 increase without bounds, in the interval $[-t_3, t_3]$, we see that the term $\cos(\omega_z(t_2, t_0)t) \approx$
 586 1 and $\sin(\omega_z(t_2, t_0)t) \approx \omega_z(t_2, t_0)t \approx 0$, given that $\omega_z(t_2, t_0)t = \frac{Kt_3}{t_0} < 1$. Hence we can write Eq. 51
 587 as follows.

$$R(t_0) \approx \cos(\omega_z(t_2, t_0)3t_0) \int_{-\infty}^{3t_0} E_0(t)e^{-2\sigma t}dt \quad (53)$$

588 For sufficiently large t_0 , the integral $R(t_0) \approx \cos(\omega_z(t_2, t_0)3t_0) \int_{-\infty}^{3t_0} E_0(t)e^{-2\sigma t}dt$ remains finite,
 589 because the order of $\cos(\omega_z(t_2, t_0)3t_0)$ is 1 and $\int_{-\infty}^{\infty} E_0(t)e^{-2\sigma t}dt > 0$ (Appendix B.1) and **does not**
 590 approach zero exponentially, as real t_0 increases to a larger and larger finite value without bounds.
 591 This is explained in detail in Section 5.1.

592

593 The term $e^{\sigma t_0}$ in $Q(t_0)$ in Eq. 51 increases to a larger and larger finite value **exponentially** and
 594 hence the term $Q(t_0)$ approaches a larger and larger finite value exponentially and hence $Q(t_0)$ and
 595 $P_{odd}(t_2, t_0) + P_{odd}(t_2, -t_0)$ **cannot** equal zero in this case.

596

597 Hence **Statement C** is **false** and the order of $\omega_z(t_2, t_0)t_0$ is **not** 1, as finite t_0 increases without
 598 bounds. Given that $\omega_z(t_2, t_0)$ is a **continuous** function of variable t_0 and t_2 , for all $|t_0| < \infty$ and
 599 $|t_2| < \infty$ (Section 4), we see that the the order of $\omega_z(t_2, t_0)t_0$ is **greater than** 1, as finite t_0 increases
 600 without bounds.

601

602 5.1. $A(t_0) = \int_{-\infty}^{3t_0} E_0(t)e^{-2\sigma t} \cos(\omega_z(t_2, t_0)t)dt$ **does not have exponential fall off rate**

603

604 In this section, we compute the minimum value of the integral $A(t_0) = \int_{-\infty}^{3t_0} E_0(t)e^{-2\sigma t} \cos(\omega_z(t_2, t_0)t)dt$
 605 for sufficiently large t_3 and $t_0 \gg t_3$ and $0 < \sigma < \frac{1}{2}$. We split $A(t_0)$ as follows.

$$\begin{aligned}
A(t_0) &= A_1(t_0) + A_2(t_0) + A_3(t_0) \\
A_1(t_0) &= \int_{-\infty}^{-t_3} E_0(t) e^{-2\sigma t} \cos(\omega_z(t_2, t_0)t) dt, \quad A_2(t_0) = \int_{-t_3}^{t_3} E_0(t) e^{-2\sigma t} \cos(\omega_z(t_2, t_0)t) dt \\
A_3(t_0) &= \int_{t_3}^{3t_0} E_0(t) e^{-2\sigma t} \cos(\omega_z(t_2, t_0)t) dt
\end{aligned}$$

(54)

We derive $A(t_0) \geq K_0 - K_1 - K_2$ where K_0 is the minimum value of $A_2(t_0)$ and K_1 is the maximum value of $A_3(t_0)$ and K_2 is the maximum value of $A_1(t_0)$.

We choose $t_3 = 10$ such that $E_0(t)e^{-2\sigma t}$ is vanishingly small and approximates zero for $|t| > t_3$. Given that $E_0(t) > 0$ for $|t| < \infty$ (Appendix B.1), for $0 < \sigma < \frac{1}{2}$, we see that the integral $\int_{-t_3}^{t_3} E_0(t) e^{-2\sigma t} dt > 2 \int_0^{t_3} E_0(t) e^{-|t|} dt > K_{00} = 0.1055$ where K_{00} is computed by considering the first 5 terms $n = 1, 2, 3, 4, 5$ in $E_0(t) = \sum_{n=1}^{\infty} [4\pi^2 n^4 e^{4t} - 6\pi n^2 e^{2t}] e^{-\pi n^2 e^{2t}} e^{\frac{t}{2}}$.

Given that $\omega_z(t_2, t_0) = \frac{K}{t_0}$ where $K < \frac{\pi}{2}$ and $t_0 \gg t_3$, we see that $\omega_z(t_2, t_0)t \leq \frac{Kt_3}{t_0} \approx 0$ in the interval $|t| \leq t_3$ and hence $\cos(\omega_z(t_2, t_0)t) > \frac{1}{2}$ because $\cos(\omega_z(t_2, t_0)t) \approx 1$ in the interval $|t| \leq t_3$. Hence we can write $A_2(t_0) = \int_{-t_3}^{t_3} E_0(t) e^{-2\sigma t} \cos(\omega_z(t_2, t_0)t) dt > \frac{K_{00}}{2} = K_0 = 0.05275$.

Next we consider the integral $A_3(t_0) = \int_{t_3}^{3t_0} E_0(t) e^{-2\sigma t} \cos(\omega_z(t_2, t_0)t) dt \leq \int_{t_3}^{3t_0} E_0(t) e^{-2\sigma t} dt < \int_{t_3}^{\infty} E_0(t) e^{-2\sigma t} dt < \int_{t_3}^{\infty} E_0(t) dt = K_{10}$. We see that $E_0(t)$ has a fall-off rate of $e^{-\pi e^{2t}} e^{\frac{5t}{2}} > e^{-\pi} e^{-\pi 2t} e^{\frac{5t}{2}} > e^{-\pi} e^{-(2\pi - \frac{5}{2})t} > e^{-\pi} e^{-3.5t}$ which is higher than a minimum fall-off rate of e^{-2t} . Hence we can write $K_{10} < E_0(t_3) e^{2t_3} \int_{t_3}^{\infty} e^{-2t} dt = -\frac{1}{2} E_0(t_3) e^{2t_3} [e^{-2t}]_{t_3}^{\infty} = E_0(t_3) \frac{1}{2} e^{2t_3} e^{-2t_3} = E_0(t_3) \frac{1}{2} = K_1$. For $t_3 = 10$, we see that $K_1 = E_0(t_3) \frac{1}{2} < 1$, given that $E_0(0) < 0.5$ and $E_0(t)$ is a strictly decreasing function for $t > 0$. (Section 6)

Similarly, we see that $A_1(t_0) = \int_{-\infty}^{-t_3} E_0(t) e^{-2\sigma t} \cos(\omega_z(t_2, t_0)t) dt = \int_{t_3}^{\infty} E_0(t) e^{2\sigma t} \cos(\omega_z(t_2, t_0)t) dt \leq \int_{t_3}^{\infty} E_0(t) e^t dt = K_{20}$. We see that $E_0(t)$ has a minimum fall-off rate of e^{-2t} . Hence we can write $K_{20} < E_0(t_3) e^{2t_3} \int_{t_3}^{\infty} e^{-t} dt = -E_0(t_3) e^{2t_3} [e^{-t}]_{t_3}^{\infty} = E_0(t_3) e^{t_3} = K_2$. For $t_3 = 10$, we see that $K_2 = E_0(t_3) e^{t_3} < 1$, given that $E_0(0) < 0.5$ and $E_0(t)$ is a strictly decreasing function for $t > 0$ (Section 6).

Hence we see that $A(t_0) = \int_{-\infty}^{3t_0} E_0(t) e^{-2\sigma t} \cos(\omega_z(t_2, t_0)t) dt > K_0 - K_1 - K_2 = 0.05275 - K_1 - K_2 \approx 0.05275$. As t_0 increases without bounds, we see that $A(t_0) > 0.05275$ and **does not** have exponential fall off rate.

6. Strictly decreasing $E_0(t)$ for $t > 0$

Let us consider $E_0(t) = \Phi(t) = \sum_{n=1}^{\infty} [4\pi^2 n^4 e^{4t} - 6\pi n^2 e^{2t}] e^{-\pi n^2 e^{2t}} e^{\frac{t}{2}}$ whose Fourier Transform is given by the entire function $E_{0\omega}(\omega) = \xi(\frac{1}{2} + i\omega)$. It is known that $\Phi(t)$ is positive for $|t| < \infty$ and its first derivative is negative for $t > 0$ and hence $\Phi(t)$ is a **strictly decreasing** function for $t > 0$ (link). This is shown below.

$$E_0(t) = \Phi(t) = \sum_{n=1}^{\infty} [4\pi^2 n^4 e^{4t} - 6\pi n^2 e^{2t}] e^{-\pi n^2 e^{2t}} e^{\frac{t}{2}} = \sum_{n=1}^{\infty} [4\pi^2 n^4 e^{4t} - 6\pi n^2 e^{2t}] e^{-\pi n^2 e^{2t}} e^{\frac{t}{2}}$$

$$E_0(t) = \sum_{n=1}^{\infty} 2\pi n^2 e^{-\pi n^2 e^{2t}} e^{\frac{t}{2}} [2\pi n^2 e^{4t} - 3e^{2t}]$$

(55)

We show that $X(t) = \frac{E_0(t)}{2}$ is a **strictly decreasing** function for $t > 0$ as follows.

• In Section 6.1, it is shown that the first derivative of $X(t)$, given by $\frac{dX(t)}{dt} < 0$ for $t > t_z$ where $t_z = \frac{1}{2} \log \frac{y_z}{\pi}$ and $y_z = 3.16$.

• In Section 6.2, it is shown that, $\frac{dX(t)}{dt} < 0$ for $0 < t \leq t_z$.

Hence $\frac{dX(t)}{dt} < 0$ for all $t > 0$ and hence $X(t)$ is strictly decreasing for all $t > 0$ and $E_0(t) = 2X(t)$ is **strictly decreasing** for all $t > 0$.

6.1. $\frac{dX(t)}{dt} < 0$ **for** $t > t_z$

We consider $X(t) = \frac{E_0(t)}{2} = \sum_{n=1}^{\infty} \pi n^2 e^{-\pi n^2 e^{2t}} e^{\frac{t}{2}} [2\pi n^2 e^{4t} - 3e^{2t}]$ and take the first derivative of $X(t)$ as follows. We note that $E_0(t)$ is an analytic function for $|t| \leq \infty$ and is infinitely differentiable in that interval.

$$\frac{dX(t)}{dt} = \sum_{n=1}^{\infty} \pi n^2 e^{-\pi n^2 e^{2t}} e^{\frac{t}{2}} [8\pi n^2 e^{4t} - 6e^{2t} + (2\pi n^2 e^{4t} - 3e^{2t}) (\frac{1}{2} - 2\pi n^2 e^{2t})]$$

$$\frac{dX(t)}{dt} = \sum_{n=1}^{\infty} \pi n^2 e^{-\pi n^2 e^{2t}} e^{\frac{t}{2}} [8\pi n^2 e^{4t} - 6e^{2t} + (\pi n^2 e^{4t} - \frac{3}{2}e^{2t} - 4\pi^2 n^4 e^{6t} + 6\pi n^2 e^{4t})]$$

$$\frac{dX(t)}{dt} = \sum_{n=1}^{\infty} \pi n^2 e^{-\pi n^2 e^{2t}} e^{\frac{t}{2}} [-4\pi^2 n^4 e^{6t} + 15\pi n^2 e^{4t} - \frac{15}{2}e^{2t}]$$

$$\frac{dX(t)}{dt} = \sum_{n=1}^{\infty} \pi n^2 e^{-\pi n^2 e^{2t}} e^{\frac{t}{2}} e^{2t} [-4\pi^2 n^4 e^{4t} + 15\pi n^2 e^{2t} - \frac{15}{2}]$$

(56)

We substitute $y = \pi e^{2t}$ in Eq. 56 and define $A(y)$ such that $\frac{dX(t)}{dt} = \pi e^{\frac{5t}{2}} A(y)$. [8]

$$A(y) = \sum_{n=1}^{\infty} n^2 e^{-n^2 y} [-4n^4 y^2 + 15n^2 y - \frac{15}{2}]$$

(57)

We see that $A(y) = 0$ at $y = \pi$, given that $\frac{dX(t)}{dt} = 0$ at $t = 0$, because $X(t)$ is an even function of variable t . The quadratic expression $B(y, n) = (-4n^4 y^2 + 15n^2 y - \frac{15}{2})$ in Eq. 57 has roots at

661 $y = \frac{-15n^2 \pm \sqrt{225n^4 - 120n^4}}{-8n^4} = \frac{(15 \pm \sqrt{105})}{8n^2}$. We see that the second derivative of $B(y, n)$ given by $-8n^4$, is
 662 negative for all y and n and hence $B(y, n)$ is a concave down function for each n , which reaches a
 663 maximum at $y = \frac{15}{8n^2}$ and given the dominant term $-4n^4y^2$ in Eq. 57, we see that $B(y, n) < 0$,
 664 for $y > \frac{(15 + \sqrt{105})}{8} > 3.16 = y_z$, for $n \geq 1$ and hence $A(y) < 0$ for $y > y_z$. Hence $\frac{dX(t)}{dt} < 0$ for
 665 $t > \frac{1}{2} \log \frac{y_z}{\pi} = t_z$ (**Result 1**).

666
 667 We show in the next section that $\frac{dX(t)}{dt} < 0$ for $0 < t \leq t_z$. It suffices to show that $\frac{dA(y)}{dy} < 0$ for
 668 $\pi \leq y \leq 3.16$ and hence $A(y) < 0$ for $\pi < y \leq 3.16$, given that $A(y) = 0$ at $y = \pi$.

669 **6.2. $\frac{dX(t)}{dt} < 0$ for $0 < t \leq t_z$**

670
 671 It is shown in this section that $\frac{dA(y)}{dy} < 0$ for $\pi \leq y \leq 3.16$ and hence $A(y) < 0$ for $\pi < y \leq 3.16$
 672 [8], given that $A(y) = 0$ at $y = \pi$. We take the derivative of $A(y)$ in Eq. 57 and take the factor n^2
 673 out of the brackets, as follows.

$$\begin{aligned} \frac{dA(y)}{dy} &= \sum_{n=1}^{\infty} n^2 e^{-n^2 y} [-8n^4 y + 15n^2 + (-4n^4 y^2 + 15n^2 y - \frac{15}{2})(-n^2)] \\ \frac{dA(y)}{dy} &= \sum_{n=1}^{\infty} n^4 e^{-n^2 y} [-8n^2 y + 15 + 4n^4 y^2 - 15n^2 y + \frac{15}{2}] = \sum_{n=1}^{\infty} n^4 e^{-n^2 y} [4n^4 y^2 - 23n^2 y + \frac{45}{2}] \end{aligned}$$

(58)

674
 675 We examine the term $C(y, n) = n^4 e^{-n^2 y} (4n^4 y^2 - 23n^2 y + \frac{45}{2})$ in Eq. 58 in the interval $\pi \leq y \leq 3.16$
 676 and show that $\frac{dA(y)}{dy} = C(y, 1) + \sum_{n=2}^{\infty} C(y, n) < 0$, as follows.

677
 678 For $n = 1$, we see that $C(y, 1) = e^{-y} (4y^2 - 23y + \frac{45}{2}) < 0$ in the interval $\pi \leq y \leq 3.16$. Given
 679 that $3.16 < 4$ and $3.16^2 < 10$ and $\pi > 3$ in the interval $\pi \leq y \leq 3.16$, we see that $C(y, 1) <$
 680 $e^{-3} (4 * 10 - 23 * 3 + \frac{45}{2}) < e^{-3} (40 - 69 + 23) = -6e^{-3} = C_{max}(1)$ where $C_{max}(1)$ is the maximum
 681 value of $C(y, 1)$ in the interval $\pi \leq y \leq 3.16$.

$$C(y, 1) = e^{-y} (4y^2 - 23y + \frac{45}{2}) < -6e^{-3}, \quad \pi \leq y \leq 3.16$$

(59)

682
 683 For $n > 1$, in the interval $\pi \leq y \leq 3.16$, we can write $C(y, n)$ as follows, given that $-23n^2 y + \frac{45}{2} < 0$
 684 and $\pi > 3$ and $3.16^2 < 10$.

$$C(y, n) = n^4 e^{-n^2 y} (4n^4 y^2 - 23n^2 y + \frac{45}{2}) < n^4 e^{-\pi n^2} (4n^4 (3.16)^2) < 40n^8 e^{-\pi n^2} < 40n^8 e^{-3n^2}$$

(60)

685
 686 We want to show that $\frac{dA(y)}{dy} = C(y, 1) + \sum_{n=2}^{\infty} C(y, n) < 0$ in the interval $\pi \leq y \leq 3.16$. Using
 687 Eq. 59 and Eq. 60, we write

$$\begin{aligned}\frac{dA(y)}{dy} &= C(y, 1) + \sum_{n=2}^{\infty} C(y, n) < -6e^{-3} + \sum_{n=2}^{\infty} 40n^8 e^{-3n^2} \\ e^3 \frac{dA(y)}{dy} &< -6 + \sum_{n=2}^{\infty} 40n^8 e^{3-3n^2}\end{aligned}$$

688

(61)

689 We want to show that $e^3 \frac{dA(y)}{dy} < 0$ in the interval $\pi \leq y \leq 3.16$. We compute $\log(n^8 e^{3-3n^2})$ as
 690 follows. We note that $f(x) = \log x$ is a concave down function whose second derivative given by
 691 $-\frac{1}{x^2} < 0$ for $|x| < \infty$ and we can write $f(x) = \log x \leq f(x_0) + f'(x_0)(x - x_0)$ using its tangent line
 692 equation. We set $x = n$ and $x_0 = 2$ below.

$$\begin{aligned}\log(n^8 e^{3-3n^2}) &= 8 \log n + (3 - 3n^2) \leq 8(\log 2 + \frac{1}{2}(n - 2)) + (3 - 3n^2) \\ \log(n^8 e^{3-3n^2}) &\leq 8 \log 2 + 4n - 5 - 3n^2\end{aligned}$$

693

(62)

694 We note that $g(x) = 4x - 5 - 3x^2$ in Eq. 62 is a concave down function whose second derivative
 695 given by $-6 < 0$ for all x and we can write $g(x) \leq g(x_0) + g'(x_0)(x - x_0)$ using its tangent line
 696 equation. We set $x = n$ and $x_0 = 2$ and write Eq. 62 as follows.

$$\begin{aligned}\log(n^8 e^{3-3n^2}) &\leq 8 \log 2 - 9 - 8(n - 2) \leq 8 \log 2 - 1 + 8(1 - n) \\ n^8 e^{3-3n^2} &\leq 2^8 e^{-1} e^{8(1-n)}\end{aligned}$$

697

(63)

698 We substitute the result in Eq. 63 in Eq. 61 as follows.

$$\begin{aligned}e^3 \frac{dA(y)}{dy} &< -6 + 40 * 2^8 e^{-1} \sum_{n=2}^{\infty} e^{8(1-n)} \\ e^3 \frac{dA(y)}{dy} &< -6 + 40 * 2^8 e^{-1} * e^8 \sum_{n=2}^{\infty} e^{-8n} \\ e^3 \frac{dA(y)}{dy} &< -6 + 40 * 2^8 e^{-1} * e^8 \frac{e^{-8*2}}{1 - e^{-8}} \\ e^3 \frac{dA(y)}{dy} &< -6 + 40 * 2^8 e^{-1} * \frac{e^{-8}}{1 - e^{-8}} \\ e^3 \frac{dA(y)}{dy} &< -6 + 40 * 2^8 e^{-1} * \frac{1}{e^8 - 1}\end{aligned}$$

699

(64)

700 We multiply Eq. 64 by $\frac{(e^8 - 1)}{6}$ and write as follows.

$$e^3 \frac{dA(y)}{dy} \frac{(e^8 - 1)}{6} < -e^8 + 1 + 40e^{-1} * \frac{256}{6} \approx -2352 \quad (65)$$

We see that $e^3 \frac{dA(y)}{dy} \frac{(e^8 - 1)}{6} < 0$ in Eq. 65 and hence $\frac{dA(y)}{dy} < 0$, in the interval $\pi \leq y \leq 3.16$. Given that $A(y) = 0$ at $y = \pi$, we see that $A(y) < 0$ in Eq. 57, for $\pi < y \leq 3.16$ and $\frac{dX(t)}{dt} = \pi e^{\frac{5t}{2}} A(y) < 0$ in the interval $0 < t \leq t_z$. (**Result 2**)

In Section 6.1, it is shown that $\frac{dX(t)}{dt} < 0$ for $t > t_z$ (from Result 1). In this section, we have shown that $\frac{dX(t)}{dt} < 0$ for $0 < t \leq t_z$. Hence $\frac{dX(t)}{dt} < 0$ for all $t > 0$.

Hence $E_0(t) = 2X(t)$ is a **strictly decreasing function** for $t > 0$.

6.3. **Result** $E_0(t - t_{2c}) - E_0(t + t_{2c}) > 0$

It is shown in Section 6 that $E_0(t)$ is **strictly decreasing** for $t > 0$. In this section, it is shown that $E_0(t - t_{2c}) - E_0(t + t_{2c}) > 0$, for $0 < t < t_{0c}$ and $t_{2c} = 2t_{0c}$ in Eq. 34.

Given that $E_0(t)$ is a **strictly decreasing** function for $t > 0$ and $E_0(t)$ is an **even** function of variable t , and $t_{2c} = 2t_{0c}$, we see that, in the interval $0 < t < t_{0c}$, $E_0(t + t_{2c}) = E_0(t + 2t_{0c})$ ranges from $E_0(2t_{0c})$ to $E_0(3t_{0c})$, which is **less than** $E_0(t - t_{2c}) = E_0(t - 2t_{0c})$ which ranges from $E_0(-2t_{0c})$ to $E_0(-t_{0c})$ respectively. Hence we see that $E_0(t - t_{2c}) > E_0(t + t_{2c})$, in the interval $0 < t < t_{0c}$. At $t = 0$, $E_0(t - t_{2c}) = E_0(t + t_{2c})$. At $t = t_{0c}$, $E_0(t - t_{2c}) > E_0(t + t_{2c})$ because $E_0(-t_{0c}) > E_0(3t_{0c})$.

Hence $E_0(t - t_{2c}) - E_0(t + t_{2c}) > 0$ for $0 < t < t_{0c}$ in Eq. 34, for $t_{0c} > 0$ and $t_{2c} = 2t_{0c}$.

7. Hurwitz Zeta Function and related functions

We can show that the new method is **not** applicable to Hurwitz zeta function and related zeta functions and **does not** contradict the existence of their non-trivial zeros away from the critical line with real part of $s = \frac{1}{2}$. The new method requires the **symmetry** relation $\xi(s) = \xi(1 - s)$ and hence $\xi(\frac{1}{2} + i\omega) = \xi(\frac{1}{2} - i\omega)$ when evaluated at the critical line $s = \frac{1}{2} + i\omega$. This means $\xi(\frac{1}{2} + i\omega) = E_{0\omega}(\omega) = E_{0\omega}(-\omega)$ and $E_0(t) = E_0(-t)$ where $E_0(t) = \sum_{n=1}^{\infty} [4\pi^2 n^4 e^{4t} - 6\pi n^2 e^{2t}] e^{-\pi n^2 e^{2t}} e^{\frac{t}{2}}$ and this condition is satisfied for Riemann's Zeta function.

It is **not** known that Hurwitz Zeta Function given by $\zeta(s, a) = \sum_{m=0}^{\infty} \frac{1}{(m+a)^s}$ satisfies a symmetry relation similar to $\xi(s) = \xi(1 - s)$ where $\xi(s)$ is an entire function, for $a \neq 1$ and hence the condition $E_0(t) = E_0(-t)$ is **not** known to be satisfied^[6]. Hence the new method is **not** applicable to Hurwitz zeta function and **does not** contradict the existence of their non-trivial zeros away from the critical line.

Dirichlet L-functions satisfy a symmetry relation $\xi(s, \chi) = \epsilon(\chi) \xi(1 - s, \bar{\chi})$ ^[7] which does **not** translate to $E_0(t) = E_0(-t)$ required by the new method and hence this proof is **not** applicable to them. This proof does not need or use Euler product.

We know that $\zeta(s) = \sum_{m=1}^{\infty} \frac{1}{m^s}$ diverges for real part of $s \leq 1$. Hence we derive a convergent and

entire function $\xi(s)$ using the well known theorem $F(x) = 1 + 2 \sum_{n=1}^{\infty} e^{-\pi n^2 x} = \frac{1}{\sqrt{x}} (1 + 2 \sum_{n=1}^{\infty} e^{-\pi \frac{n^2}{x}})$,

where $x > 0$ is real^[4] and then derive $E_0(t) = \sum_{n=1}^{\infty} [4\pi^2 n^4 e^{4t} - 6\pi n^2 e^{2t}] e^{-\pi n^2 e^{2t}} e^{\frac{t}{2}}$ (link). In the case of **Hurwitz zeta** function and **other zeta functions** with non-trivial zeros away from the critical line, it is **not** known if a corresponding relation similar to $F(x)$ exists, which enables derivation of a convergent and entire function $\xi(s)$ and results in $E_0(t)$ as a Fourier transformable, real, even and analytic function. Hence the new method presented in this paper is **not** applicable to Hurwitz zeta function and related zeta functions.

The proof of Riemann Hypothesis presented in this paper is **only** for the specific case of Riemann's Zeta function and **only** for the **critical strip** $0 \leq |\sigma| < \frac{1}{2}$. This proof requires both $E_p(t)$ and $E_{p\omega}(\omega)$ to be Fourier transformable where $E_p(t) = E_0(t)e^{-\sigma t}$ is a real analytic function and uses the fact that $E_0(t)$ is an **even** function of variable t and $\int_{-\infty}^{\infty} E_0(t)dt > 0$ for $|t| < \infty$ (Appendix B.1) and $E_0(t)$ is **strictly decreasing** function for $t > 0$ (Section 6). These conditions may **not** be satisfied for many other functions including those which have non-trivial zeros away from the critical line and hence the new method may **not** be applicable to such functions.

References

- [1] Bernhard Riemann, On the Number of Prime Numbers less than a Given Quantity.(Ueber die Anzahl der Primzahlen untereiner gegebenen Grosse.) Monatsberichte der Berliner Akademie, November 1859. (Link to Riemann's 1859 paper)
- [2] Hardy, G.H., Littlewood, J.E. The zeros of Riemann's zeta-function on the critical line. Mathematische Zeitschrift volume 10, pp.283 to 317 (1921).
- [3] E. C. Titchmarsh, The Theory of the Riemann Zeta Function. (1986) pp.254 to 255
- [4] Fern Ellison and William J. Ellison, Prime Numbers (1985).pp147 to 152
- [5] J. Brian Conrey, The Riemann Hypothesis (2003). (Link to Brian Conrey's 2003 article)
- [6] Mathworld article on Hurwitz Zeta functions. (Link)
- [7] Wikipedia article on Dirichlet L-functions. (Link)
- [8] Thomas Browning. draft article

Appendix A. Derivation of $E_p(t)$

Let us start with Riemann's Xi Function $\xi(s)$ evaluated at $s = \frac{1}{2} + i\omega$ given by $\xi(\frac{1}{2} + i\omega) = E_{0\omega}(\omega)$. Its inverse Fourier Transform is given by $E_0(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} E_{0\omega}(\omega) e^{i\omega t} d\omega = \sum_{n=1}^{\infty} [4\pi^2 n^4 e^{4t} - 6\pi n^2 e^{2t}] e^{-\pi n^2 e^{2t}} e^{\frac{t}{2}}$ (link). This is re-derived in link.

We will show in this section that the inverse Fourier Transform of the function $\xi(\frac{1}{2} + \sigma + i\omega) = E_{p\omega}(\omega)$, is given by $E_p(t) = E_0(t)e^{-\sigma t}$ where $0 \leq |\sigma| < \frac{1}{2}$ is real.

$$\begin{aligned}\xi\left(\frac{1}{2} + \sigma + i\omega\right) &= \xi\left(\frac{1}{2} + i(\omega - i\sigma)\right) = E_{p\omega}(\omega) = E_{0\omega}(\omega - i\sigma) \\ E_p(t) &= \frac{1}{2\pi} \int_{-\infty}^{\infty} E_{p\omega}(\omega) e^{i\omega t} d\omega = \frac{1}{2\pi} \int_{-\infty}^{\infty} E_{0\omega}(\omega - i\sigma) e^{i\omega t} d\omega\end{aligned}$$

(A.1)

We substitute $\omega' = \omega - i\sigma$ in Eq. A.1 as follows.

$$E_p(t) = e^{-\sigma t} \frac{1}{2\pi} \int_{-\infty - i\sigma}^{\infty - i\sigma} E_{0\omega}(\omega') e^{i\omega' t} d\omega' \quad (\text{A.2})$$

We can evaluate the above integral in the complex plane using contour integration, substituting $\omega' = z = x + iy$ and we use a rectangular contour comprised of C_1 along the line $x = [-\infty, \infty]$, C_2 along the line $y = [\infty, \infty - i\sigma]$, C_3 along the line $x = [\infty - i\sigma, -\infty - i\sigma]$ and then C_4 along the line $y = [-\infty - i\sigma, -\infty]$. We can see that $E_{0\omega}(z) = \xi(\frac{1}{2} + iz)$ has no singularities in the region bounded by the contour because $\xi(\frac{1}{2} + iz)$ is an entire function in the Z-plane.

In **Appendix B.1**, we show that $\int_{-\infty}^{\infty} |E_p(t)| dt$ is finite and $E_p(t) = E_0(t) e^{-\sigma t}$ is an absolutely integrable function, for $0 \leq |\sigma| < \frac{1}{2}$.

We use the fact that $E_{0\omega}(z) = \xi(\frac{1}{2} + iz) = \xi(\frac{1}{2} - y + ix) = \int_{-\infty}^{\infty} E_0(t) e^{-izt} dt = \int_{-\infty}^{\infty} E_0(t) e^{yt} e^{-ixt} dt$, **goes to zero** as $x \rightarrow \pm\infty$ when $-\sigma \leq y \leq 0$, as per Riemann-Lebesgue Lemma (link), because $E_0(t) e^{yt}$ is a absolutely integrable function in the interval $-\infty \leq t \leq \infty$. Hence the integral in Eq. A.2 **vanishes** along the contours C_2 and C_4 . Using Cauchy's Integral theroem, we can write Eq. A.2 as follows.

$$\begin{aligned}E_p(t) &= e^{-\sigma t} \frac{1}{2\pi} \int_{-\infty}^{\infty} E_{0\omega}(\omega') e^{i\omega' t} d\omega' \\ E_p(t) &= E_0(t) e^{-\sigma t} = \sum_{n=1}^{\infty} [4\pi^2 n^4 e^{4t} - 6\pi n^2 e^{2t}] e^{-\pi n^2 e^{2t}} e^{\frac{t}{2}} e^{-\sigma t}\end{aligned}$$

(A.3)

Thus we have arrived at the desired result $E_p(t) = E_0(t) e^{-\sigma t}$.

Appendix B. Properties of Fourier Transforms

Appendix B.1. $E_p(t), h(t)$ are absolutely integrable functions and their Fourier Transforms are finite.

The inverse Fourier Transform of the function $E_{p\omega}(\omega) = \xi(\frac{1}{2} + \sigma + i\omega)$ is given by $E_p(t) = E_0(t) e^{-\sigma t} = \frac{1}{2\pi} \int_{-\infty}^{\infty} E_{p\omega}(\omega) e^{i\omega t} d\omega$. We see that $E_0(t) = \sum_{n=1}^{\infty} [4\pi^2 n^4 e^{4t} - 6\pi n^2 e^{2t}] e^{-\pi n^2 e^{2t}} e^{\frac{t}{2}} > 0$ for all $0 \leq t < \infty$. Given that $E_0(t) = E_0(-t)$, we see that $E_0(t) > 0$ and $E_p(t) = E_0(t) e^{-\sigma t} > 0$ for all $-\infty < t < \infty$.

As $t \rightarrow \infty$, $E_p(t)$ goes to zero, due to the term $e^{-\pi n^2 e^{2t}}$. As $t \rightarrow -\infty$, $E_p(t)$ goes to zero, because for every value of n , the term $e^{-\pi n^2 e^{2t}} e^{\frac{5t}{2}} e^{-\sigma t}$ goes to zero, for $0 \leq |\sigma| < \frac{1}{2}$. Hence $E_p(t) = E_0(t)e^{-\sigma t}$ goes to zero, at $t \rightarrow \pm\infty$ and we showed that $E_p(t) > 0$ for all $-\infty < t < \infty$. Hence $E_{p\omega}(\omega) = \int_{-\infty}^{\infty} E_p(t)e^{-i\omega t}dt$, evaluated at $\omega = 0$ **cannot** be zero. Hence $E_{p\omega}(\omega)$ **does not have a zero** at $\omega = 0$ and hence $\omega_0 \neq 0$.

Given that $\xi(\frac{1}{2} + \sigma + i\omega) = E_{p\omega}(\omega)$ is an entire function in the whole of s-plane, it is finite for $|\omega| \leq \infty$ and also for $\omega = 0$. Hence $\int_{-\infty}^{\infty} E_p(t)dt$ is finite. We see that $E_p(t) \geq 0$ for all $|t| \leq \infty$. Hence we can write $\int_{-\infty}^{\infty} |E_p(t)|dt$ is finite and $E_p(t)$ is an absolutely **integrable function** and its Fourier transform $E_{p\omega}(\omega)$ goes to zero as $\omega \rightarrow \pm\infty$, as per Riemann Lebesgue Lemma (link).

Using the arguments in above paragraph, we see that $E_0(t)$ and $E_0(t)e^{-2\sigma t}$ are absolutely **integrable** functions and the integrals $\int_{-\infty}^{\infty} |E_0(t)|dt < \infty$ and $\int_{-\infty}^{\infty} |E_0(t)e^{-2\sigma t}|dt < \infty$.

We can see that $h(t) = e^{\sigma t}u(-t) + e^{-\sigma t}u(t)$ is an absolutely **integrable function** because $\int_{-\infty}^{\infty} |h(t)|dt = \int_{-\infty}^{\infty} h(t)dt = [\int_{-\infty}^{\infty} h(t)e^{-i\omega t}dt]_{\omega=0} = [\frac{1}{\sigma-i\omega} + \frac{1}{\sigma+i\omega}]_{\omega=0} = \frac{2}{\sigma}$, is finite for $0 < \sigma < \frac{1}{2}$ and its Fourier transform $H(\omega)$ goes to zero as $\omega \rightarrow \pm\infty$, as per Riemann Lebesgue Lemma (link).

Appendix B.2. Convolution integral convergence

Let us consider $h(t) = e^{\sigma t}u(-t) + e^{-\sigma t}u(t)$ whose **first derivative is discontinuous** at $t = 0$. The second derivative of $h(t)$ given by $h_2(t)$ has a Dirac delta function $A_0\delta(t)$ where $A_0 = -2\sigma$ and its Fourier transform $H_2(\omega)$ has a constant term A_0 , corresponding to the Dirac delta function.

This means $h(t)$ is obtained by integrating $h_2(t)$ twice and its Fourier transform $H(\omega)$ has a term $-\frac{A_0}{\omega^2}$ (link) and has a **fall off rate** of $\frac{1}{\omega^2}$ as $|\omega| \rightarrow \infty$ and $\int_{-\infty}^{\infty} H(\omega)d\omega$ converges.

Let us consider the function $g(t, t_2, t_0) = f(t, t_2, t_0)e^{-\sigma t}u(-t) + f(t, t_2, t_0)e^{\sigma t}u(t)$. We can see that $G(\omega, t_2, t_0), H(\omega)$ have **fall-off rate** of $\frac{1}{\omega^2}$ as $|\omega| \rightarrow \infty$ because the **first derivatives** of $g(t, t_2, t_0), h(t)$ are **discontinuous** at $t = 0$. Hence the convolution integral below converges to a finite value for $|\omega| \leq \infty$.

$$F(\omega, t_2, t_0) = \frac{1}{2\pi} \int_{-\infty}^{\infty} G(\omega', t_2, t_0)H(\omega - \omega')d\omega' = \frac{1}{2\pi} [G(\omega, t_2, t_0) * H(\omega)] \quad (\text{B.1})$$

Appendix B.3. Fall off rate of Fourier Transform of functions

Let us consider a real Fourier transformable function $P(t) = P_+(t)u(t) + P_-(t)u(-t)$ whose $(N-1)^{th}$ **derivative is discontinuous** at $t = 0$. The $(N)^{th}$ derivative of $P(t)$ given by $P_N(t)$ has a Dirac delta function $A_0\delta(t)$ where $A_0 = [\frac{d^{N-1}P_+(t)}{dt^{N-1}} - \frac{d^{N-1}P_-(t)}{dt^{N-1}}]_{t=0}$ and its Fourier transform $P_N(\omega)$ has a constant term A_0 , corresponding to the Dirac delta function.

This means $P(t)$ is obtained by integrating $P_N(t)$, N times and its Fourier transform $P(\omega)$ has a term $\frac{A_0}{(i\omega)^N}$ (link) and has a **fall off rate** of $\frac{1}{\omega^N}$ as $|\omega| \rightarrow \infty$.

We have shown that if the $(N-1)^{th}$ **derivative** of the function $P(t)$ is **discontinuous** at $t = 0$ then its Fourier transform $P(\omega)$ has a **fall-off rate** of $\frac{1}{\omega^N}$ as $|\omega| \rightarrow \infty$.

848 *Appendix B.4. Payley-Weiner theorem and Exponential Fall off rate of analytic func-*
849 *tions.*
850

851 We know that Payley-Weiner theorem relates analytic functions and exponential decay rate of
852 their Fourier transforms (link). Using similar arguments, we will show that the functions $E_0(t)$, $E_p(t)$
853 and $x(t) = E_0(t)e^{-2\sigma t}$ have fall-off rates **at least** $\frac{1}{t^3}$ as $|t| \rightarrow \infty$ for $0 < \sigma < \frac{1}{2}$.
854

855 We know that the order of Riemann's Xi function $\xi(\frac{1}{2} + i\omega) = E_{0\omega}(\omega) = \Xi(\omega)$ is given by
856 $O(\omega^A e^{-\frac{|\omega|\pi}{4}})$ where A is a constant^[3] (link). Hence both $E_{0\omega}(\omega)$ and $E_{p\omega}(\omega) = \xi(\frac{1}{2} + \sigma + i\omega) =$
857 $E_{0\omega}(\omega - i\sigma)$ have **exponential fall-off** rate $O(\omega^A e^{-\frac{|\omega|\pi}{4}})$ as $|\omega| \rightarrow \infty$ and they are absolutely inte-
858 grable and Fourier transformable, given that they are derived from an entire function $\xi(s)$.
859

860 Given that $\xi(s)$ is an entire function in the s-plane, we see that $E_{0\omega}(\omega)$ and $E_{p\omega}(\omega)$ are **analytic**
861 functions which are infinitely differentiable which produce no discontinuities for all $|\omega| \leq \infty$ and
862 $0 < \sigma < \frac{1}{2}$. Hence their respective **inverse Fourier transforms** $E_0(t)$, $E_p(t)$ have fall-off rates faster
863 than $\frac{1}{t^M}$ as $M \rightarrow \infty$, as $|t| \rightarrow \infty$ (Appendix B.3) and hence it should have **exponential fall-off**
864 rates as $|t| \rightarrow \infty$.
865

866 We can use similar arguments to show that $x(t) = E_0(t)e^{-2\sigma t}$ has a fall-off rate of **at least** $\frac{1}{t^3}$ as
867 $|t| \rightarrow \infty$, because its Fourier transform is an **analytic** function for all $|\omega| \leq \infty$ with **exponential**
868 **fall-off** rate $O(\omega^A e^{-\frac{|\omega|\pi}{4}})$ as $|\omega| \rightarrow \infty$.
869

870 We can show that $x(t) = t^r x(t)$ has exponential fall-off rate as $|t| \rightarrow \infty$ for $r = 0, 1, 2$, given that
871 its Fourier transform given by $\frac{d^r X(\omega)}{d\omega^r} (\frac{1}{-i2\pi})^r$ is an analytic function for all $|\omega| \leq \infty$ with **exponential**
872 **fall-off** rate $O(\omega^A e^{-\frac{|\omega|\pi}{4}})$ as $|\omega| \rightarrow \infty$.