

A unified dataset of co-located sewage pollution, periphyton, and benthic macroinvertebrate community and food web structure from Lake Baikal (Siberia)

Michael F. Meyer^{1*}

Ted Ozersky²

Kara H. Woo³

Kirill Shchapov²

Aaron W. E. Galloway⁴

Julie B. Schram⁴

Daniel D. Snow⁵

Maxim A. Timofeyev⁶

Dmitry Yu. Karnaukhov⁶

Matthew R. Brousil³

Stephanie E. Hampton³

¹. School of the Environment, Washington State University, Pullman, WA, USA

². Large Lakes Observatory, University of Minnesota - Duluth, Duluth, MN, USA

³. Center for Environmental Research, Education, and Outreach, Washington State University, Pullman, WA, USA

⁴. Oregon Institute of Marine Biology, University of Oregon, Charleston, OR, USA

⁵. School of Natural Resources, University of Nebraska-Lincoln, Lincoln, NE, USA

⁶. Biological Research Institute, Irkutsk State University, Irkutsk, Irkutsk Oblast, Russia

*corresponding author: michael.f.meyer@wsu.edu

Author Contribution Statement

Conceptualized the project: MFM, SEH, TO

Collected samples in the field: MFM, TO, KHW, SEH

Processed samples: MFM, KS, JBS, DDS, TO, AWEG, SEH

Wrote and Reviewed R scripts: MFM, MRB, KHW

Data management: MFM, MRB

Wrote and edited the manuscript: All authors

Approved the final manuscript: All authors

Grant sponsor information:

Funding was provided by the National Science Foundation (NSF-DEB-1136637) to S.E.H., a Fulbright Fellowship to M.F.M., a NSF Graduate Research Fellowship to M.F.M. (NSF-DGE-1347973), and the Russian Ministry of Science and Education (N FZZE-2020-0026).

Key Words: pharmaceuticals, microplastics, fatty acids, stable isotopes, amphipod, mollusk, diatom, spirogyra

URL of the Dataset and Metadata with permanent identifier:

- Environmental Data Initiative: doi:10.6073/pasta/76f43144015ec795679bac508efa044b
- Open Science Framework: <https://doi.org/10.17605/OSF.IO/9TA8Z>

Code URL with permanent identifier:

- Environmental Data Initiative: doi:10.6073/pasta/76f43144015ec795679bac508efa044b
- Open Science Framework: <https://doi.org/10.17605/OSF.IO/9TA8Z>

Measurement(s): Chlorophyll a, Fatty Acids, Pharmaceuticals and Personal Care Products, Microplastics, Periphyton community abundance, benthic macroinvertebrate abundance, Stable Isotopes, nitrate, ammonium, total phosphorus

Technology Type(s): GC/MS, LC/MS, Spectrophotometry

Temporal range: 19 – 23 August 2015

Frequency or sampling interval: single snapshot in time

Spatial scale: site-based

Abstract (150 of 150 words)

Sewage released from lakeside development can introduce nutrients and micropollutants that can restructure aquatic ecosystems. Lake Baikal, the world's most ancient, biodiverse, and voluminous lake, has been experiencing localized sewage pollution from lakeside settlements.

Nearby increasing filamentous algal abundance suggests benthic communities are responding to this localized pollution. We surveyed 40-km of Lake Baikal's southwestern shoreline 19-23 August 2015 for sewage indicators, including pharmaceuticals, personal care products, and microplastics with co-located periphyton, macroinvertebrate, stable isotope, and fatty acid samplings. Unique identifiers corresponding to sampling locations are retained throughout all data files to facilitate interoperability among the dataset's 150+ variables. The data are structured in a tidy format (a tabular arrangement familiar to limnologists) to encourage reuse. For Lake Baikal studies, these data can support continued monitoring and research efforts. For global studies of lakes, these data can help characterize sewage prevalence and ecological consequences of anthropogenic disturbance across spatial scales.

Background and Motivation

Globally, sewage pollution is a common and often concentrated source of nitrogen and phosphorus inputs that can reshape aquatic ecosystems. Sewage inputs are often associated with increased primary production (Edmondson 1970; Moore et al. 2003), which can eventually lead to nuisance algal blooms (Hall et al. 1999; Lapointe et al. 2015). Even in instances where sewage pollution is mitigated, restoring systems can be complicated and necessitate system-specific (Jeppesen et al. 2005), long-term mitigation strategies (Hall et al. 1999; Tong et al. 2020). As such, effective sewage monitoring can require merging a suite of chemical, biological, and ecological data to synthesize locations and timing of inputs with associated shifts in ecological communities (Rosenberger et al. 2008; Hampton et al. 2011).

Definitively identifying sewage as the source of excess nutrients in a system can be challenging. Nutrients can originate from multiple sources, such as agriculture (Powers et al. 2016) or melting permafrost (Turetsky et al. 2000), which can obfuscate wastewater signals. Unlike nutrients,

sewage-specific indicators, such as enhanced $\delta^{15}\text{N}$ stable isotope signatures (Costanzo et al. 2001; Camilleri and Ozersky 2019), pharmaceuticals and personal care products (PPCPs) (Bendz et al. 2005; Rosi-Marshall and Royer 2012; Meyer et al. 2019) and microplastics (Barnes et al. 2009), can be highly specific to human wastewater. Accordingly, sewage-associated micropollutants have garnered global attention for their usefulness in identifying presence and quantifying magnitude of wastewater inputs. While indicators may accumulate differentially in certain taxa (Gartner et al. 2002; Green 2016; Vendel et al. 2017; Richmond et al. 2018), acutely dangerous concentrations are not common in most aquatic systems (Kolpin et al. 2002; Focazio et al. 2008; Yang et al. 2018). However, chronic exposure to microplastics and PPCPs at even minute concentrations (e.g., $\mu\text{g/L}$) can still disrupt ecological processes (Richmond et al. 2017). For example, oxazepam can increase feeding rate and decrease sociability of river perch (Brodin et al. 2013), and microplastics can release dissolved organic carbon, thereby altering microbial communities (Romera-Castillo et al. 2018). The pervasiveness and diversity of sewage-associated micropollutants in tandem with their potency as ecologically disrupting compounds necessitates investigation within and across systems, thereby enabling synthesis of how micropollutants alter ecosystems.

When assessing biological responses to increased nutrient loading, littoral benthic algal, and macroinvertebrate communities often respond most markedly, as their physical proximity to the shoreline puts them in the path of sewage pollution entering the lake (Rosenberger et al. 2008; Hampton et al. 2011). Filamentous algae, for example, can quickly increase in abundance near sewage sources (Rosenberger et al. 2008; Hampton et al. 2011). As algal communities change, food webs can also restructure. For example, change in algal communities can alter the nutritional value of primary producers or cause changes in the relative abundance of different feeding groups (e.g., increased representation of detritivores). Among the suite of food quality metrics, availability of essential fatty acids (EFAs) offers a nuanced understanding of food quality as primary producers usually maintain a consistent EFA signatures (Taipale et al. 2013) and consumers acquire EFAs by grazing (Dalsgaard et al. 2003) or trophic upgrading (Sargent and Falk-Petersen 1988; Dalsgaard et al. 2003).

Together, food web structure, community composition, and sewage indicator data can be powerful tools to assess biological impacts of sewage pollution. Despite their utility, these data are not often available for many limnological systems. PPCPs, for example, have historically been less measured in lake environments (Meyer et al. 2019). In instances where data are available, efficiently merging disparate data into a single, analytically-friendly format can be challenging and require relatively complex, computationally intensive workflows (Meyer et al. 2020a).

To offer a template for harmonizing sewage indicator and biological data, we present a unified data product, which contains disparate data collected from 14 littoral and 3 pelagic sites at Lake Baikal from 19 through 23 August 2015 (Figure 1). Located in Siberia, Lake Baikal is the oldest, most voluminous, and deepest freshwater lake in the world (Hampton et al. 2018). Lake Baikal also has the global distinction of being the most biodiverse lake, with the highest endemism (Moore et al. 2009). The lake is experiencing rapid warming associated with climate change, including decrease in ice cover duration (Moore et al. 2009), and it exhibits offshore plankton community changes associated with warming (Hampton et al. 2008; Katz et al. 2015; Izmet'seva et al. 2016). Less is known of the change occurring in the nearshore of Lake Baikal, where not only climatic changes (Swann et al. 2020) but also human activity (Timoshkin et al. 2018) alter the environment.

Nearshore change is particularly important to understand in Lake Baikal, since the majority of the lake's biodiversity and endemic species occur in the littoral zone (Kozhova and Izvest'eva 1998). While Lake Baikal's pelagic zone is generally ultra-oligotrophic (Yoshida et al. 2003; O'Donnell et al. 2017), littoral areas abutting lakeside settlements have recently shown distinct signs of eutrophication (Timoshkin et al. 2016; Volkova et al. 2018).

As a means of identifying sewage from small, concentrated lakeside towns and the associated ecological responses, we assembled a dataset consisting of over 150 variables collected at 14 littoral and 3 pelagic sampling sites. We structured the dataset in a tidy format, where each row is a sample, each column is a variable, and each CSV file is an observable unit, where more similar variables are contained within an individual file (Wickham 2014). Independent CSV files can be merged using unique locational identifiers as relational keys, enabling future researchers to customize analyses around a particular suite of variables. As a result of the dataset's interoperability, reproducibility, and extensive variable content, it is well poised for future reuse as supporting evidence of sewage pollution in Lake Baikal. Additionally, the data's flexibility and consistent structure enable it to be merged with similar datasets, so as to synthesize biological responses to sewage across systems and scales.

To our knowledge, no raw data on Lake Baikal macroinvertebrates, periphyton, or nearshore water quality are public in a machine-readable format, for any variable (i.e. abundance, fatty acid content, stable isotopes, nutrient and pollutant concentration), and no georeferenced data on pharmaceuticals and personal care products or microplastics appear to be publicly available for any boreal, subarctic, or arctic lakes or rivers in Siberia. Thus, the dataset fills a substantial gap for future studies, providing a window into nearshore biotic assemblages and water quality in a unique, ancient ecosystem that holds 20% of the world's liquid surface water (Moore et al. 2009).

Data Description

The final, replicate-level data products are available on the Environmental Data Initiative (EDI), where they can be freely accessed without potential barriers such as paywalls or account registrations. The final data are provided as 11 separate CSV files, each structured in a tabular format and containing a "site" column that can be used to merge tables. The repository also contains a compressed folder of R scripts (scripts.tar.gz), which were used in the main analysis of the dataset (Meyer et al., Under Review).

chlorophylla.csv

This file contains chlorophyll a concentrations as well as fluorometric corrections for each littoral and pelagic sampling location.

site

Unique alphanumeric identifier for a sampling location.

replicate

Replicate number.

filtered_volume_ml

Lake water volume filtered in milliliters for a given replicate.

sample_volume_ml

Sample volume filtered for chlorophyll a extraction.

raw_fluo

Raw, uncorrected fluorometric reading for chlorophyll analysis.

adjusted_raw

Corrected fluorometric reading for chlorophyll analysis.

chl_conc

Chlorophyll a concentration in milligrams per liter.

distance_weighted_population_metrics.csv

This file contains human population data for each sampled location. Although the majority of sites do not have adjacent shoreline human developments, we calculated inverse distance weighted (IDW) population for each sampling location. IDW population is a generalized representation of the size of and proximity to a sampling location's neighboring human settlements. A full description of the methods used to calculate IDW population can be found in the companion manuscript Meyer et al.(Under Review).

site

Unique alphanumeric identifier for a sampling location.

distance_weighted_population

Inverse distance weighted population for a given sampling location and estimated as number of people. Because this interpolation process is a function of the size of and proximity to neighboring developed sites, values can contain decimal values.

fatty_acid.csv

This file contains fatty acid concentrations for various benthic macroinvertebrate genera, periphyton, and endemic *Draparnaldia* spp. benthic algae collected from the 14 littoral sampling locations.

site

Unique alphanumeric identifier for a sampling location.

Genus

Genus of the analyzed organism.

Species

Species of the analyzed organism. When organism was identified solely to genus, the Species value is NA.

c12_0

Concentration of 12:0 fatty acid as micrograms of fatty acid per milligram of tissue.

i_14_0

Concentration of i-14:0 fatty acid as micrograms of fatty acid per milligram of tissue.

c14_0

Concentration of 14:0 fatty acid as micrograms of fatty acid per milligram of tissue.

c14_4w5

Concentration of 14:4n-5 fatty acid as micrograms of fatty acid per milligram of tissue.

i_15_0

Concentration of i-15:0 fatty acid as micrograms of fatty acid per milligram of tissue.

a_15_0

Concentration of a-15:0 fatty acid as micrograms of fatty acid per milligram of tissue.

c15_0

Concentration of 15:0 fatty acid as micrograms of fatty acid per milligram of tissue.

c15_1w7

Concentration of 15:1 ω 7 fatty acid as micrograms of fatty acid per milligram of tissue.

i_16_0

Concentration of i-16:0 fatty acid as micrograms of fatty acid per milligram of tissue.

c16_0

Concentration of 16:0 fatty acid as micrograms of fatty acid per milligram of tissue.

c16_1w9

Concentration of 16:1 ω 9 fatty acid as micrograms of fatty acid per milligram of tissue.

c16_1w8

Concentration of 16:1 ω 8 fatty acid as micrograms of fatty acid per milligram of tissue.

c16_1w7

Concentration of 16:1 ω 7 fatty acid as micrograms of fatty acid per milligram of tissue.

c16_1w6

Concentration of 16:1 ω 6 fatty acid as micrograms of fatty acid per milligram of tissue.

c16_1w5

Concentration of 16:1 ω 5 fatty acid as micrograms of fatty acid per milligram of tissue.

i_17_0

Concentration of i-17:0 fatty acid as micrograms of fatty acid per milligram of tissue.

a_17_0

Concentration of a-17:0 fatty acid as micrograms of fatty acid per milligram of tissue.

c17_0

Concentration of 17:0 fatty acid as micrograms of fatty acid per milligram of tissue.

c17_1w7

Concentration of 17:1n-7 fatty acid as micrograms of fatty acid per milligram of tissue.

c16_2w7

Concentration of 16:2 ω 7 fatty acid as micrograms of fatty acid per milligram of tissue.

c16_2w6

Concentration of 16:2 ω 6 fatty acid as micrograms of fatty acid per milligram of tissue.

c16_2w4

Concentration of 16:2 ω 4 fatty acid as micrograms of fatty acid per milligram of tissue.

c16_3w6

Concentration of 16:3 ω 6 fatty acid as micrograms of fatty acid per milligram of tissue.

c16_3w4

Concentration of 16:3 ω 4 fatty acid as micrograms of fatty acid per milligram of tissue.

c16_3w3

Concentration of 16:3 ω 3 fatty acid as micrograms of fatty acid per milligram of tissue.

c16_4w3

Concentration of 16:4 ω 3 fatty acid as micrograms of fatty acid per milligram of tissue.

c16_4w1

Concentration of 16:4 ω 1 fatty acid as micrograms of fatty acid per milligram of tissue.

c18_0

Concentration of 18:0 fatty acid as micrograms of fatty acid per milligram of tissue.

c18_1w9

Concentration of 18:1 ω 9 fatty acid as micrograms of fatty acid per milligram of tissue.

c18_1w7

Concentration of 18:1 ω 7 fatty acid as micrograms of fatty acid per milligram of tissue.

c18_2w6t

Concentration of 18:2 ω 6t fatty acid as micrograms of fatty acid per milligram of tissue.

c18_2w6

Concentration of 18:2 ω 6 fatty acid as micrograms of fatty acid per milligram of tissue.

c18_3w6

Concentration of 18:3 ω 6 fatty acid as micrograms of fatty acid per milligram of tissue.

c18_3w3

Concentration of 18:3 ω 3 fatty acid as micrograms of fatty acid per milligram of tissue.

c18_4w4

Concentration of 18:4 ω 4 fatty acid as micrograms of fatty acid per milligram of tissue.

c18_4w3

Concentration of 18:4 ω 3 fatty acid as micrograms of fatty acid per milligram of tissue.

c18_5w3

Concentration of 18:5 ω 3 fatty acid as micrograms of fatty acid per milligram of tissue.

c20_0

Concentration of 20:0 fatty acid as micrograms of fatty acid per milligram of tissue.

c20_1w9

Concentration of 20:1 ω 9 fatty acid as micrograms of fatty acid per milligram of tissue.

c20_1w7

Concentration of 20:1 ω 7 fatty acid as micrograms of fatty acid per milligram of tissue.

c20_2w5_11

Concentration of 20:2-5-11 fatty acid as micrograms of fatty acid per milligram of tissue.

c20_2w5_13

Concentration of 20:2-5-13 fatty acid as micrograms of fatty acid per milligram of tissue.

c20_2w6

Concentration of 20:2 ω 6 fatty acid as micrograms of fatty acid per milligram of tissue.

c20_3w6

Concentration of 20:3 ω 6 fatty acid as micrograms of fatty acid per milligram of tissue.

c20_4w6

Concentration of 20:4 ω 6 fatty acid as micrograms of fatty acid per milligram of tissue.

c20_3w3

Concentration of 20:3 ω 3 fatty acid as micrograms of fatty acid per milligram of tissue.

c20_4w3

Concentration of 20:4 ω 3 fatty acid as micrograms of fatty acid per milligram of tissue.

c20_5w3

Concentration of 20:5 ω 3 fatty acid as micrograms of fatty acid per milligram of tissue.

c22_0

Concentration of 22:0 fatty acid as micrograms of fatty acid per milligram of tissue.

c22_1w9

Concentration of 22:1 ω 9 fatty acid as micrograms of fatty acid per milligram of tissue.

c22_1w7

Concentration of 22:1 ω 7 fatty acid as micrograms of fatty acid per milligram of tissue.

c22_2w6

Concentration of 22:2 ω 6 fatty acid as micrograms of fatty acid per milligram of tissue.

c22_4w6

Concentration of 22:4 ω 6 fatty acid as micrograms of fatty acid per milligram of tissue.

c22_5w6

Concentration of 22:5 ω 6 fatty acid as micrograms of fatty acid per milligram of tissue.

c22_3w3

Concentration of 22:3 ω 3 fatty acid as micrograms of fatty acid per milligram of tissue.

c22_4w3

Concentration of 22:4 ω 3 fatty acid as micrograms of fatty acid per milligram of tissue.

c22_5w3

Concentration of 22:5 ω 3 fatty acid as micrograms of fatty acid per milligram of tissue.

c22_6w3

Concentration of 22:6 ω 3 fatty acid as micrograms of fatty acid per milligram of tissue.

c24_0

Concentration of 24:0 fatty acid as micrograms of fatty acid per milligram of tissue.

comments

Quality flag column. Two samples spilled during fatty acid extraction. These samples are flagged as such. Although concentrations are lower than other samples, proportions between fatty acids are consistent.

invertebrates.csv

This file contains abundance for benthic macroinvertebrates collected at each of the 14 littoral sampling locations. Only amphipod taxa were identified to species. Mollusks and isopods were identified to genus.

site

Unique alphanumeric identifier for a sampling location.

replicate

Replicate for sampling location. While three replicates were collected in the field, some samples were poorly preserved, and invertebrates were not enumerated so as to prevent potential errors.

Acroloxidae

Mollusk genus

Asellidae

Endemic isopod genus

Baicaliidae

Mollusk genus, most of which are endemic

Benedictidae

Mollusk genus, most of which are endemic

Brandtia_latissima

Endemic amphipod species

Brandtia_parasitica_parasitica

Endemic amphipod species

Caddisflies

General grouping; were not identified to species.

Cryptoropus_inflatus

Endemic amphipod species

Cryptoropus_pachytus

Endemic amphipod species

Cryptoropus_rugosus

Endemic amphipod species

Eulimnogammarus_capreolus

Endemic amphipod species

Eulimnogammarus_cruentes
Endemic amphipod species

Eulimnogammarus_cyaneus
Endemic amphipod species

Eulimnogammarus_grandimanus
Endemic amphipod species

Eulimnogammarus_juveniles
Endemic amphipod genus. Identification kept at genus level so as to prevent misclassification.

Eulimnogammarus_maackii
Endemic amphipod species

Eulimnogammarus_marituji
Endemic amphipod species

Eulimnogammarus_verucossus
Endemic amphipod species

Eulimnogammarus_viridis_viridis
Endemic amphipod species

Eulimnogammarus_vittatus
Endemic amphipod species

Flatworms
Not identified beyond order.

Leeches
Not identified beyond order, although 12 endemic species do exist.

Maackia
Mollusk genus, most of which are endemic

Pallasea_brandtia_brandtia
Endemic amphipod species

Pallasea_brandtii_tenera
Endemic amphipod species

Pallasea_cancelloides
Endemic amphipod species

Pallasea_cancellus

Endemic amphipod species

Pallasea_viridis

Endemic amphipod species

Planorbidae

Mollusk genus, most of which are endemic

Poekilogammarus_crassimus

Endemic amphipod species

Poekilogammarus_ephippiatus

Endemic amphipod species

Poekilogammarus_juveniles

Endemic amphipod genus. Identifying to species introduced risk of misclassification.

Poekilogammarus_megonychus_perpolitus

Endemic amphipod species

Poekilogammarus_pictus

Endemic amphipod species

Valvatidae

Mollusk genus, most of which are endemic

metadata.csv

This file contains metadata for each of the pelagic and littoral sampling locations. Missing data are assigned as NA.

year

Year sampling occurred.

month

Month sampling occurred.

day

Day sampling occurred.

time

Time sampling occurred as Hours:Minutes.

site

Unique alphanumeric identifier for a sampling location.

lat

Latitude of sampling location in decimal degrees.

long

Longitude of sampling location in decimal degrees.

site_description

Researchers' description of sampling location at the time of sampling.

distance_to_shore_m

Distance from *in situ* sampled location to the shoreline in meters.

depth_m

Depth at *in situ* sampling location in meters.

air_temp_celsius

Temperature of air at sampling location in Celsius.

surface_temp_celsius

Temperature of water's surface at sampling location in Celsius.

mid_temp_celsius

Temperature of water midway between surface and bottom at sampling location in Celsius.

bottom_temp_celsius

Temperature of water near sediment at sampling location in Celsius.

comments

Notes in the field describing sampling conditions.

shore_photo

Whether or not photos of the shoreline were taken. Photos are available on the project's Open Science Framework page (Meyer et al. 2015).

substrate_photo

Whether or not photos of the substrate were taken.

sponges

Whether or not sponges were present at a sampling location.

brandtia

Whether or not *Brandtia spp.* (endemic amphipod) was present at a sampling location.

microplastics.csv

This file contains microplastics counts for each of the pelagic and littoral sampling locations.

site

Unique alphanumeric identifier for a sampling location.

replicate

Replicate for a given sampling location. Replicate values of “C” indicate a control.

fragments

Number of microplastic fragments observed.

fibers

Number of microplastic fibers observed.

beads

Number of microplastic beads observed.

comments

Observer comments while enumerating microplastics

volume_filtered_ml

Volume in milliliters for a given replicate filtered.

nutrients.csv

This file contains nutrient concentrations for each of the associated sampling locations. Nutrient samples were not filtered prior to analysis, meaning that nitrogen concentrations have the potential to include intracellular nitrogen. Therefore, nitrogenous species’ concentrations may be spurious.

site

Unique alphanumeric identifier for a sampling location.

replicate

Replicate for a given sampling location.

nh4_mg_dm3

Ammonium concentration in milligrams of ammonium per cubic decimeter.

no3_mg_dm3

Nitrate concentration in milligrams of nitrate per cubic decimeter

tp_mg_dm3

Total phosphorus concentration in milligrams of phosphorus per cubic decimeter.

tpo43_mg_dm3

Total phosphate concentration as phosphate in milligrams per cubic decimeter.

periphyton.csv

This file contains periphyton abundance data for each of the sampled littoral locations. For poorly preserved samples, counts are listed as NA for each taxonomic grouping, and a note in the “comments” column is provided.

site

Unique alphanumeric identifier for a sampling location.

replicate

Replicate number for a given sampling site.

subsamples_counted

Number of 10 microliter subsamples counted for a given replicate.

diatom

Number of diatom cells counted for a given replicate.

spirogyra

Number of *Spirogyra* spp. cells counted for a given replicate.

spirogyra_filament

Number of *Spirogyra* spp. filaments counted for a given replicate.

ulothrix

Number of *Ulothrix* spp. cells counted for a given replicate.

ulothrix_filament

Number of *Ulothrix* spp. filaments counted for a given replicate.

tetrasporales

Number of *Tetrasporales* spp. cells counted for a given replicate

pediastrum

Number of *Pediastrum* spp. cells counted for a given replicate.

desmidales

Number of *Desmidales* spp. cells counted for a given replicate.

comments

Notes from the observer.

ppcp.csv

This file contains Pharmaceutical and Personal Care Product (PPCP) concentrations for each littoral and pelagic sampling location. Detection limits are estimated to be 0.001 µg/L based on a 500 mL sample volume.

site

Unique alphanumeric identifier for a sampling location.

paraxanthine

Concentration of paraxanthine, also known as 1,7-dimethylxanthine, in micrograms per liter. Paraxanthine is the main human metabolite of caffeine

acetaminophen

Concentration of acetaminophen, also known as paracetamol, in micrograms per liter.

amphetamine

Concentration of amphetamine in micrograms per liter.

caffeine

Concentration of caffeine in micrograms per liter.

carbamazepine

Concentration of carbamazepine in micrograms per liter.

cimetidine

Concentration of cimetidine in micrograms per liter.

cotinine

Concentration of cotinine, which is the main human metabolite of nicotine, in micrograms per liter.

diphenhydramine

Concentration of diphenhydramine in micrograms per liter.

mda

Concentration of methylenedioxyamphetamine in micrograms per liter.

mdma

Concentration of methylenedioxymethamphetamine in micrograms per liter.

methamphetamine

Concentration of methamphetamine in micrograms per liter.

morphine

Concentration of morphine in micrograms per liter.

phenazone

Concentration of phenazone in micrograms per liter.

sulfachloropyridazine

Concentration of sulfachloropyridazine in micrograms per liter.

sulfamethazine

Concentration of *sulfamethazine* in micrograms per liter.

sulfamethoxazole

Concentration of sulfamethoxazole in micrograms per liter.

thiabendazole

Concentration of thiabendazole in micrograms per liter.

trimethoprim

Concentration of trimethoprim in micrograms per liter.

collection_year

Year sample was collected in the field.

collection_month

Month sample was collected in the field.

collection_day

Day sample was collected in the field.

analysis_year

Year sample was analyzed.

analysis_month

Month sample was analyzed.

analysis_day

Day sample was analyzed.

stable_isotopes.csv

This file contains carbon ($\delta^{13}\text{C}$) and nitrogen ($\delta^{15}\text{N}$) values for various benthic macroinvertebrate genera and periphyton collected from the 14 littoral sampling locations.

C13

Carbon ($\delta^{13}\text{C}$) stable isotope values in parts per thousand.

N15

Nitrogen ($\delta^{15}\text{N}$) stable isotope values in parts per thousand.

site

Unique alphanumeric identifier for a sampling location.

Genus

Genus of the analyzed organism.

Species

Species of the analyzed organism. When organism was identified solely to genus, the Species value is NA.

comments

Quality flag column where $\delta^{13}\text{C}$ samples were outside of the range of standards.

total_lipid.csv

site

Unique alphanumeric identifier for a sampling location.

Genus

Genus of the analyzed organism.

Species

Species of the analyzed organism. When organism was identified solely to genus, the Species value is NA.

total_lipid_mg_per_g

Total amount of lipids in a sample in milligrams of Lipid per gram of tissue.

deviation

Samples were weighed three times and standard deviation in measurement was calculated. All values are reported in milligrams of Lipid per gram of tissue.

comments

Quality flag column. Two samples spilled during fatty acid extraction. These samples are flagged as such.

Methods

Inverse distance weighted (IDW) population calculation for each sampling location

We recognized that sewage indicator concentrations at each sampling location may be related to a sampling location's spatial position relative to both the size and proximity of neighboring developed sites. Therefore, we created the inverse distance weighted (IDW) population metric to compress, into a single metric, information about human population size, density, and location along the shoreline as well as distance between developed sites and sampling locations.

Our workflow for calculating IDW population required five steps. First, we traced polygons of each lakeside development's perimeter and line geometries of each development's shorelines from satellite imagery for each developed site in Google Earth. Second, polygon and line geometries were downloaded from Google Earth as a .kml file. Third, the .kml file was imported into the R statistical environment (R Core Team 2019), where using the sf package (Pebesma 2018) we calculated shoreline length, polygon area, and centroid location for each developed site. Fourth, we joined point locations of each sampling site with the spatial polygons to calculate the distance from each sampling location to each developed site's centroid. Fifth, we calculated IDW population for each sampling location, using formula (1).

$$(1) I_j = \frac{\frac{P_{LI} * L_{LI}}{A_{LI}}}{D_{j,LI}} + \frac{\frac{P_{BK} * L_{BK}}{A_{BK}}}{D_{j,BK}} + \frac{\frac{P_{BGO} * L_{BGO}}{A_{BGO}}}{D_{j,BGO}}$$

where I is the IDW population at sampling location j , P is the population at each of the three developed sites Listvyanka (LI), Bolshie Koty (BK), Bolshoe Goloustnoe (BGO), A is the area of a developed site in km^2 , L is the shoreline length at a developed site in km, and D is the distance from developed site j to each developed site's centroid in km.

Nutrients

Water samples for nutrient analyses were collected in 150 mL glass jars that had been washed with phosphate-free soap and rinsed three times with water from the sampling location. Samples were collected in duplicates and immediately frozen at -20°C until processing at the A.P. Vinogradov Institute of Geochemistry (Siberian Branch of the Russian Academy of Sciences, Irkutsk). Samples were not filtered prior to freezing, meaning that nitrogen and ammonium concentrations may potentially include intracellular nitrogen and overestimate dissolved nitrogenous forms in the water column.

For each water sample, nitrate, ammonium, and total phosphorus concentrations were measured. For ammonium (2016a) and nitrate (2017) concentrations, samples were analyzed with a spectrophotometer following the addition of Nessler's reagent and disulfuric acid respectively. Total phosphorus concentration was measured with a spectrophotometer following the addition of persulfate (2016b).

Chlorophyll a

Water samples were collected in 1.5 L plastic bottles from a depth of approximately 0.75 m. Within 12 h of collection, three subsamples (up to 150 mL each) were filtered through 25-mm diameter, $0.2 \mu\text{m}$ pore size nitrocellulose filters. Filters were then placed in a 35 mm petri dish and frozen in the dark until processing.

Chlorophyll samples were processed in a manner similar to that of Parsons and Strickland (1963) and Lorenzen (1967). Nitrocellulose filters were ground in 90% acetone, in which chlorophyll extraction was allowed to proceed overnight. Samples were then centrifuged for 15-20 minutes. After centrifugation, absorbance of the chlorophyll extract was measured in a spectrophotometer at 630, 645, 665, and 750 nm. Concentrations were calculated using the formula: $C = 11.64(A_{665} - A_{750}) - 2.16(A_{645} - A_{750}) - 0.1(A_{630} - A_{750}) / (V_2/V_1)$; where A is the absorbance value of a

particular wavelength, V_1 is the volume of the filtered water, and V_2 is the volume of extract. Concentrations are reported as mg/L.

Pharmaceuticals and Personal Care Products (PPCPs)

Water samples for PPCP analysis were collected in 250 mL amber glass bottles that were rinsed with either methanol or acetone and then three times with sample water prior to collections. Following collection, samples were refrigerated and kept in the dark until solid phase extraction (SPE).

Within 12 h of collection, samples were filtered directly from the amber glass bottle using an in-line Teflon filter holder with glass microfiber GMF (1.0 μm pore size, WhatmanGrad 934-AH) in tandem with a solid phase extraction (SPE) cartridge (200 mg HLB, Waters Corporation, Milford, MA) connected to a 1-liter vacuum flask. Lab personnel wore gloves and face masks to minimize contamination. Prior to filtration, SPE cartridges were primed with at least 5 mL of either methanol or acetone and then washed with at least 5 mL of sample water. Rate of extraction was maintained at approximately 1 drop per second. Extraction proceeded until water could no longer pass through the SPE cartridge or until all collected water was filtered. Cartridges were stored in Whirlpaks at -20°C until analysis for 18 PPCP residues using liquid chromatography tandem mass spectrometry (LC-MS-MS) following methods of Lee et al. (2016) and D'Alessio et al (2018). Concentrations are reported in $\mu\text{g/L}$.

Microplastics

At each location, samples were collected in triplicate using 1.5 L clear plastic bottles that were washed thoroughly with sample water before each collection. Samples were collected by hand for each littoral site and with a metal bucket from aboard the ship for pelagic sites.

For processing, each sample was vacuum filtered on to a 47-mm diameter GF/F filter. During filtration, aluminum foil was used to cover the filtration funnel to prevent contamination from airborne microplastic particles. After filtration, filters were dried under vacuum pressure and then stored in 50-mm petri dishes. Following filtration of all three replicates, the filtrate was collected and then re-filtered through a GF/F filter as a control for contamination from the plastic vacuum funnel or potentially airborne microplastics.

Microplastic counting involved visual inspection of the entire GF/F in a similar manner to methods described in Hanvey et al. (2017). Visual enumeration was conducted under a stereo microscope with $\sim 100\times$ magnification, and microplastics were classified into one of three categories: fibers, fragments, or beads. For all categories, plastics were defined as observed objects with apparent artificial colors, so as to not enumerate plastics potentially contributed from the sampling bottle itself. Fibers were defined as smooth, long plastics with consistent diameters. Fragments were defined as plastics with irregularly sharp or jagged edges. Beads were defined as spherical plastics. Although we did not measure microplastic size, this technique likely allowed us to reliably quantify microplastics as small as $\sim 300\ \mu\text{m}$ (Hanvey et al. 2017). During enumeration, GF/Fs remained covered in the petri dish to minimize potential for contamination from the air.

It is worth noting that since the time of our field sampling, evidence has accumulated that our methods likely dramatically underestimated microplastic abundance (Wang and Wang 2018; Brandon et al. 2020). Recent investigations of microplastics in Lake Baikal near Bolshie Koty (BK) used analogous methods and measured similar microplastic concentrations (Karnaukhov et al. 2020). Future studies aiming to use these data for comparison or supplementing potential data gaps should consider the minimum microplastic size that could be reliably detected by the method, so as to ensure data are comparable across methods.

Periphyton abundance

At each littoral site, we haphazardly selected three rocks representative of local substrate. A plastic stencil was used to define a surface area of each rock from which we scraped a standardized 14.5 cm² patch of periphyton. Samples were preserved with Lugol's solution and stored in plastic scintillation vials. Additional periphyton was collected in composite from each site for fatty acid and stable isotope analysis.

Periphyton taxonomic identification and enumeration was performed by subsampling 10 µL aliquots from each preserved sample. Cells, filaments, and colonies were counted for each taxonomic group until at least 300 cells were identified. For all 10 µL aliquots, the entire subsample was counted, even if 300 cells were counted prior to completing the aliquot. Taxa were classified into broad categories consistent with Baikal algal taxonomy (Izhboldina 2007), using coarse groupings to capture general patterns in relative algal abundance. As a result, algal groups consisted of diatoms, *Ulothrix*, *Spirogyra*, and the green algal Order Tetrasporales.

Benthic macroinvertebrate abundance

Three kick-net samples were collected for assessment of benthic community composition and abundance. Using a D-net, we collected macroinvertebrates by flipping over 1-3 rocks, and then sweeping five times in a left-to-right motion across approximately 1 m. After the series of sweeps, the catch was rinsed into a plastic bucket. For each replicate, bucket contents were concentrated using a 64-µm mesh and placed in glass jars with 40% ethanol (vodka; the only preservative available to us at the time) for preservation and refrigerated at 4°C aboard the research vessel. The 40% ethanol preservative was replaced with ~80% ethanol upon return to the lab within 24 to 48 hours, and samples were stored at ~4°C.

Invertebrate taxonomic identification and enumeration were performed under a stereo microscope. All invertebrates were identified to species with the exception of juveniles (Takhteev and Didorenko (2015) for amphipods; Sitnikova (2012) for mollusks; Table 2). All samples contained oligochaetes and polychaetes, but due to poor preservation, these taxa were not counted. Six samples of the 42 collected were not well-preserved and were excluded from further analyses, in order to reduce errors in identification. KD-1 and LI-1 were the only sites with 1 sample counted. BK-2 and KD-2 each had two samples counted.

Stable Isotope Analysis

Measurements of $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ were performed on an elemental analyzer-isotope ratio mass spectrometer (EA-IRMS; Finnigan DELTApplus XP, Thermo Scientific) at the Large Lakes Observatory, University of Minnesota Duluth.

Fatty Acid Analysis

Following freeze-drying, samples were transferred to 10 mL glass centrifuge vials, and 2 mL of 100% chloroform was added to each under nitrogen gas. Samples were allowed to sit in chloroform overnight at -80°C . Fatty acid extractions generally involved three phases: (1) 100% chloroform extraction, (2) chloroform-methanol extraction, and (3) fatty acid methylation. Fatty acid extraction methods were adapted from Schram et al. (2018).

After overnight chloroform extraction, samples underwent a chloroform-methanol extraction three times. To each sample, we added 1 mL cooled 100% methanol, 1 mL chloroform:methanol solution (2:1), and 0.8 mL 0.9% NaCl solution. Samples were inverted three times and sonicated on ice for 10 minutes. Next, samples were vortexed for 1 minute, and centrifuged for 5 minutes (3,000 rpm) at 4°C . Using a double pipette technique, the lower organic layer was removed and kept under nitrogen. After the third extraction, samples were evaporated under nitrogen flow, and resuspended in 1.5 mL chloroform and stored at -20°C overnight.

Once resuspended in chloroform, 1 mL of chloroform extract was transferred to a glass centrifuge tube with a glass syringe as well as an internal standard of 4 μL of 19-carbon fatty acid. Samples were then evaporated under nitrogen, and then 1 mL of toluene and 2 mL of 1% sulfuric acid-methanol was added. The vial was closed under nitrogen gas and then incubated in 50°C water bath for 16 hours. After incubation, samples were removed from the bath, allowed to reach room temperature and stored on ice. Next, we performed a potassium carbonate-hexane extraction twice. To each sample, we added 2 mL of 2% potassium bicarbonate and 5 mL of 100% hexane, inverting the capped vial so as to mix the solution. Samples were centrifuged for 3 minutes (1,500 rpm) at 4°C . The upper hexane layer was then removed and placed in a vial to evaporate under nitrogen flow. Once almost evaporated, 1 mL of 100% hexane was added and stored in a glass amber autosampler vial for GC/MS quantification. GC/MS quantification was performed with a Shimadzu QP2020 GC/MS following Schram et al. (2018).

Technical Validation

The dataset had three main validation procedures: taxonomic, analytical, and reproducible.

For taxonomic validation, all phylogenetic groupings were based off most recent identification keys. Amphipods were identified according to Takhteev & Didorenko (2015). Mollusks were identified according to Sitnikova (2012). Algal taxa were identified according to Izhboldina (2007). For consistency, all taxa were identified by one person (Michael F. Meyer), who was trained by experts in Baikal algal and macroinvertebrate taxonomy.

For analytical validation, internal standards were used for all mass-spectroscopy analyses. PPCP analyses involved labeled internal standards ($^{13}\text{C}_3$ -caffeine, methamphetamine-d8, MDMA-d8, morphine-d3, and $^{13}\text{C}_6$ -sulfamethazine). Stable isotope values were calibrated against certified

reference materials including L-glutamic acid (NIST SRM 8574), low organic soil and sorghum flour (standards B-2153 and B-2159 from Elemental Micro-analysis Ltd., Okehampton, UK) and in-house standards (acetanilide and caffeine). Replicate analyses of external standards showed a mean standard deviation of 0.06 ‰ and 0.09 ‰, for $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$, respectively. Finally, fatty acid estimations used an internal 19:0 standard to assess oxidation of fatty acids during extraction, methylation, and quantification.

For data reproducibility, data aggregation and harmonization procedures were conducted in the R statistical environment (R Core Team 2019), using the tidyverse (Wickham et al. 2019) packages. We also implemented a protocol for systematically reviewing all R scripts. All data, including .kml files used to calculate IDW metric, are publicly available from the Environmental Data Initiative repository (Meyer et al. 2020b), and all R scripts are available from the GitHub repository of this project's Open Science Framework account (Meyer et al. 2015). We systematically reviewed all R code to format and aggregate source data. An initial cleaning script (00_disaggregated_data_cleaning.R) removed incorrect spellings, erroneous data values, and inconsistent column names from raw data. This step created the standardized .csv files detailed above, which are available on the EDI repository. Raw data files are available on the project's Open Science Framework page (DOI) but are not included in the EDI repository to prevent confusion or incorrect usage. Data hosted on EDI are at the replicate-level but can be aggregated to the sampling-site-level using script 01_data_cleaning.R. In addition to aggregation scripts, six R scripts used for analyses in Meyer et al. (*Under Review*) are also available on the EDI repository within the compressed entity "scripts.tar.gz". All R code for data aggregation was written by one person (Michael F. Meyer) and then independently reviewed by two others (Matthew R. Brouil and Kara H. Woo) to confirm that code performed as intended, was well documented, and annotations were complete.

A commitment to FAIR and TRUST principles

Throughout the dataset's development, we strove to incorporate both FAIR (Findable, Accessible, Interoperable, and Reproducible) and TRUST (Transparency, Responsibility, User Focus, Sustainability, and Technology) principles where applicable.

With respect to FAIR principles (Wilkinson et al. 2016), the data are openly accessible in a standardized, replicate-level format on the EDI portal. The 11 CSV files contained within the dataset are entirely interoperable using the "site" column, enabling all variables to efficiently be merged together. Finally, all analytical and some data wrangling scripts are available on the EDI portal in a compressed format, such that future users can reproduce data manipulation and analyses described in Meyer et al. (*Under Review*).

With respect to TRUST principles (Lin et al. 2020), we strove to document additional metadata and data-cleaning practices in a public Open Science Framework (OSF) repository (Meyer et al. 2015). These steps are not necessarily critical to the core EDI dataset, but provide increased transparency for future users wishing to recreate the dataset de novo. All "raw" data are provided in the OSF portal, including an initial cleaning script (00_disaggregated_data_cleaning.R) to remove incorrect spellings, erroneous data values, and inconsistent column names. This repository also includes photographs of both field notes as well as photographs of shoreline and substrate from sampling

locations. To empower and expedite future reuse, all directories are accompanied with documentation that details directory contents, and all associated scripts are documented and annotated. While many of the files are redundant from the EDI repository, the OSF repository is meant to supplement the EDI repository, so as to enable sustainable, user-focused transparency of how data were collected and cleaned from their raw formats.

Data Use and Recommendations for Reuse

Recognizing the potential for continued low-level, sewage pollution at Lake Baikal (Timoshkin et al. 2016, 2018; Volkova et al. 2018) and lakes worldwide (Yang et al. 2018; Meyer et al. 2019), the final dataset can be applied to a suite of research questions pertaining to ecological responses to human disturbance. We highlight two main areas for immediate application.

First, the final data products can be harmonized with other littoral sampling efforts throughout Lake Baikal, so as to enhance spatial coverage and data diversity. Since 2010, Lake Baikal has experienced increasing filamentous algal abundance, especially near larger lakeside developments (Kravtsova et al. 2014; Timoshkin et al. 2016, 2018; Volkova et al. 2018). Recent benthic algal surveys throughout Lake Baikal's entirety have suggested that cosmopolitan filamentous algae, such as *Spirogyra spp.*, tend to be more abundant near larger lakeside developments (Timoshkin et al. 2016; Volkova et al. 2018). For example, Listvyanka is a small town located at the beginning of the Angara River, Lake Baikal's only surface outflow. While Listvyanka's permanent population is approximately 2,000 persons, the town is a growing tourism hub, and hosts over 1.2 million tourists per year (Interfax-Tourism 2018). Surveys conducted near Listvyanka have suggested increased *Spirogyra spp.* abundance is associated with wastewater release (Timoshkin et al. 2016). Although wastewater inputs are likely low and are diluted to negligible concentrations offshore (Meyer et al., 202X), combining monitoring efforts across spatial and temporal scales are necessary to evaluate the spatial and temporal extent of wastewater entering Lake Baikal. As such, our data could complement previous, current, and future monitoring efforts, where observations may be missing.

Second, the final data products are useful to expanding freshwater PPCP, microplastic, and associated biological responses across large spatial scales. Recent syntheses of the PPCP literature have reported that studies involving lakes are less abundant relative to those focused on lotic systems (Meyer et al. 2019). Likewise, microplastic studies have noted that freshwater environments are less represented in the literature relative to marine ecosystems (Horton et al. 2017). For both PPCPs and microplastics, toxic responses to even minute concentrations can be uncertain and differ between ecosystem types (e.g., Rosi-Marshall et al. 2013 for lotic and Shaw et al. 2015 for lentic). As a result of PPCPs and microplastics garnering increasing attention worldwide, sampling of PPCPs and microplastics with co-located biological data across multiple spatial and temporal scales would be necessary to synthesize biotic responses to micropollutants across systems. Although our data constitute a limited sample number of PPCP and microplastic data that exist globally, our final data products are highly structured and flexible for merging with similar datasets. Additionally, our dataset's sequential harmonization workflow could be adopted by similar monitoring efforts, thereby facilitating data interoperability. Through integration with similar monitoring efforts, our dataset can contribute to global synthesis of emerging contaminant consequences, especially in a region of the world that is often not easily accessible to many researchers.

References

- Barnes, D. K. A., F. Galgani, R. C. Thompson, and M. Barlaz. 2009. Accumulation and fragmentation of plastic debris in global environments. *Philos Trans R Soc Lond B Biol Sci* **364**: 1985–1998. doi:10.1098/rstb.2008.0205
- Bendz, D., N. A. Paxéus, T. R. Ginn, and F. J. Loge. 2005. Occurrence and fate of pharmaceutically active compounds in the environment, a case study: Höje River in Sweden. *Journal of Hazardous Materials* **122**: 195–204. doi:10.1016/j.jhazmat.2005.03.012
- Brandon, J. A., A. Freibott, and L. M. Sala. 2020. Patterns of suspended and salp-ingested microplastic debris in the North Pacific investigated with epifluorescence microscopy. *Limnology and Oceanography Letters* **5**: 46–53. doi:10.1002/lol2.10127
- Brodin, T., J. Fick, M. Jonsson, and J. Klaminder. 2013. Dilute Concentrations of a Psychiatric Drug Alter Behavior of Fish from Natural Populations. *Science* **339**: 814–815. doi:10.1126/science.1226850
- Camilleri, A. C., and T. Ozersky. 2019. Large variation in periphyton $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values in the upper Great Lakes: Correlates and implications. *Journal of Great Lakes Research* **45**: 986–990. doi:10.1016/j.jglr.2019.06.003
- Costanzo, S. D., M. J. O'Donohue, W. C. Dennison, N. R. Loneragan, and M. Thomas. 2001. A New Approach for Detecting and Mapping Sewage Impacts. *Marine Pollution Bulletin* **42**: 149–156. doi:10.1016/S0025-326X(00)00125-9
- D'Alessio, M., S. Onanong, D. D. Snow, and C. Ray. 2018. Occurrence and removal of pharmaceutical compounds and steroids at four wastewater treatment plants in Hawai'i and their environmental fate. *Science of The Total Environment* **631–632**: 1360–1370. doi:10.1016/j.scitotenv.2018.03.100

- Dalsgaard, J., M. St. John, G. Kattner, D. Müller-Navarra, and W. Hagen. 2003. Fatty acid trophic markers in the pelagic marine environment, p. 225–340. *In* *Advances in Marine Biology*. Elsevier.
- Edmondson, W. T. 1970. Phosphorus, Nitrogen, and Algae in Lake Washington after Diversion of Sewage. *Science* **169**: 690–691.
- Fellows, I., and using the Jm. library by J. P. Stotz. 2019. OpenStreetMap: Access to Open Street Map Raster Images,.
- Focazio, M. J., D. W. Kolpin, K. K. Barnes, E. T. Furlong, M. T. Meyer, S. D. Zaugg, L. B. Barber, and M. E. Thurman. 2008. A national reconnaissance for pharmaceuticals and other organic wastewater contaminants in the United States - II) Untreated drinking water sources. *SCIENCE OF THE TOTAL ENVIRONMENT* **402**: 201–216.
doi:10.1016/j.scitotenv.2008.02.021
- Gartner, A., P. Lavery, and A. J. Smit. 2002. Use of delta N-15 signatures of different functional forms of macroalgae and filter-feeders to reveal temporal and spatial patterns in sewage dispersal. *Mar. Ecol.-Prog. Ser.* **235**: 63–73. doi:10.3354/meps235063
- Green, D. S. 2016. Effects of microplastics on European flat oysters, *Ostrea edulis* and their associated benthic communities. *Environmental Pollution* **216**: 95–103.
doi:10.1016/j.envpol.2016.05.043
- Hall, R. I., P. R. Leavitt, R. Quinlan, A. S. Dixit, and J. P. Smol. 1999. Effects of agriculture, urbanization, and climate on water quality in the northern Great Plains. *Limnology and Oceanography* **44**: 739–756. doi:10.4319/lo.1999.44.3_part_2.0739

- Hampton, S. E., S. C. Fradkin, P. R. Leavitt, and E. E. Rosenberger. 2011. Disproportionate importance of nearshore habitat for the food web of a deep oligotrophic lake. *Marine and Freshwater Research* **62**: 350. doi:10.1071/MF10229
- Hampton, S. E., L. R. Izmet'eva, M. V. Moore, S. L. Katz, B. Dennis, and E. A. Silow. 2008. Sixty years of environmental change in the world's largest freshwater lake - Lake Baikal, Siberia. *Global Change Biology* **14**: 1947–1958. doi:10.1111/j.1365-2486.2008.01616.x
- Hampton, S. E., S. McGowan, T. Ozersky, and others. 2018. Recent ecological change in ancient lakes. *Limnology and Oceanography* **63**: 2277–2304. doi:10.1002/lno.10938
- Hanvey, J. S., P. J. Lewis, J. L. Lavers, N. D. Crosbie, K. Pozo, and B. O. Clarke. 2017. A review of analytical techniques for quantifying microplastics in sediments. *Anal. Methods* **9**: 1369–1383. doi:10.1039/C6AY02707E
- Horton, A. A., A. Walton, D. J. Spurgeon, E. Lahive, and C. Svendsen. 2017. Microplastics in freshwater and terrestrial environments: Evaluating the current understanding to identify the knowledge gaps and future research priorities. *Science of The Total Environment* **586**: 127–141. doi:10.1016/j.scitotenv.2017.01.190
- Interfax-Tourism. 2018. Байкал с января по август 2018 года посетили 1,2 миллиона туристов (1.2 million tourists visited Baikal from January through August 2018). Interfax-Tourism, October 25
- Izhboldina, L. A. 2007. Guide and Key to Benthic and Periphyton Algae of Lake Baikal (meio- and macrophytes) with Brief Notes on Their Ecology, Nauka-Centre.
- Izmet'eva, L. R., M. V. Moore, S. E. Hampton, and others. 2016. Lake-wide physical and biological trends associated with warming in Lake Baikal. *Journal of Great Lakes Research* **42**: 6–17. doi:10.1016/j.jglr.2015.11.006

- Jeppesen, E., M. Søndergaard, J. P. Jensen, and others. 2005. Lake responses to reduced nutrient loading – an analysis of contemporary long-term data from 35 case studies. *Freshwater Biology* **50**: 1747–1771. doi:10.1111/j.1365-2427.2005.01415.x
- Karnaukhov, D., S. Biritskaya, E. Dolinskaya, M. Teplykh, N. Silenko, Y. Ermolaeva, and E. Silow. 2020. POLLUTION BY MACRO- AND MICROPLASTIC OF LARGE LACUSTRINE ECOSYSTEMS IN EASTERN ASIA. *Pollution Research* **2**: 353–355.
- Kassambara, A. 2019. ggpubr: “ggplot2” Based Publication Ready Plots,.
- Katz, S. L., L. R. Izmet'seva, S. E. Hampton, T. Ozersky, K. Shchapov, M. V. Moore, S. V. Shimaraeva, and E. A. Silow. 2015. The “Melosira years” of Lake Baikal: Winter environmental conditions at ice onset predict under-ice algal blooms in spring. *Limnology and Oceanography* **60**: 1950–1964. doi:10.1002/lno.10143
- Kolpin, D. W., E. T. Furlong, M. T. Meyer, E. M. Thurman, S. D. Zaugg, L. B. Barber, and H. T. Buxton. 2002. Pharmaceuticals, Hormones, and Other Organic Wastewater Contaminants in U.S. Streams, 1999–2000: A National Reconnaissance. *Environmental Science & Technology* **36**: 1202–1211. doi:10.1021/es011055j
- Kozhova, O. M., and L. R. Izmet'seva. 1998. Lake Baikal: Evolution and Biodiversity, Backhuys Publishers.
- Kravtsova, L. S., L. A. Izboldina, I. V. Khanaev, and others. 2014. Nearshore benthic blooms of filamentous green algae in Lake Baikal. *Journal of Great Lakes Research* **40**: 441–448. doi:10.1016/j.jglr.2014.02.019
- Lapointe, B. E., L. W. Herren, D. D. Debortoli, and M. A. Vogel. 2015. Evidence of sewage-driven eutrophication and harmful algal blooms in Florida's Indian River Lagoon. *Harmful Algae* **43**: 82–102. doi:10.1016/j.hal.2015.01.004

- Lee, S. S., A. M. Paspalof, D. D. Snow, E. K. Richmond, E. J. Rosi-Marshall, and J. J. Kelly. 2016. Occurrence and Potential Biological Effects of Amphetamine on Stream Communities. *Environmental Science & Technology* **50**: 9727–9735. doi:10.1021/acs.est.6b03717
- Lin, D., J. Crabtree, I. Dillo, and others. 2020. The TRUST Principles for digital repositories. *Scientific Data* **7**: 144. doi:10.1038/s41597-020-0486-7
- Lorenzen, C. J. 1967. Determination of Chlorophyll and Pheo-Pigments: Spectrophotometric Equations. *Limnology and Oceanography* **12**: 343–346. doi:https://doi.org/10.4319/lo.1967.12.2.0343
- Meyer, M. F., S. G. Labou, A. N. Cramer, M. R. Brousil, and B. T. Luff. 2020a. The global lake area, climate, and population dataset. *Scientific Data* **7**: 174. doi:10.1038/s41597-020-0517-4
- Meyer, M. F., T. Ozersky, K. H. Woo, and others. 2020b. A unified dataset of co-located sewage pollution, periphyton, and benthic macroinvertebrate community and food web structure from Lake Baikal (Siberia). doi:10.6073/PASTA/76F43144015EC795679BAC508EFA044B
- Meyer, M. F., T. Ozersky, K. H. Woo, and others. Effects of spatially heterogeneous lakeside development on nearshore biotic communities in a large, deep, oligotrophic lake (Lake Baikal, Siberia).
- Meyer, M. F., S. M. Powers, and S. E. Hampton. 2019. An Evidence Synthesis of Pharmaceuticals and Personal Care Products (PPCPs) in the Environment: Imbalances among Compounds, Sewage Treatment Techniques, and Ecosystem Types. *Environ. Sci. Technol.* **53**: 12961–12973. doi:10.1021/acs.est.9b02966
- Meyer, M., T. Ozersky, K. Woo, A. W. E. Galloway, M. R. Brousil, and S. Hampton. 2015. Baikal Food Webs. doi:10.17605/OSF.IO/9TA8Z

- Moore, J. W., D. E. Schindler, M. D. Scheuerell, D. Smith, and J. Frodge. 2003. Lake eutrophication at the urban fringe, Seattle region, USA. *AMBIO: A Journal of the Human Environment* **32**: 13–18.
- Moore, M. V., S. E. Hampton, L. R. Izmet'seva, E. A. Silow, E. V. Peshkova, and B. K. Pavlov. 2009. Climate Change and the World's "Sacred Sea"-Lake Baikal, Siberia. *Bioscience* **59**: 405–417. doi:10.1525/bio.2009.59.5.8
- O'Donnell, D. R., P. Wilburn, E. A. Silow, L. Y. Yampolsky, and E. Litchman. 2017. Nitrogen and phosphorus colimitation of phytoplankton in Lake Baikal: Insights from a spatial survey and nutrient enrichment experiments. *Limnology and Oceanography* **62**: 1383–1392. doi:10.1002/lno.10505
- Parsons, T. R., and J. D. H. Strickland. 1963. Discussion of spectrophotometric determination of marine-plant pigments, with revised equations for ascertaining chlorophylls and carotenoids. *Journal of Marine Research*.
- Pebesma, E. 2018. Simple Features for R: Standardized Support for Spatial Vector Data. *The R Journal* **10**: 439–446. doi:10.32614/RJ-2018-009
- Powers, S. M., T. W. Bruulsema, T. P. Burt, and others. 2016. Long-term accumulation and transport of anthropogenic phosphorus in three river basins. *Nature Geoscience* **9**: 353–356. doi:10.1038/ngeo2693
- R Core Team. 2019. R: A Language and Environment for Statistical Computing,.
- Richmond, E. K., M. R. Grace, J. J. Kelly, A. J. Reisinger, E. J. Rosi, and D. M. Walters. 2017. Pharmaceuticals and personal care products (PPCPs) are ecological disrupting compounds (EcoDC). *Elem Sci Anth* **5**: 66. doi:10.1525/elementa.252

- Richmond, E. K., E. J. Rosi, D. M. Walters, J. Fick, S. K. Hamilton, T. Brodin, A. Sundelin, and M. R. Grace. 2018. A diverse suite of pharmaceuticals contaminates stream and riparian food webs. *Nature Communications* **9**: 4491. doi:10.1038/s41467-018-06822-w
- Romera-Castillo, C., M. Pinto, T. M. Langer, X. A. Álvarez-Salgado, and G. J. Herndl. 2018. Dissolved organic carbon leaching from plastics stimulates microbial activity in the ocean. *Nat Commun* **9**: 1–7. doi:10.1038/s41467-018-03798-5
- Rosenberger, E. E., S. E. Hampton, S. C. Fradkin, and B. P. Kennedy. 2008. Effects of shoreline development on the nearshore environment in large deep oligotrophic lakes. *Freshwater Biology* **53**: 1673–1691. doi:10.1111/j.1365-2427.2008.01990.x
- Rosi-Marshall, E. J., D. W. Kincaid, H. A. Bechtold, T. V. Royer, M. Rojas, and J. J. Kelly. 2013. Pharmaceuticals suppress algal growth and microbial respiration and alter bacterial communities in stream biofilms. *Ecological Applications* **23**: 583–593. doi:10.1890/12-0491.1
- Rosi-Marshall, E. J., and T. V. Royer. 2012. Pharmaceutical Compounds and Ecosystem Function: An Emerging Research Challenge for Aquatic Ecologists. *Ecosystems* **15**: 867–880. doi:10.1007/s10021-012-9553-z
- Sargent, J. R., and S. Falk-Petersen. 1988. The lipid biochemistry of calanoid copepods. *Hydrobiologia* **167–168**: 101–114. doi:10.1007/BF00026297
- Schram, J. B., J. N. Kobelt, M. N. Dethier, and A. W. E. Galloway. 2018. Trophic Transfer of Macroalgal Fatty Acids in Two Urchin Species: Digestion, Egestion, and Tissue Building. *Front. Ecol. Evol.* **6**. doi:10.3389/fevo.2018.00083
- Shaw, L., C. Phung, and M. Grace. 2015. Pharmaceuticals and personal care products alter growth and function in lentic biofilms. *Environmental Chemistry* **12**: 301. doi:10.1071/EN14141

- Sitnikova, T. Ya. 2012. Определитель брюхоногих моллюсков бухты Большие Коты (юго-западное побережье озера Байкал) [Key of the Gastropod Molluscs in the Bay of Bolshie Koty (South-West shoreline of Lake Baikal)], Irkutsk State University.
- Slowikowski, K. 2019. ggrepel: Automatically Position Non-Overlapping Text Labels with “ggplot2.”
- Swann, G. E. A., V. N. Panizzo, S. Piccolroaz, and others. 2020. Changing nutrient cycling in Lake Baikal, the world’s oldest lake. PNAS **117**: 27211–27217. doi:10.1073/pnas.2013181117
- Taipale, S., U. Strandberg, E. Peltomaa, A. W. E. Galloway, A. Ojala, and M. T. Brett. 2013. Fatty acid composition as biomarkers of freshwater microalgae: analysis of 37 strains of microalgae in 22 genera and in seven classes. Aquatic Microbial Ecology **71**: 165–178. doi:10.3354/ame01671
- Takhteev, V. V., and D. I. Didorenko. 2015. Fauna and ecology of amphipods of Lake Baikal: A Training manual, V.B. Sochava Institute of Geography SB RAS.
- Timoshkin, O. A., M. V. Moore, N. N. Kulikova, and others. 2018. Groundwater contamination by sewage causes benthic algal outbreaks in the littoral zone of Lake Baikal (East Siberia). Journal of Great Lakes Research. doi:10.1016/j.jglr.2018.01.008
- Timoshkin, O. A., D. P. Samsonov, M. Yamamuro, and others. 2016. Rapid ecological change in the coastal zone of Lake Baikal (East Siberia): Is the site of the world’s greatest freshwater biodiversity in danger? Journal of Great Lakes Research **42**: 487–497. doi:10.1016/j.jglr.2016.02.011
- Tong, Y., M. Wang, J. Peñuelas, and others. 2020. Improvement in municipal wastewater treatment alters lake nitrogen to phosphorus ratios in populated regions. Proc Natl Acad Sci USA **117**: 11566–11572. doi:10.1073/pnas.1920759117

- Turetsky, M. R., R. K. Wieder, C. J. Williams, and D. H. Vitt. 2000. Organic matter accumulation, peat chemistry, and permafrost melting in peatlands of boreal Alberta. *Écoscience* **7**: 115–122. doi:10.1080/11956860.2000.11682608
- Vendel, A. L., F. Bessa, V. E. N. Alves, A. L. A. Amorim, J. Patrício, and A. R. T. Palma. 2017. Widespread microplastic ingestion by fish assemblages in tropical estuaries subjected to anthropogenic pressures. *Marine Pollution Bulletin* **117**: 448–455. doi:10.1016/j.marpolbul.2017.01.081
- Volkova, E. A., N. A. Bondarenko, and O. A. Timoshkin. 2018. Morphotaxonomy, distribution and abundance of *Spirogyra* (Zygnematophyceae, Charophyta) in Lake Baikal, East Siberia. *Phycologia* **57**: 298–308. doi:10.2216/17-69.1
- Wang, W., and J. Wang. 2018. Investigation of microplastics in aquatic environments: An overview of the methods used, from field sampling to laboratory analysis. *TrAC Trends in Analytical Chemistry* **108**: 195–202. doi:10.1016/j.trac.2018.08.026
- Wickham, H. 2014. Tidy Data. *Journal of Statistical Software* **59**: 1–23. doi:10.18637/jss.v059.i10
- Wickham, H., M. Averick, J. Bryan, and others. 2019. Welcome to the tidyverse. *Journal of Open Source Software* **4**: 1686. doi:10.21105/joss.01686
- Wilke, C. O. 2019. cowplot: Streamlined Plot Theme and Plot Annotations for “ggplot2,.”
- Wilkinson, M. D., M. Dumontier, Ij. J. Aalbersberg, and others. 2016. The FAIR Guiding Principles for scientific data management and stewardship. *Sci Data* **3**. doi:10.1038/sdata.2016.18
- Yang, Y., W. Song, H. Lin, W. Wang, L. Du, and W. Xing. 2018. Antibiotics and antibiotic resistance genes in global lakes: A review and meta-analysis. *Environment International* **116**: 60–73. doi:10.1016/j.envint.2018.04.011

Yoshida, T., T. Sekino, M. Genkai-Kato, and others. 2003. Seasonal dynamics of primary production in the pelagic zone of southern Lake Baikal. *Limnology* **4**: 53–62.

doi:10.1007/s10201-002-0089-3

2016a. Methods for determination of nitrogen-containing matters (with corrections) (Методы определения азотсодержащих веществ (с Поправками)).

2016b. Methods for determination of phosphorus-containing matters (with corrections) (Методы определения фосфорсодержащих веществ).

2017. Nitrate concentration in waters: Photometric methods with Giress reagent following stabilization in a cadmium reducer (Массовая концентрация нитратного азота в водах: Методика измерений фотометрическим методом с реактивом Грисса после восстановления в кадмиевом редукторе).

Acknowledgements

We would like to thank the faculty, students, staff, and mariners of the Irkutsk State University's Biological Research Institute Biostation for their expert field, taxonomic, and laboratory support; Marianne Moore and Bart De Stasio for helpful advice; the researchers and students of the Siberian Branch of the Russian Academy of Sciences Limnological Institute for expert taxonomic and logistical assistance; Oleg A. Timoshkin, Tatiana Ya. Sitnikova, Irina V. Mekhanikova, Nina A. Bondorenko, Ekaterina Volkova, and Vadim V. Takhteev for offering insights and taxonomic training throughout the development of this project. Funding was provided by the National Science Foundation (NSF-DEB-1136637) to S.E.H., a Fulbright Fellowship to M.F.M., a NSF Graduate Research Fellowship to M.F.M. (NSF-DGE-1347973), and Russian Ministry of Science and Education (N FZZE-2020-0026). This work serves as one chapter of M.F.M.'s doctoral dissertation in Environmental and Natural Resource Sciences at Washington State University.




Figure 1: Map of all sampling locations with sites labeled with unique alphanumeric code. The entire transect included three developed sites (i.e., Listvyanka (LI), Bolshie Koty (BK), Bolshoe Goloustnoe (BGO)). Three offshore sites (OS) were also sampled to compare pelagic sewage signals to those in the littoral. Sites without adjacent lakeside development included Emelyanikha Bay (EM), Maloe Kadilnoe (KD), Mys Soboliny (MS), Sredny Mys (SM). Littoral sampling locations were all 8.9-20.75 m from shore and at a depth approximately of 0.75 m, whereas pelagic sites were approximately 2-5 km from shore and ranged in depth from 900 to 1300 m. This map was created using the R statistical environment (R Core Team 2019) and the tidyverse (Wickham et al. 2019), OpenStreetMap (Fellows and Stotz 2019), ggpubr (Kassambara 2019), cowplot (Wilke 2019), and ggrepel (Slowikowski 2019) packages.

Site	Latitude	Longitude	Depth (m)	Distance to shore (m)
BK-1	51.90316	105.074	0.7	10
BK-2	51.90365	105.069	0.9	17.5
BK-3	51.90536	105.0957	0.8	10
BGO-1	52.02693	105.401	0.9	18
BGO-2	52.0197	105.3771	1.1	14
BGO-3	52.02649	105.4358	0.7	21
OS-1	51.98559	105.4724	900	NA
KD-1	51.92646	105.245	0.8	20.75
KD-2	51.91807	105.2146	0.9	14.5
MS-1	51.89863	105.1502	0.6	10.5
SM-1	51.87152	104.9801	0.9	11.5
LI-1	51.86825	104.8304	0.6	8.9
LI-2	51.84626	104.8736	0.8	9.4
LI-3	51.85407	104.8622	0.7	9.25
EM-1	51.86005	104.94	0.7	15.5
OS-2	51.8553	104.8148	1300	NA
OS-3	51.85911	105.0769	1400	5000

Table 1: Locational information for each of the 17 sampling stations. “OS” refers to pelagic locations (i.e., “Offshore”), whereas other site abbreviations refer to littoral sampling locations.