
Core Mathematics

A Practical Guide on Mathematics for Physics Learners

Sho Iwamoto

v0.0.1 – July 21, 2026

Lecture notes for first-year university students in physics or physical science, covering mathematical tools you the students need most. Calculus, logic, complex numbers, and vectors, with introductions to vector calculus, differential equations, and statistical analysis. Exercises are central to these notes. Practice each topic repeatedly until you can work through problems quickly and accurately.

Preface

Physics is an activity to describe the world, but how? At present, mathematics (+ *broken* English) is the only language we can use. So, *unfortunately*, we need to learn mathematics! (and English... 🙄) But it means we are *fortunate*: we have a language to describe the world. In fact, physics have developed together with mathematics. We understand physics better if we study mathematics well!

When you do DIY, you need to use the tools properly, smoothly, and accurately. It is the same here; when we use math as a tool, it must be **logical**, **quick**, and **accurate**. To be logical, you must think each problem carefully; otherwise your discussion might be incomplete and you can't convince others. To be quick and accurate, you must "*drill*" repeatedly, as if top NBA players do shooting practice almost everyday. You should reach the point where basic calculations feel automatic.

This is not a math textbook. **This is rather** a math "drill" book. Rigorous proofs are mostly omitted, but drill problems are the core of this document, through which students are expected to achieve better conceptual understanding.

Target of this document

This document is primarily for first-year undergraduate students in their second semester, who

- completed basic calculus (differentiation and integration),
- are going to major in physics or physical science, and
- do not hesitate to do repetitive practice to achieve better conceptual understanding.

How to use this document

1. Buy a A4-sized paper notebook.

Digital notebooks on tablets are not suitable (less effective).

2. Read the text and solve **Quizzes**.

The texts are kept short, so please read all the text carefully. Quizzes should be solved while reading, but you don't have to write down the solution process for Quizzes.

3. Solve **Drill Problems**, until you feel "I can solve them *fluently and accurately*".

This is to improve your fluency ($\neq$ speed) and accuracy. Write down solution process clearly.

4. Solve other **Problems**.

This is to have better conceptual understanding, but also to develop your writing skills. Write down the solution process clearly *as an English text*, taking care of *logical completeness*.

Optionally, you are encouraged to prepare a reference book, so that you can consult it when you want to know more. Sho supposes [Boas] listed below is the best option.



This document is licensed under [the Creative Commons CC-BY-NC 4.0 International Public License](https://creativecommons.org/licenses/by-nc/4.0/).

You may use this document only if you do in compliance with the license.

Visit <https://github.com/misho104/LecturePublic> for further information, updates, and to report issues.

Problems

 **Drill problems**, marked with , are simple and easy. They help you practice the basic calculations until you can do them fluently and accurately. If they feel too easy for you, feel free to skip some of them.

- ★ These are **minimal basic** problems. To pass this course, you should be able to solve all of them. If you struggle with them, please come talk to Sho—don't wait until the exam!
- *** These **basic** problems are the same difficulty as the ★-problems. Solve these for extra practice, to check that you really understand the ★-problems and were not just lucky.
- ** These are **intermediate** problems. Solve these if you are aiming for an A+. Note also that, at this level, you are asked to write answers carefully with clear explanations.
- * These are **advanced** problems, beyond the scope of the lecture, for motivated students who are interested in research activities, a top-level engineering career, or demanding analytical-thinking work.

Bibliography

This document is written under the influence of the following references.

- [The Math Book by Hal Tasaki](#) (数学：物理を学び楽しむために, in Japanese)
A free book with full description, but unfortunately in Japanese. Topic selections and the depth of descriptions are based on this reference.
- [First-Year Math Practice](#) (in Japanese)
This document is grounded in a “learn steadily” approach (“じっくり” in Japanese), which originates from Prof. Gocho and Prof. Kiyono, who were instructors of Sho in his freshman. Problems and rigorous math descriptions are taken from their works.
- **[Boas]** Mary L. Boas, *Mathematical Methods in the Physical Sciences*, 3rd ed., Wiley, 2006.
- **[AWH]** George B. Arfken, Hans J. Weber, and Frank E. Harris,
Mathematical Methods for Physicists, 7th ed., Academic Press, 2023.
These two books are widely used in physics departments around the world. [Boas] seems more friendly to beginners, while [AWH] seems more detailed and complete. Sho recommends you to study [Boas] *in parallel with* this document, using it as a reference, and to consult [AWH] when you want more details or more exercises.

Other references are:

- [1] Bureau International des Poids et Mesures, “The International System of Units (SI),” BIPM. [Online]. Available: <https://doi.org/10.59161/AUEZ1291>

Symbols

SI Units

Historically, different civilizations used different units. Today, science and engineering worldwide use the **SI units** (Système International d’Unités)—the single international standard. It is built from seven **SI base units**:

Table 1: SI base units and their symbols for dimension [1]. Recall that units are case-sensitive.

quantity	symbol and name	symbol for dimension
time	s (second)	T
length	m (meter)	L
mass	kg (kilogram)	M
electric current	A (ampere)	I
temperature	K (kelvin)	Θ
amount of substance	mol (mole)	N
luminous intensity	cd (candela)	J

Mathematical Notations

Chapter 1		Chapter 3	
$\mathbb{C}$	complex numbers	$\wedge$	logical “and”
$\mathbb{R}$	real numbers	$\vee$	logical “or”
$\mathbb{Q}$	rational numbers	$\neg$	logical “not”
$\mathbb{Z}$	integers	$\implies$	implication
$\mathbb{N}^0$	natural numbers (0, 1, 2, ...)	$\iff$	logical equivalence
$\mathbb{N}^+$	natural numbers (1, 2, ...)	$\forall$	for all
$x \in A$	a member of	$\exists$	there exists
$A \cup B$	union	$:=$	is defined as
$A \cap B$	intersection		
$A \setminus B$	set difference		
$f'(x)$	derivative $\left(\text{also } \frac{df(x)}{dx} \right)$		
$f^{(n)}(x)$	n -th derivative $\left(\text{also } \frac{d^n f(x)}{dx^n} \right)$		

Greek symbols

A	α	alpha	H	η	eta	N	ν	nu	T	τ	tau
B	β	beta	Θ	θ	theta	Ξ	ξ	xi	Y	υ	upsilon
Γ	γ	gamma	I	ι	iota	O	$\omicron$	omicron	Φ	$\varphi(\phi)$	phi
Δ	δ	delta	K	κ	kappa	Π	π	pi	X	χ	chi
E	ε	epsilon	Λ	λ	lambda	P	ρ	rho	Ψ	ψ	psi
Z	ζ	zeta	M	μ	mu	Σ	σ	sigma	Ω	ω	omega

Gray-outed ones are almost never used, but others are almost always used. You need to write them so that **others can distinguish each from others**, but Sho thinks Japanese- or Chinese-speakers are OK with it, as we everyday read and write 日 vs 日 vs 白 or 土 vs 土 vs 工 properly!

Quiz

Q1. Read out the following words.

- | | | |
|---|----------------------------------|---|
| (1) α -particle | (2) β -decay | (3) Γ -function |
| (4) ε - δ definition | (5) angle θ and φ | (6) μ - and τ -leptons |
| (7) $\psi(x)$ and $\varphi(x)$ | (8) ζ and ξ | (9) metric $\eta_{\mu\nu}$ |
| (10) X-ray and γ -ray | (11) Ω -baryon | (12) $\varepsilon_{\mu\nu\rho\sigma}$ -tensor |

Problems

 0.1 Write the following letters so that people can distinguish each from others.

- | | | | |
|--------------------|------------------------------------|---------------------|-------------------------|
| (1) a, α | (2) b, β | (3) c, C | (4) e, E, ε |
| (5) ζ, ξ | (6) $\Theta, \theta, Q, \sigma, 6$ | (7) $g, q, 9$ | (8) $i, l, 1, I$ |
| (9) k, K, κ | (10) m, μ | (11) p, ρ | (12) s, S |
| (13) t, τ, T | (14) u, v, ν | (15) w, W, ω | (16) x, X, χ |

Lecture Plan

The content is designed for 150-minute $\times$ 14-week lectures, as it is originally for a lecture *Mathematics for Fundamental Physics* (基礎物理數學) in National Sun Yat-sen University (國立中山大學).

1. Derivative
2. Units. Significant Digits.
3. Logic. Theorems. a^x and $\log_a x$.
4. Logic. Theorems. a^x and $\log_a x$.
5. Vectors are arrows.
6. $a \cdot b$ and $a \times b$
7. Matrices (rotation, reflection, scaling)
8. Complex numbers (polar form, Euler's formula, multi-valuedness)
9. Complex vectors, Hermitian inner product, bra-ket notation
10. Vector calculus (grad, div, rot; example of point charge)
11. ODE basics
12. ODE basics
13. Probability and error analysis
14. Probability and error analysis

♣TODO♣ a bit on (abstract) vector space to prep for QM?

Contents

Preface	2
Bibliography	3
Symbols	4
Lecture Plan	6
Index	9
1 Derivative (Review)	10
1.1 The first step	10
1.2 Typical Pitfalls	11
1.3 Taylor expansion	13
1.4 Mathematical Notation	15
2 Units and Significant figures	17
2.1 Physical quantity	17
2.2 Units	18
2.3 Significant figures	20
2.4 Calculation with Significant figures	21
3 Logic	24
3.1 Basic Logics	24
3.2 Implication $\Rightarrow$	25
3.3 Equivalent Statements—What does “solve” mean?	26
3.4 A few more notes about logic and proof	30
3.5 Facts, Definitions, Assumptions, and Conclusions	31
4 Power, Exponential, and Logarithm	34
4.1 Powers	34
4.2 Exponential functions and Logarithmic functions	37
4.3 Napier’s number and Natural logarithm	38
4.4 Exponential Growth and Decay	40

List of Theorems and Definitions

Example 1.1	13	Example 4.2	35
Statement 1.2	14	Example 4.3	35
Statement 2.1	18	Example 4.4	35
Example 2.2	22	Example 4.5	36
Example 2.3	22	Definition 4.6 ...	36
Example 3.1	25	Definition 4.7 ...	36
Definition 3.2 ...	26	Theorem 4.8	37
Theorem 3.3	30	Definition 4.9 ...	37
Example 3.4	30	Theorem 4.10 ...	38
Example 3.5	31	Theorem 4.11 ...	38
Definition 4.1 ...	35		

List of tables

Table 1 SI base units and their symbols for dimension [1]. Recall that units are case-sensitive. . . 4

List of figures

Index

absolute uncertainty	17	or operator	25
and operator	25	order	12
base	34	physical quantity	17
central value	17	power	34
conjunction	→ see <i>and</i>	proof	24
constant	11	proof by contradiction	31
contrapositive	30	proof by contrapositive	30
converse	30	proposition	33
counterexample	25	quantifier	29
de Morgan s theorem	29 , 30	radian	12
definition	13	rate of change	13
derivative	10	rational number	15
dimensionless	18	relative uncertainty	17
disjunction	→ see <i>or</i>	rounding	21
e	→ see <i>Napier's number</i>	scientific notation	20
empty set	16	set	16
equivalent	26	set difference	16
error	→ see <i>uncertainty</i>	SI base unit	4
exponent	34	SI unit	4
exponential decay	40	significant figure	17 , 20
exponential function	37	sufficient condition	27
exponential growth	40	syllogism	33
false	24	symbols for dimension	19
half-life	40	tangent line	13
Hyperbolic function	40	transfinite induction	31
iff	27	trigonometric function	13
implication	26	true	24
infinite descent	31	truth table	25
integer	15	uncertainty	17
intersection	16	absolute	17
inverse	30	relative	17
logarithm	37	union	16
mathematical induction	31	unit	17
Napier s number	38	variable	11
natural logarithm	38		
necessary condition	27		
negation	→ see <i>not</i>		
not operator	25		

Chapter 1

Derivative (Review)

1.1 The first step

Birds sing, fish swim, flowers bloom, stars twinkle, and university students calculate derivatives. Let us begin with basic **derivatives**. Try the next quiz—and note how long it takes.

Quiz

Q1. Calculate the first derivatives of the following functions, **measuring how many minutes it takes**.

(1) $(x + 1)^3$

(2) $\tan x$

(3) $2 \cos^2 x$

(4) $2 \cos x^2$

(5) $x^3 \cos x$

(6) $\frac{3 \sin x}{x}$

(7) $x^{3/2}$

(8) $\sqrt{\sin x}$

- Less than two minutes? Amazing! It is as fast as Sho!
- Less than four minutes? Great, it is exactly as Sho anticipates. Please continue your effort!
- Even if you took more than four minutes, do not worry. At least you don't make any mistakes. A little more practice will help you. Try more problems in [the Derivative Boot Camp](#).^{#1}
- If you make any mistakes in these eight calculations, then it is a serious issue—just as serious as forgetting how to do the calculations at all. In university, you will perform similar calculations more than 100 times. You need to be both fast and accurate.



At university, students often underestimate the importance of basic calculations. In physics, simple calculations appear constantly, so your speed and accuracy directly affect how well you follow lectures, how efficiently you study, and ultimately your grade. Both can be improved with practice.

This course will help you strengthen these basic calculations—some of which you already know from high school—while also introducing new topics that are essential for physics. You will solve many problems and drills, just as an athlete repeats the same move hundreds of times to master it.

This document (this lecture course) is designed for first-year students in their second semester, **who have already studied derivatives well** in their first-semester calculus. As we do not cover the details of derivatives here, if you are not confident with derivatives, please first study [the Derivative Boot Camp](#).

^{#1}Visit <https://misho104.github.io/LecturePublic> and find gp1_boot1_deriv_true.pdf.

1.2 Typical Pitfalls

1.2.1 Notations

Consider a function $f(x)$. As we vary the value of x , x is called a **variable**. For a function $f(x)$, there are several equivalent ways to write its derivative:

$$f'(x) = \frac{df}{dx}(x) = \frac{df(x)}{dx} = \frac{d}{dx}f(x) \quad [\text{All means the same thing}]. \quad (1.1)$$

Similarly, the value of $f'(x)$ at a specific point $x = 3$ can be written as

$$f'(3) = f'(x)\Big|_{x=3} = \frac{df}{dx}\Big|_{x=3} = \frac{df}{dx}(3) = \frac{d}{dx}f(3), \quad (1.2)$$

and they are all equivalent. For example, if $f(x) = 3x^2 + 6$, then $f'(x) = 6x$ and $f'(3) = 18$.

Most of students get confused when a **constant** a appears. If a is *declared as a constant*, we can define a function such as $g(x) = ax^2 + 2a$, for which

$$g'(x) = 2ax. \quad (1.3)$$

We can evaluate $g'(x)$ at $x = a$. The result is $g'(a) = 2a^2$, written as

$$g'(a) = g'(x)\Big|_{x=a} = \frac{dg}{dx}(a) = \frac{d}{dx}g(a) = 2a^2. \quad (1.4)$$

Be careful: The notation $\frac{d}{dx}g(a)$ does **not** mean $\frac{d}{dx}[g(a)] = \frac{d}{dx}(a^3 + 2a) = 0$.

Fail safe note: If you can't see Eq. (1.3), try setting $a = 3$. Then, you notice $g(x)$ is the same function as $f(x)$, and $g'(x)$ should equal $f'(x)$.

Quiz

Q2. Let a be a constant and $g(x) = ax^2 + 2a$. Calculate

- | | | | |
|-------------|-------------|--------------|------------------------|
| (1) $g(1)$ | (2) $g(a)$ | (3) $g(0)$ | (4) $g(7)$ |
| (5) $g'(1)$ | (6) $g'(a)$ | (7) $g''(2)$ | (8) $\frac{d}{dx}g(7)$ |

One more bad news. *Physicists are often lazy* and write $f(x)$ as f . If $f(x) = 2x^2 + 1$, we may write $f' = 4x$ and $f'' = 4$. For $g(t) = 2t^2 - 1$, you may write $g' = 4t$ and $g'' = 4$. Therefore, when you see a function f , you must **identify its variable from the context**.

In physics, a variable can be a function of another variable. The kinetic energy $K(v) = mv^2/2$ is a good example. Its variable v is a function of time t —we write this fact by $v = v(t)$ —and thus

$$K(v) = \frac{1}{2}mv^2 \quad \text{but also} \quad K(t) = K(v(t)) = \frac{1}{2}mv(t)^2. \quad (1.5)$$

Now, what does K' mean? —It is ambiguous and we must avoid such notation. Even though, if you see it in a textbook, you need to guess the author's intention from the context.

Advanced note: This rewrite, $K(t)$ and $K(v)$, is common in physics. However, mathematicians do not like this practice because they consider K a unique object: a function should always mean the same rule applied to its input. If they see $K(v) = mv^2/2$, they would say [K is an object that converts its input into $m/2 \times (\text{input})^2$, and thus if you feed t into K , you must get $K(t) = mt^2/2$]. This is of course not what we mean. For example, imagine $v(t) = \alpha t$ with α a constant. Then

$$\text{physicists: } K(v) = \frac{1}{2}mv^2, \quad K(t) = \frac{1}{2}m\alpha^2 t^2; \quad (1.6)$$

but mathematicians would say, [since it returns $m\alpha^2/2 \times (\text{input})^2$, it is a different object] and hence use a different name for it, such as $\tilde{K}$:

$$\text{mathematicians: } K(v(t)) = \frac{1}{2}m \times v(t)^2 = \tilde{K}(t) = \frac{1}{2}m\alpha^2 t^2. \quad (1.7)$$

Both conventions are reasonable, and eventually you will get used to both.

Higher-order derivatives is written as

$$\frac{d}{dx} \left(\frac{df}{dx} \right) = \frac{d^2 f}{dx^2} = f''(x) = f^{(2)}(x), \quad \frac{d}{dx} \left(\frac{d}{dx} \left(\frac{df}{dx} \right) \right) = \frac{d^3 f}{dx^3} = f'''(x) = f^{(3)}(x), \quad (1.8)$$

etc. Please be careful on the position of “2” and “3”.

1.2.2 Radian and trigonometric functions

At university, angles are almost always measured in **radians**:

$$360 \text{ degree (or: } 360^\circ) = 2\pi \text{ radian (or: } 2\pi \text{ rad)} \quad (1.9)$$

Furthermore, we usually omit “radian” (because we are lazy!). So,

$$\text{a right angle is } \pi/2. \quad \text{The sum of the interior angles of a triangle is } \pi. \quad (1.10)$$

Quiz

Q3. Express the following angles in radians, and radians in angles.

- | | | | | |
|-----------------|----------------|----------------|---------------|-----------------|
| (1) 180° | (2) 45° | (3) 60° | (4) 90 | (5) -30° |
| (6) 1° | (7) 1 | (8) -0.3 | (9) x° | (10) x |

Why do insist on radians? The answer comes from the derivative formula

$$\frac{d}{dx} \sin x = \cos x. \quad (1.11)$$

Because we have chosen radians as the standard, $(\sin x)'$ becomes this simple.

Advanced note: The simpleness of Eq. (1.11) originates in the fact $\sin(0.01 \text{ rad}) \approx 0.01$. If we used degrees, we would have $\sin(0.01^\circ) \approx 0.01 \times 0.017453$ and everything is messed up with this number 0.017453.

There are a few remarks in the notation of **trigonometric functions**:

$$\begin{aligned} \sin^2 x &\neq \sin x^2. & \text{Namely, } (\sin x)^2 = \sin^2 x &\neq \sin x^2 = \sin(x^2). \\ \tan^{-1} x &\neq \frac{1}{\tan x}. & \text{Namely, } \tan^{-1} x = \arctan x &\neq (\tan x)^{-1} = \frac{1}{\tan x} = \cot x. \end{aligned} \quad (1.12)$$

The following expressions are not incorrect but confusing;

$$\sin^{-2} x, \quad \sin^{1/2} x, \quad \sin(x)^2, \quad \sin(x+1)^2, \quad \dots$$

We should avoid ambiguity, so please *never* use confusing these notations. Sho thinks we should use $\sin^k x$ only for $k = 2, 3, 4, \dots$, and use $\arcsin x$ instead of $\sin^{-1} x$.

1.2.3 Several interpretations of derivatives

When you, physics learners, discuss $f'(t)$, you should have the following three interpretations:

- Regarding t as the time, $f'(t)$ is the **rate of change** of $f(t)$ per unit time. If $f'(t) > 0$, the function f is increasing at the time t , while $f'(t) < 0$ means it is decreasing.
- Consider a graph of $f(t)$. Then, $f'(t)$ is the slope of the **tangent line** to the graph at the point t .
- Mathematically, $f'(t)$ is defined by (Check that they are equivalent.)

$$f'(t) := \lim_{\Delta t \rightarrow 0} \frac{f(t + \Delta t) - f(t)}{\Delta t} = \lim_{h \rightarrow 0} \frac{f(t + h) - f(t - h)}{2h} = \lim_{s \rightarrow t} \frac{f(s) - f(t)}{s - t}. \quad (1.13)$$

If you are unsure of them, please consult your first-year Calculus textbooks for further information.

Quiz

Q4. What is the definition of $f'(x)$? Explain.

In physics, it is important to memorize and understand the **definition** of each concept.

1.3 Taylor expansion

Reviewing the definition of $f'(x)$, we have

$$f'(x) := \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x}, \quad f'(a) := \lim_{\delta \rightarrow 0} \frac{f(a + \delta) - f(a)}{\delta}. \quad (1.14)$$

The second equation is interpreted as follows:

$$\text{If } \delta \approx 0, \quad f'(a) \approx \frac{f(a + \delta) - f(a)}{\delta}, \quad \text{i.e., } f(a + \delta) \approx f(a) + \delta f'(a) \quad (1.15)$$

and this interpretation is useful in the following example:

Example 1.1

Find the approximate value of the following expressions without using calculators. Then, check your answer with a calculator.

1. $\sqrt{1.002}$

2. $(1.002)^{10}$

3. $\sqrt{4.004}$

Solution

(1) Apply Eq. (1.15) for $f(x) = \sqrt{x}$, $a = 1$, and $\delta = 0.002$. Then,

$$f'(x) = \frac{1}{2\sqrt{x}}, \quad f'(a) = f'(1) = \frac{1}{2}, \quad f(a + \delta) \approx f(a) + \delta f'(a) = 1 + \frac{\delta}{2} = \underline{1.001}.$$

(2) Doing the same thing for $g(x) = x^{10}$ with $a = 1$ and $\delta = 0.002$,

$$g'(x) = 10x^9, \quad g'(a) = g'(1) = 10, \quad g(a + \delta) \approx g(a) + \delta g'(a) = 1^{10} + 10\delta = \underline{1.02}.$$

(3) Doing the same thing for $h(x) = \sqrt{x}$, $a = 4$, and $\delta = 0.004$, we have

$$h'(4) = \frac{1}{4}, \quad h'(4.004) \approx h(4) + 0.004 \times \frac{1}{4} = \underline{2.001}.$$

Advanced note: “Solutions” contain not only the answer but also how you reached the answer. You are, of course, asked to write such explanations when you solve problems.

Namely, with this technique, you can find **the value of $f(x)$ around a point $x = a$** . This technique, Eq. (1.15), is a special case of Taylor’s theorem, which is discussed in **♣TODO♣???**.

Quiz

Q5. Find the approximate value of the following expressions without using calculators. Then, check your answer with a calculator.

(1) $\sqrt{1.001}$

(2) $(1.0001)^{30}$

(3) $\frac{1}{1.001}$

To summarize, we have the following statement:

Statement 1.2 (Taylor expansion: basic)

For a physicists-friendly function $f(x)$,

$$f(a + \delta) \approx f(a) + \delta f'(a), \tag{1.16}$$

or as an equivalent expression,

$$f(x) \approx f(x_0) + (x - x_0)f'(x_0). \tag{1.17}$$

Quiz

Q6. Check these two equations are equivalent. You will be ready to use either of them in the future.

Q7. Review how to derive Eq. (1.16) and summarize it in your own words.

Problems

 **1.1** Calculate the following expression.

- (1) $\sin 30^\circ$ (2) $\cos 45^\circ$ (3) $\cos 120^\circ$ (4) $\sin 150^\circ$ (5) $\tan 180^\circ$
 (6) $\cos 0$ (7) $\sin 2\pi$ (8) $\cos \pi$ (9) $\sin(\pi/3)$ (10) $\cos(-\pi/4)$
 (11) $\tan(2\pi/3)$ (12) $\sin(\pi/6)$ (13) $\tan(5\pi/6)$ (14) $\cos(-\pi/6)$ (15) $\tan(-\pi/2)$

 **1.2** For the following $f(x)$, calculate $f'(x)$, $f''(x)$, $f^{(3)}(x)$, $f^{(4)}(x)$, and $f^{(5)}(x)$.

- (1) x^3 (2) $\sin 2x$ (3) e^{2x} (4) $\ln 2x$

 **1.3** Find the approximate value for the following expressions without calculators. Then, check your answer with a calculator.

- (1) $(1.002)^7$ (2) $(2.002)^7$ (3) $(0.998)^4$ (4) $(10.01)^3$ (5) $1 \div 0.998$
 (6) $\sqrt{0.998}$ (7) $\sqrt{4.001}$ (8) $(0.999)^{1/2}$ (9) $(0.999)^{-1/2}$ (10) $(0.999)^{-3/2}$
 (11) $\frac{1}{(2.02)^2}$ (12) $\tan 0.001$ (13) $\cos 0.002$ (14) $\sin 0.003$ (15) $\ln 1.001$

****1.4** Find the approximate value for the following expressions without calculators. Then, check your answer with a calculator.

- (1) $\sqrt[3]{1.001}$ (2) $\sqrt[3]{8.012}$ (3) $1.001^{0.5}$ (4) $1.001^{-0.9}$ (5) $\sin(3.14)$
 (6) $\cos(1.57)$ (7) $\sin 1^\circ$ (8) $\cos 61^\circ$ (9) $\ln 0.999$ (10) $\log_{10}(10.01)$
 (11) $\sqrt{1.001} \sin 0.001$

Fail safe note: If you are not sure about $\sqrt[3]{1.001}$, $(1.001)^{-0.9}$, etc., study [Chapter 4](#) first.

****1.5** For the following $f(x)$, calculate $f^{(n)}(x)$ for all positive integers $n = 1, 2, 3, \dots$

- (1) x^{10} (2) $\sin 2x$ (3) e^{2x} (4) $\ln x^2$ (5) $\sqrt{x}$

***1.6** Write down the definition of the second derivative $f''(x)$. Repeat the discussion of Eq. (1.15) to find the expansion $f'(a + \varepsilon) \approx f'(a) + \varepsilon f''(a) + (\varepsilon^2/2)f''(a)$.

Advanced note: The two equations in Eq. (1.14) have similar but different meanings: the first one defines a new function $f'(x)$, while the second one defines a number that is eventually equal to $f'(x)|_{x=a}$. Anyway, we don't care the difference.

1.4 Mathematical Notation

As *physicists are lazy*, we usually use the following notation:

$x \in \mathbb{C}$	means “ x is a complex number”.	$x \in \mathbb{Z}$	means “ x is an integer ”. ^{#2}
$x \in \mathbb{R}$	means “ x is a real number”.	$x \in \mathbb{N}^+$	means “ x is a positive integer”.
$x \in \mathbb{Q}$	means “ x is a rational number ”.	$x \in \mathbb{N}^0$	means “ x is a non-negative integer”.

^{#2} $\mathbb{Z}$ comes from “Zahlen” (Zahl).

The symbol “ $\in$ ” means “*is a member of*”, and $\mathbb{N}^+$ means a **set** of positive integers, $\{1, 2, 3, 4, \dots\}$. So,

$x \in \mathbb{N}^+$ and $x \in \{1, 2, 3, 4, 5, \dots\}$ means the same thing: “ x is a positive integer”.

Remark: In mathematics, “natural numbers” usually mean $\mathbb{N}^0$. Meanwhile, physicists and other people tend to think natural numbers mean $\mathbb{N}^+$.

Quiz

Q8. What is the difference between $\mathbb{N}^+$ and $\mathbb{N}^0$?

Q9. What is the difference between $\mathbb{N}^0$ and $\mathbb{Z}$?

Q10. What is “rational numbers”? What is the difference between $\mathbb{Z}$ and $\mathbb{Q}$?

We write “ x is a positive number” by $x \in \mathbb{R}, x > 0$ but also by $x > 0$. Similarly,

$x \in \mathbb{R}, x \leq 0$ or $x \leq 0$ means “ x is a non-positive number”.

Notice that, when we say “positive” or “negative”, we implicitly assume the number is real.

Advanced note: This is because we cannot compare complex numbers with 0. The operator $>$ is defined only for real numbers. When we write $a > b$, we implicitly consider that a and b are real numbers.

Problems

****1.7** The **intersection** of two sets A and B is a set made by elements both in A and B ; we write it by $A \cap B$. The **union** of A and B is a set made by elements in A , B , or both; we write it by $A \cup B$. The **set difference**, $A \setminus B$, is a set made by elements in A but not in B .

For example, let $A = \{1, 2, 3\}$, $B = \{2, 3, 4\}$, and $C = \{2, 4\}$. Then,

$$A \cap B = \{2, 3\}, \quad A \cup B = \{1, 2, 3, 4\}, \quad A \setminus C = \{1, 3\}, \quad B \setminus C = \{3\}, \quad C \setminus B = \{\},$$

where $\{\}$ (or $\emptyset$) means the **empty set**, the set without any elements.

(1) Write the following sets.

$$(a) A \cap C \quad (b) B \cap C \quad (c) A \cup C \quad (d) B \cup C \quad (e) A \cup B \cup C$$

$$(f) B \setminus A \quad (g) C \setminus A \quad (h) C \setminus B \quad (i) (B \setminus A) \cup A \quad (j) \mathbb{N}^0 \setminus \mathbb{N}^+$$

(2) Express the following statements only with $\mathbb{R}$, $\mathbb{Q}$, $\mathbb{Z}$, $\mathbb{N}^+$, $\mathbb{N}^0$, $\cap$, $\cup$, $\setminus$, and “ $x \in$ ”.

(a) x is a rational number.

(b) x is an irrational number.

(c) x is a rational number but not an integer.

(d) x is a non-positive integer.

(e) x is a negative integer.

(f) x is a non-zero integer.

Units and Significant figures

A **physical quantity** is usually not just a number, but also has a **unit** and an **uncertainty** (error). For example, “ $(72 \pm 2) \text{ kg}$ ” has a number “72” and a unit “kg”, which compose the **central value** “72 kg”. It also has uncertainty because every measurement has an uncertainty. In experimental physics, uncertainty is usually more important than the central value, and thus uncertainty handling is a fundamental skill of physicists.

In this chapter, we first review physical quantities and how to handle units. Then, we discuss **significant figures**, a simple and convenient method to express the uncertainty. Further discussions on uncertainty analysis are given in Chapter ♣[TODO](#)♣chapter.

2.1 Physical quantity

The expression “ $(72 \pm 2) \text{ kg}$ ” means the central value is 72 kg and the **absolute uncertainty** is 2 kg. Accordingly, its **relative uncertainty** is given by $(2 \text{ kg})/(72 \text{ kg}) = 0.028$. Let’s see other examples.

	unit	central value	absolute uncertainty	relative uncertainty
$10\text{ m} \pm 1\text{ cm}$	m (meter)	10 m	1 cm = 0.01 m	0.001 (or 0.1%)
$1.6\text{ A} \pm 0.04\text{ A}$	A (ampere)	1.6 A	0.04 A	0.025 (or 2.5%)
$(5 \pm 1) \times 10^{-3}\text{ C}$	C (coulomb)	0.005 C	0.001 C	0.2 (or 20%)
$(50 \pm 1)\text{ m/s}$	m/s	50 m/s	1 m/s	0.02 (or 2%)

Remark: Upper- and lowercase letters are distinguished. “M” **does not** mean “meter”. The unit “A” is “ampere”, not “Ampere”. The unit “coulomb” is “C”, not “c”.

Quiz

Q1. For each of the following values, find its unit, central value, absolute uncertainty, and relative uncertainty.

- (1) $50 \text{ kg} \pm 500 \text{ g}$ (2) $(0.05 \pm 0.001) \text{ A}$
(3) $(1.6 \pm 0.001) \times 10^{-19} \text{ C}$ (4) $72 \text{ km/h} \pm 1 \text{ m/s}$

Q2. The next passage has six (6) errors in the use of uppercase and lowercase letters. Find them out.

In the SI system, temperature is expressed in k (kelvin), a unit named after Lord Kelvin. The unit of force is N (newton), named after the british scientist Isaac newton. The units A (Ampere) and c (coulomb) are named after French scientists. In contrast, Kg (kilogram) is not named after a person. , but comes from Greek.

As an undergraduate student, you need to follow the following rules:

Statement 2.1 (Rules for Physical Quantities: Basic)

1. Numbers are always with units, even in calculations.
2. Include units in a symbol.
3. For physical quantities, use decimals (e.g., 1.6 A). Do not use fractions like (8/5) A.
4. Use significant figures to express the measurement precision.

Be careful: We use *decimals for physical quantities* and *fractions for mathematical concepts*. Since this lecture is mainly on math and math concepts, you will use decimals **♣TODO♣only in this section**.

2.2 Units

When we educate kids, we write “my height is cm”, or “my height is h cm”, where $h = 172$ is just a number. But this is not nice! We want to convert the units freely and write equations such as $1.72 \text{ m} = 172 \text{ cm} = 0.00172 \text{ km}$. So, **we always include units in symbols** $h = 172 \text{ cm}$. Then

we can write: $h = 172 \text{ cm} = 1.72 \text{ m} = 0.00172 \text{ km} = 1.82 \times 10^{-16} \text{ light-year}$.

Similarly, if $m = 110 \text{ g}$ and $g = 9.8 \text{ m/s}^2$,

we may write: $w = mg = 110 \text{ g} \times 9.8 \text{ m/s}^2 = 1.1 \text{ kg} \cdot \text{m/s}^2$
 but not: $w = mg = 0.11 \times 9.8 = 1.1 \text{ kg} \cdot \text{m/s}^2$

This second equation is incorrect because m is not equal to 0.11; m is equal to 0.11 kg or 110 g.

Remark: Usually, physicists use upright fonts for units and *italic fonts* for quantities. For example, m , T , and C are quantities, which can be mass, temperature, capacitance, etc. Meanwhile, m, T, and C are units: meter, tesla, and coulomb, respectively.

Quiz

- Q3.** (1) If $mr\omega^2 = 10.0 \text{ kg} \cdot \text{m/s}^2$, $m = 5.0 \text{ kg}$, and $r = 1.0 \text{ m}$, what is ω ?
- (2) If $v_0 t + \frac{1}{2}at^2 = 5.0 \text{ m}$, where $a = -4.0 \text{ m/s}^2$ and $v_0 = 7.0 \text{ m/s}$, what is t ?
- (3) At time $t = 0$, a particle is at $(x, y) = (2.0 \text{ m}, 0)$. It moves with a constant velocity $(v_x, v_y) = (3.0, 4.0) \text{ m/s}$. What is its position at $t = 1.0 \text{ s}$?

Every physical concept has its own unit. For example, speed has m/s, acceleration has m/s^2 , and energy has $\text{kg} \cdot \text{m/s}^2 = \text{N} \cdot \text{m} = \text{J}$. The table below lists the quantities you have learned. Notice that angle (rad) has no dimension. It is called a **dimensionless** quantity.

concept	dimension	typical unit
speed	$L T^{-1}$	m/s
acceleration	$L T^{-2}$	m/s^2
linear momentum	$M L T^{-1}$	$kg \cdot m/s$
force	$M L T^{-2}$	$kg \cdot m/s^2$ (= N)
energy / work	$M L^2 T^{-2}$	$kg \cdot m^2/s^2$ (= J)
frequency	T^{-1}	1/s (= Hz)
angle	1 (dimensionless)	rad

Advanced note: To understand why angle has no dimension, you may consider the definition of the radian: it is defined as the ratio of the arc length to the radius, so the units cancel (meter / meter = 1).

Here, please do not write: The force is $kg \cdot m^2/s^2$. This is also called newton (N)., because we cannot mix units and concepts. A correct (and easy) way is to use English words:

the unit of the force is $kg \cdot m/s^2$ or the force has the unit $kg \cdot m/s^2$.

Similarly, the following statement is incorrect because it mixes concepts and units:

Since $m = kg$ and $g = m/s^2$, $mg = kg \cdot m/s^2 = N$.

Instead, you need to write

Since m has a unit of kg and g has a unit of m/s^2 , mg has a unit of $kg \cdot m/s^2 = N$.

Notice the last equation: units can be equal to other units, so we may write $N = J/m = kg \cdot m/s^2$

$m = J/N$ $s = \sqrt{kg \cdot m/N}$. These are scientifically correct equations.

Remark: We sometimes use informal notations, such as $[F] = N$ or $F \sim N$ to express “the unit of F is N (newton)”. Still, using an equal symbol ($=$) is **not allowed**. Namely, $F = N$ is always incorrect.

Quiz

Q4. What are the units of the following concepts? Write English sentences to explain it.

(1) speed (2) velocity (3) force (4) energy (5) power

Advanced note: In very formal situations, we use **symbols for dimensions** (see Table 1 on page 4):

Since $\dim(m) = M$ and $\dim(g) = L T^{-2}$, $\dim(mg) = M L T^{-2}$, with the operator “dim” [1]. However, it seems too complicated for most situations.

Problems

★2.1 Express the following units only with SI base units, e.g., $N = \text{kg} \cdot \text{m}/\text{s}^2$.

- (1) W (watt) (2) Pa (pascal) (3) J (joule) (4) C (coulomb)

***2.2 Find a few more examples of dimensionless quantities in physics.

**2.3 Use “dim” notation to express the dimensions of the following quantities. For example, if v is speed, then $\text{dim}(v) = L T^{-1}$.

- | | |
|--------------------------|--------------------------------------|
| (1) speed v | (2) force F |
| (3) angular momentum L | (4) electric charge Q |
| (5) resistance R | (6) specific heat capacity c |
| (7) Planck constant h | (8) fine-structure constant α |

2.3 Significant figures

In researches, we need to treat uncertainties in the method given in [♣TODO♣chap](#). However, most of lectures use **significant figures** to express the accuracy of a value, so that students becomes familiar with the concept of uncertainties. For example,

3.14	means the last digit “4” is uncertain. It can be 3.16, 3.15, 3.11, 3.10,
3.141	means “3.14” is for sure but it can be 3.143, 3.142, 3.140, 3.139,
3.1415	might be 3.1416, 3.14153, etc., but the writer is sure <i>it is not</i> 3.145 or 3.135.

We write the “uncertain digit” in [a different style](#) for clarity. You should notice the difference between:

1.0000	meaning the last digit “0” is uncertain so it may be 1.0002, 1.0001, 0.9998,
1.00	meaning it can be 1.03 or 0.99, but the writer is sure <i>it is not</i> 1.2 or 0.8.

It is usually convenient to use **scientific notation**:

0.000123	is OK but we prefer	1.23×10^{-4} .	Both mean the value can be 1.22×10^{-4} etc.
0.00100	is OK but we prefer	1.00×10^{-3} .	
29979	is OK but we prefer	2.9979×10^4 .	

Fail safe note: $10^3 = 1000$ and $10^{-3} = 1/10^3 = 1 \div 1000$. Go to [♣TODO♣chap](#) for details!

We need to *avoid ambiguous notations*. Namely,

2040	is ambiguous and not nice. We are not sure the author means 2.04×10^3 or 2.040×10^3 .
120	is not nice. We are not sure the author means It may mean 1.20×10^2 or 1.2×10^2 .
42000	is not nice because it has four ways to interpret the author’s intention.

Quiz

Q5. What are the four interpretations of 42000? Write them in scientific notation.

We use **rounding-to-the-nearest** (四捨五入) when necessary. For example, if you need to convert to three significant figures, it will be

$1.234 \rightarrow 1.23$, $1.235 \rightarrow 1.24$, $31.98 \rightarrow 32.0$, $1234 \rightarrow 1.23 \times 10^3$, and so on.

Quiz

Q6. Round the following numbers to three significant figures.

- | | | | |
|------------|---------------|------------|-------------------------|
| (1) 41.11 | (2) 98.76 | (3) 100.12 | (4) 15.449 |
| (5) 2.2360 | (6) 0.0123456 | (7) 9876 | (8) 9.999×10^4 |

Remark: Programmers use $2.99\text{e}8$ (or $2.99\text{E}8$) to mean 2.99×10^8 , or $1.6\text{e}-12$ (or $1.6\text{E}-12$) to mean 1.6×10^{-12} . However, we **should not** use them in handwriting.

Problems

 2.4 Write the following numbers in scientific notation. However, some of them are ambiguous, so answer “ambiguous” if so.

152 340 1000 9999 43210 0.0300 0.00213 31.0 31.00 31

 2.5 The expressions 0.123, 1.23, and 123 are numbers with **three** significant figures. Similarly, 1234 and 1.234×10^{-3} are numbers with **four** significant figures. How about the following expressions?

- | | | | |
|------------|--------------|--------------------------|----------------------------|
| (1) 3.14 | (2) 0.11 | (3) 1.1×10^{-2} | (4) 1.1×10 |
| (5) 1.0008 | (6) 1.0020 | (7) 1.0000 | (8) 4.00×10^{-10} |
| (9) 0.0008 | (10) 0.00080 | (11) 12345 | (12) 67890 |

2.4 Calculation with Significant figures

We need to do calculations of numbers with uncertainties, such as $1.23 + 4.56$ or $1.23/4.56$, but how? Here we discuss a simple method for

- addition ($x + y$) and subtraction ($x - y$)
- multiplication ($x \times y$) and division ($x \div y$ or x/y)

Other calculations, such as $\sqrt{x}$, e^x or $\sin(x)$, need the professional method given in  **TODO**  chap.

Multiplication and Division

As a rule, if x has m significant figures and y has n significant figures, then $x \times m$ or $x \div y$ should be rounded to have $\min(m, n)$ significant figures. ...But the rule is tough to understand. Probably you should see the example, and learn through practice.

Example 2.2

Calculate the following expressions, using calculators.

1. 8.912×2.4

2. $5.0 \div 3.14$

3. 2.01×49.8

4. 1.210^3

Solution

Here we use a shorthand notation “3-SF” to mean “three significant figures”.

(1) Since 8.912 has 4-SF and 2.4 has 2-SF, we round the result to (the smaller) 2-SF:

$$8.912 \times 2.4 = 21.3888 \rightarrow \underline{21}.$$

(2) Since 5.0 has 2-SF and 3.14 has 3-SF, we round the result to 2-SF:

$$5.0 \div 3.14 = 1.592... \rightarrow \underline{1.6}.$$

(3) Both 2.01 and 49.8 have 3-SF, so we keep 3-SF, but don't forget to avoid ambiguity!

$$2.01 \times 49.8 = 100.098 \rightarrow 100 \rightarrow \underline{1.00 \times 10^2}.$$

(4) $1.210^3 = 1.210 \times 1.210 \times 1.210$, so we round the result to 4-SF:

$$1.210^3 = 1.771561 \rightarrow \underline{1.772}.$$

If you want, we can justify this treatment by carrying out a hand calculation of the long multiplication.

$$\begin{array}{r} 1.20 \\ \times 1.31 \\ \hline 120 \\ 360 \\ 120 \\ \hline 15720 \end{array}$$

$$\begin{array}{r} 8.912 \\ \times 2.4 \\ \hline 35648 \\ 17824 \\ \hline 213888 \end{array}$$

$$\begin{array}{r} 2.01 \\ \times 49.8 \\ \hline 1608 \\ 1809 \\ 804 \\ \hline 100.098 \end{array}$$

Observe which digits are uncertain, and how they “pollute” the results in each step. In 1.20×1.31 , the last **1** pollutes the part of **1** + 6 and we get an uncertain number **7**. So, the result is **1.57**.

Addition and subtraction

Different rules are applied for addition and subtraction. You can understand the rules easily if you do long addition and observe which digits are polluted.

Example 2.3

Calculate the following expressions.

1. $12.3 + 4.56$

2. $123 + 4.56$

3. $0.50 - 0.032$

4. $1.20 \times 10^3 - 27$

Solution

$$\begin{array}{r} 12.3 \\ + 4.56 \\ \hline 16.86 \end{array}$$

$$\begin{array}{r} 123 \\ + 4.56 \\ \hline 127.56 \end{array}$$

$$\begin{array}{r} 0.50 \\ - 0.032 \\ \hline 0.468 \end{array}$$

$$\begin{array}{r} 1200 \\ - 27 \\ \hline 1173 \end{array}$$

and we round the results. So, the answers are **16.9**, **128**, **0.47**, and **1.17×10^3** .

Problems

 **2.6** Calculate the following, taking care of significant figures. You may use calculators.

- | | | | |
|--|---|------------------------|-----------------------------|
| (1) 1.23×4.5 | (2) 1.50×2.0 | (3) 2.00×5.00 | (4) 4.0×2.5 |
| (5) $1.23 + 4.5$ | (6) $1.50 + 2.0$ | (7) $2.00 + 8.00$ | (8) $100.0 + 0.123$ |
| (9) $1.23 \div 4.5$ | (10) $6.0 \div 3.00$ | (11) $9.00 \div 4.5$ | (12) $1.50 \div 3.0$ |
| (13) $1.23 - 4.5$ | (14) $3.10 - 0.10$ | (15) $3.10 - 0.1$ | (16) $3.10 - 3.0$ |
| (17) $131 + 69$ | (18) $131 + 6.9$ | (19) $131 + 0.69$ | (20) $1.3 \times 10^2 + 69$ |
| (21) $(1.2 \times 10^5) \times 9.4$ | (22) $1.2 \div (5.2 \times 10^5)$ | | |
| (23) $(3.33 \times 10^5) \times (6.3 \times 10^4)$ | (24) $(3.33 \times 10^5) \times (6.3 \times 10^{-4})$ | | |
| (25) $(3.33 \times 10^5) \div (6.3 \times 10^4)$ | (26) $(3.33 \times 10^5) \div (6.3 \times 10^{-4})$ | | |
| (27) $1.00 \times 10^4 + 2.5 \times 10^3$ | (28) $3.00 \times 10^3 - 4.50 \times 10^2$ | | |
| (29) $2.99 \times 10^2 + 0.814$ | (30) $1.23 \times 10^{-4} + 4.5 \times 10^{-6}$ | | |

****2.7** Calculate the following, taking care of significant figures. You may use calculators.

- | | | | |
|-------------------------------------|--------------------------------|---|-----------------|
| (1) $1.50 \times 2.000 \times 3.50$ | (2) $8.00 \div 4.00 \div 0.25$ | (3) 5.0^3 | (4) 2.10^3 |
| (5) $1.0 + 2.0 + 3.0 \times 4.0$ | (6) $17 - 8.0 - 3.0$ | (7) $3.0^3 + 1.2$ | (8) $5.0^3 - 5$ |
| (9) $(1.2 + 3.45) \times 2.1$ | (10) $(8.0 - 1.25) \div 2.0$ | (11) $1.2 \times 3.4 + 5.6$ | |
| (12) $18.0 - 2.5 \times 1.2$ | (13) $1.20 \times 3.40 + 5.60$ | (14) $(5.0 \times 10^2 + 3.4) \times 1.2$ | |

 **2.8** Calculate the following with calculators, taking care of significant figures and units.

- | | |
|---|--|
| (1) $3.1 \text{ m/s} \times 1.0 \text{ hour}$ | (2) $3.1 \text{ m} \div 25 \text{ cm}$ |
| (3) $1.5 \text{ kg} \times 9.8 \text{ m/s}^2$ | (4) $1.50 \text{ km} + 195 \text{ m}$ |
| (5) $(5.2 \text{ kg/m}^3) \times (4.0 \text{ }\mu\text{m})^3$ | (6) $(5.2 \text{ kg/m}^3) \times (4.0 \text{ }\mu\text{m}^3)$ |
| (7) $1.2 \text{ }\mu\text{m} \div 2.00 \text{ nm}$ | (8) $(1.7 \times 10^{-27} \text{ kg}) \times (3.00 \times 10^8 \text{ m/s})^2$ |
| (9) $2.000 \text{ kg} \times (3.0 \text{ m/s}^2)$ | (10) $2.000 \text{ kg} \times (3.0 \text{ m/s})^2$ |
| (11) $(1.60 \times 10^{-19} \text{ C}) \div 2.0 \text{ }\mu\text{s}$ | (12) $(1.60 \times 10^{-19} \text{ C}) \times 5.0 \text{ kV}$ |
| (13) $(6.63 \times 10^{-34} \text{ J} \cdot \text{s}) \times (3.0 \times 10^{12} \text{ Hz})$ | (14) $1.50 \text{ MJ} \div 20.0 \text{ m}$ |

Advanced note: Uncertainties can often be determined subjectively, but central values are also without specific definitions; it is often the averaged value of the measurements, but one may assume some probabilistic distribution and use its median, mean, or mode as the central value.

Chapter 3

Logic

Solve [Quiz 2.3](#) (on page 18) again.

- (1) If $mr\omega^2 = 10.0 \text{ kg}\cdot\text{m/s}^2$, $m = 5.0 \text{ kg}$, and $r = 1.0 \text{ m}$, what is ω ?
- (2) If $v_0 t + \frac{1}{2}at^2 = 5.0 \text{ m}$, where $a = -4.0 \text{ m/s}^2$ and $v_0 = 7.0 \text{ m/s}$, what is t ?
- (3) At time $t = 0$, a particle is at $(x, y) = (2.0 \text{ m}, 0)$. It moves with a constant velocity $(v_x, v_y) = (3.0, 4.0) \text{ m/s}$. What is its position at $t = 1.0 \text{ s}$?

The answers are

- (1) $\omega = \pm 1.4 \text{ s}^{-1}$ (2) $t = 1.0 \text{ s}, 2.5 \text{ s}$ (3) $x = 5.0 \text{ m}, y = 4.0 \text{ m}$

but **what do they mean?**

Quiz

Q1. Guess the correct meaning of each of above expressions.

- | | | |
|--|------|---|
| (1) ω is both $+1.4 \text{ s}^{-1}$ and -1.4 s^{-1} | v.s. | ω is either $+1.4 \text{ s}^{-1}$ or -1.4 s^{-1} |
| (2) t is both 1.0 s and 2.5 s | v.s. | t is either 1.0 s or 2.5 s |
| (3) x is 5.0 m and y is 4.0 m | v.s. | Either x is 5.0 m , or y is 4.0 m . |

Elementary educations do not discuss these differences because kids do not know **logical thinking**. Now, you are a grown-up university student. You need to think about it.

3.1 Basic Logics

Since elementary school, you have written many equalities and inequalities, such as

$$3 + 5 = 8, \quad 6 + 2 = 10, \quad 1 + 3 > -3, \quad 1 + 3 \neq 0, \quad 3^3 = 9, \quad 1 > 2, \quad \text{and} \quad \sin(\pi) \neq 0.$$

Quiz

Q2. The above statements are either **true** (T) or **false** (F). For each of them, state whether it is true or false.

Q3. Confirm that the following statements are all true:

- | | |
|-------------------------------|-------------------------------|
| (1) $3 + 5 = 8$ is true. | (2) $6 + 2 = 10$ is false. |
| (3) $3 \neq 3$ is false. | (4) False is false. |
| (5) “False is false” is true. | (6) “False is true” is false. |

We can only write true things, such as “ $3 + 5 = 8$ ”, “ $3^2 = 9$ is true”, or “ $6 + 2 = 10$ is false”. In other words, **when you write something, you must confirm that it is true**. Sometimes, you are asked to confirm a statement, or **prove** a statement. The process is called **proof** of the statement.

Numbers can be manipulated by operators such as $+$, $\div$. Similarly, true (T) and false (F) can be manipulated by **and**, **or**, and **not** operators.

and (conjunction) “A and B” ($A \wedge B$) is true if both A and B are true, and false otherwise.

or (disjunction) “A or B” ($A \vee B$) is true if at least one of A and B is true. False if both are false.

not (negation) “not A” ($\neg A$) is true if A is false, and false if A is true.

We can write these property as equations:

$$(T \wedge T) = T, \quad (T \wedge F) = (F \wedge T) = (F \wedge F) = F,$$

$$(F \vee F) = F, \quad (T \vee T) = (T \vee F) = (F \vee T) = T,$$

$$(\neg T) = F, \quad (\neg F) = T,$$

A	B	A and B	A or B	not A
T	T	T	T	F
T	F	F	T	F
F	T	F	T	T
F	F	F	F	T

but the **truth table**, shown to the right, is more useful.

Remark: Note that “ $1 + 1 = 2$ or $2 + 2 = 4$ ” is true, which says the word “or” is a bit different from everyday English. In daily conversation, “*Sho will eat ramen or sushi tonight*” usually means “but not both”. However, in mathematics, if Sho eats both ramen and sushi for a dinner, the statement “*Sho had sushi or ramen tonight*” is true.

Quiz

Q4. State whether it is true or false for the following statements.

(1) $1 + 2 = 3$ and $3 + 4 < 5$.

(2) $10 \div 3 > 0$ or $10 \div 2 > 0$.

(3) $\neg(3 > 5)$.

(4) $(6 + 3 > 0) \wedge \neg(5 - 2 > 0)$.

Fail safe note: Don't be confused by daily English.

“ $x = 1$ and $x = -1$ are the two solutions of $x^2 = 1$.” (correct daily English)

“ $x^2 = 0$ if and only if $x = 1$ or $x = -1$.” (correct mathematical English)

are both correct, but if you say “ $x = 1$ and $x = -1$ ”, then it means an impossible equality $x = 1 = -1$.

3.2 Implication $\implies$

We can discuss the following statements: are they true or false?

Example 3.1

- If $x > 5$, then $x > 1$ This is true. (Number larger than 5 are larger than 1.)
- If $x < 9$, then $x < 2$ This is false, because we have a counterexample $x = 5$.
- If $x^2 < 1$, then $x < 1$ This is true.
- If $x^2 = 1$, then $x = -1$ This is false, because we have a counterexample $x = 1$.

In general, it is easy to claim that a statement is false, i.e., to *refute* or *disprove* a statement. You just have to find one **counterexample**. Meanwhile, it is more difficult to claim that a statement is true. As mentioned above, you need to write a proof.

Now, let's try to find some counterexamples.

Quiz

Q5. The following statements are false. Find a counter example for each.

- (1) If $x^2 > 0$, then $x > 0$. (2) If $x^2 - x = 0$, then $x = 0$.
 (3) If $x^2 = y^2$, then $x = y$. (4) If $x + y < 1$, then $x < 1$ and $y < 1$.

We use the symbol “ $\Rightarrow$ ” to express “if ... then ...” statement.

$A \Rightarrow B$ means “If A is true, then B is true.” or equivalently, “ A implies B .”

and this is called **implication**. For example, the statements in the above quiz can be written as

$$x^2 > 0 \Rightarrow x > 0, \quad x^2 - x = 0 \Rightarrow x = 0, \quad x^2 = y^2 \Rightarrow x = y, \quad x + y < 1 \Rightarrow x < 1 \wedge y < 1.$$

Quiz

Q6. Similarly rewrite the statements in Example 3.1 with using the symbol “ $\Rightarrow$ ”.

Advanced note: In formal logic, $A \Rightarrow B$ is defined by $(\neg A) \vee B$, which means “if A is true, B must also be true; but if A is false, we do not care about B ”. So, if A is always false, $A \Rightarrow B$ is always true, no matter what B is. For example, $(x \in \mathbb{R} \wedge x^2 < -1) \Rightarrow x = 999$ is a true statement. Through this definition, it is clear that $A \Rightarrow B$ is equivalent to $\neg B \Rightarrow \neg A$, which we will see below.

3.3 Equivalent Statements—What does “solve” mean?

Since elementary school, you have solved many equations, but what does “solving an equation” mean?

Quiz

Q7. (1) Choose the true statements.

- (a) $2x - 1 = 0 \Rightarrow x = 1/2$. (b) $x^2 = 4 \Rightarrow x = 2$.
 (c) $2x - 1 = 0 \Rightarrow x > -10$. (d) $x^2 = 4 \Rightarrow x = 0$.

(2) Do you think these statements can be considered as “solving an equation”?

(b) and (d) are false statements (find counterexamples!), so we cannot write them. (a) and (c) are true statements, and (a) looks “solving an equation”, but we do not consider (c) is “solving an equation”. The difference between (a) and (c) is that

$$2x - 1 = 0 \Leftarrow x = 1/2 \text{ is true, but } 2x - 1 = 0 \Leftarrow x > -10 \text{ is false.}$$

So, for (a), both $\Rightarrow$ and $\Leftarrow$ are correct: we use $\Leftrightarrow$ to express such cases.

Definition 3.2 (Equivalent)

If both $A \Rightarrow B$ and $A \Leftarrow B$ are true, we call “ A and B are **equivalent**”, and write $A \Leftrightarrow B$.

$A \Rightarrow B$ is read by “ B if A ” in English. Similarly, $A \Leftrightarrow B$ is read by “ B if and only if A ” or “ B **iff** A ”. (namely, **if-and-only-if**).

Now you need to do a bit of practice. It is actually tough, unfortunately, but this drill is important for logical thinking.

Problems

 **3.1** Fill in the blanks with $\Rightarrow$, $\Leftarrow$, or $\Leftrightarrow$, where x is a real number.

- | | | |
|---|--|--|
| (1) $x = 2$ <input type="text"/> $x > 0$ | (2) $x > 1$ <input type="text"/> $x > 0$ | (3) $x = 1$ <input type="text"/> $x > 0$ |
| (4) $x = -3$ <input type="text"/> $x^2 = 9$ | (5) $x = 0$ <input type="text"/> $x^2 = 0$ | (6) $x < 0$ <input type="text"/> $x^2 > 0$ |
| (7) $x = 3$ <input type="text"/> $x \neq 5$ | (8) $3x = 3$ <input type="text"/> $x = 1$ | (9) $x^3 = x^2$ <input type="text"/> $x = 1$ |

If the statement “ $A \Rightarrow B$ ” is true, then

- B is called a **necessary condition** for A , because B is necessary for A ; if not B , then not A .
- A is called a **sufficient condition** for B , because if A is true, B is “sufficiently” true.

So, if A and B are equivalent, B (resp. A) is called *necessary-and-sufficient condition* for A (resp. B). For example, because $x = 1 \Rightarrow x^2 = 1$,

- $x = 1$ is sufficient to satisfy $x^2 = 1$. ($x = 1$ is a sufficient condition for $x^2 = 1$.)
- $x^2 = 1$ is necessary to satisfy $x = 1$. ($x^2 = 1$ is a necessary condition for $x = 1$.)

Quiz

Q8. Fill in the blanks with “a sufficient condition” or “a necessary condition”.

- | | |
|--|---|
| (1) $x = -1$ is <input type="text"/> for $x^2 = 1$. | (2) $x^2 = 4$ is <input type="text"/> for $x = 2$. |
| (3) $x > 1$ is <input type="text"/> for $x > 0$. | (4) $x^2 > 0$ is <input type="text"/> for $x < 0$. |
| (5) $x < 0$ is <input type="text"/> for $x^2 > 0$. | (6) $x^2 > 0$ is <input type="text"/> for $x > 0$. |



“To solve an equation” means “to find an *equivalent* equation in the form of $x = \boxed{}$ ”. Since $2x - 1 = 0 \Leftrightarrow x = 1/2$ and $x^2 = 4 \Leftrightarrow x = \pm 2$, we say $x = 1/2$ and $x = \pm 2$ are the solutions of the equations, respectively.

When you solve an equation, you have to check that the solution is **necessary and sufficient**.

- If you don’t check necessity, you may have an “incomplete solution”.
- If you don’t check sufficiency, you may have an “extraneous solution”.

Quiz

- Q9.** (1) Solve $\sqrt{x+2} = x$. You might reach an *extraneous solution*, which you need to *exclude* it.
- (2) Solve $\sqrt{x^2} = 4$. You might reach an *incomplete solution*, where you need to find more solutions. (Hint: $\sqrt{x^2} = x$ is a false statement.)

Before discussing advanced topics on logics, you should do some drills.

Problems

 **3.2** State whether each statement is true (T) or false (F). For example, $2 > 5$ is “F”.

- | | | |
|----------------------------------|------------------------------------|--------------------------------------|
| (1) $(3 > 1) \wedge (2 < 5)$ | (2) $3 < 1 \vee 2 > 5$ | (3) $\neg(3 > 1)$ |
| (4) $\neg(2 > 5)$ | (5) $(\neg(3 < 1)) \wedge (2 < 5)$ | (6) $\neg(3 < 1 \wedge 2 < 5)$ |
| (7) $(4 = 4) \wedge \neg(2 > 5)$ | (8) $(3 < 1) \vee \neg(3 = 1)$ | (9) $(\neg(3 < 1)) \vee \neg(3 > 1)$ |

 **3.3** State whether each statement is true or false. If false, give a counter example.

- | | |
|--|--|
| (1) If $x = 2$, then $x^2 = 4$. | (2) If $x^2 = 4$, then $x = 2$. |
| (3) $x^2 = 0 \implies x = 0$ | (4) $x > 2 \implies x > 0$ |
| (5) $x > 0 \implies x > 2$ | (6) $x = 3 \implies x = 3$ |
| (7) $(x = 1) \vee (x = -1) \implies x^2 = 1$ | (8) $x^2 = 1 \implies (x = 1) \vee (x = -1)$ |

★3.4 Solve each equation, assuming x is a real number.

- | | | | |
|----------------------|----------------------|-----------------|------------------|
| (1) $\sqrt{x} = 3$ | (2) $\sqrt{x-1} = 2$ | (3) $ x = 5$ | (4) $ x = -3 $ |
| (5) $\sqrt{x^2} = 3$ | (6) $ x^2 = 5$ | (7) $ x-2 = 3$ | (8) $ 2x+1 = 7$ |

****3.5** Solve each equation, assuming x is a real number.

- | | | |
|----------------------------|----------------------------|----------------------------|
| (1) $\sqrt{x-3} = 2$ | (2) $\sqrt{x-3} = x-3$ | (3) $\sqrt{x-3} = 3-x$ |
| (4) $\sqrt{(x-3)^2} = 2$ | (5) $\sqrt{(x-3)^2} = x-1$ | (6) $\sqrt{(x-3)^2} = x-3$ |
| (7) $\sqrt{(x-3)^2} = 3-x$ | (8) $\sin x = 1/2$ | (9) $\tan x = 0$ |

★3.6 Let x and y be real numbers. Fill in the blanks with $\implies$, $\impliedby$, or $\iff$. If your answer is $\implies$ or $\impliedby$, explain the reason by giving relevant counter examples.

- | | |
|---|--|
| (1) $x = 2$ ☐ $x^2 = 4$ | (2) $x^2 = 4$ ☐ $x = 2 \vee x = -2$ |
| (3) $x > 0$ ☐ $x^2 > 0$ | (4) $ x = 2$ ☐ $(x = 2 \vee x = -2)$ |
| (5) $x + 1 = 0$ ☐ $x = -1$ | (6) $x^2 = x$ ☐ $x(x-1) = 0$ ☐ $x = 0$ |
| (7) $\sqrt{x} = 123$ ☐ $x = 123^2$ | (8) $\sqrt{x^2} = 0$ ☐ $x = 0$ |
| (9) $\sqrt{x^2} = 3$ ☐ $x = 3$ | (10) $\sqrt{x^2} = 3 \wedge x > 0$ ☐ $x = 3$ |
| (11) $\sqrt{x^2} = \sqrt{y^2}$ ☐ $x = y$ | (12) $x + y > 0$ ☐ $x > 0 \wedge y > 0$ |
| (13) $xy = 0$ ☐ $x = y = 0$ | (14) $x^2 + y^2 = 0$ ☐ $x = y = 0$ |

****3.7** Let a be a real constant and x be a real number. For (1)–(4), fill in the blanks with $\implies$, $\impliedby$, or $\iff$. If your answer is $\implies$ or $\impliedby$, explain the reason by counter examples.

(1) $\sqrt{x} = 4$ $x = 16$

(2) $\sqrt{x^2} = 4$ $x = 4$

(3) $\sqrt{x} = a^2$ $x = a^4$

(4) $\sqrt{x^2} = a^2$ $x = a^2$

(5) Solve the equation $\sqrt{x^2} = a$, noting that a can be negative.

★3.8 The following “solutions” have logical errors. Identify errors and find the right answer.

(1) A student solves $x^2 = 9$ and writes $x = 3$.

(2) He solves $x^2 - 3x = 0$ by dividing both sides by x , obtaining $x - 3 = 0$, so $x = 3$.

(3) He solves $x^6 = x^4$ by dividing both sides by x^4 , obtaining $x^2 = 1$, so $x = \pm 1$.

Next, fill in the blanks with $\implies$, $\impliedby$, or $\iff$.

(4) $x^2 = 9$ $x = 3$ (5) $x^2 - 3x = 0$ $x - 3 = 0$ $x = 3$

(6) $x^6 = x^4$ $x^2 = 1$ $x = \pm 1$

****3.9** (1) Recall that $A \Rightarrow B$ is defined by $(\neg A) \vee B$. Also, recall that $A \Leftrightarrow B$ is defined by $A \Rightarrow B \wedge A \Leftarrow B$. Write a truth table (see page 25) for $A \Rightarrow B$, $A \Leftarrow B$, and $A \Leftrightarrow B$.

(2) Explain the reason we can understand $A \Leftrightarrow B$ as $A = B$.

(3) Write a truth table for the following expressions:

$$A \wedge B, \quad \neg(A \wedge B), \quad (\neg A) \vee (\neg B), \quad \neg(A \vee B), \quad (\neg A) \wedge (\neg B)$$

Explain that this truth table is considered as a *proof* of **de Morgan's theorem**

$$\neg(A \vee B) = (\neg A) \wedge (\neg B), \quad \neg(A \wedge B) = (\neg A) \vee (\neg B). \quad (3.1)$$

(4) Prove the following, which we will discuss in the next section.

$$(A \Rightarrow B) = ((\neg B) \Rightarrow (\neg A)), \quad (A \Rightarrow B) = \neg(A \wedge \neg B). \quad (3.2)$$

****3.10** This lecture does not cover **quantifiers** “for-all $\forall$ ” and “exists $\exists$ ”. Learn them by yourselves. In particular, prove the following:

(1) $\exists x, P(x) \iff \neg(\forall x, \neg P(x))$

(2) $\neg(\exists x, P(x)) \iff \forall x, \neg P(x)$

(3) $\forall x, P(x) \iff \neg(\exists x, \neg P(x))$

(4) $\neg(\forall x, P(x)) \iff \exists x, \neg P(x)$

****3.11** For each statement, write its negation, such as “not ($x \geq 0$ for all real x)”, in a natural form. Then, state whether the original statement and the negated one are true or false.

(1) $x^2 \geq 0$ for all real x .

(2) There exists a real x such that $x^2 = -1$.

(3) For all real x , $x > 0 \Rightarrow x^2 > 0$.

(4) $|x| = x$ for all real x .

3.4 A few more notes about logic and proof

There is a useful theorem for “not” operator, called **de Morgan’s theorem**.

Theorem 3.3 (De Morgan's theorem)

1. “not (A and B)” is equivalent to “(not A) or (not B)”, i.e., $\neg(A \wedge B) = (\neg A) \vee (\neg B)$.
2. “not (A or B)” is equivalent to “(not A) and (not B)”, i.e., $\neg(A \vee B) = (\neg A) \wedge (\neg B)$.

The proof is given as [Problem 3.9](#).

Quiz

Q10. Simplify the following statements by using de Morgan’s theorem.

- | | |
|--------------------------------------|-------------------------------|
| (1) not ((not A) or (not B)) | (2) not ((not A) and (not B)) |
| (3) $\neg((\neg A) \wedge (\neg B))$ | (4) $\neg(A \vee \neg B)$ |

Q11. For each of the following, write its negation (“not A”).

- | | | |
|--------------------------|---------------------------------|---------------------------------|
| (1) $x > 0$ or $x < 1$. | (2) $x \neq 4$ and $x \neq 5$. | (3) $0 < x < 1$. |
| (4) $x > 0$ or $y > 0$. | (5) $x = 0$ and $y = 0$. | (6) $x \neq 0$ and $y \neq 0$. |

In general, $A \Rightarrow B$ and $B \Rightarrow A$ are different. However, if $A \Rightarrow B$, then $\neg B \Rightarrow \neg A$ (not $B \Rightarrow$ not A) is always correct. It is called the **contrapositive** of $A \Rightarrow B$. Let’s see an example.

Example 3.4

If $0 < x < 2$, then x^2 is always less than 4. We can write it by $(0 < x < 2) \Rightarrow (x^2 < 4)$. Let A be the statement “ $0 < x < 2$ ” and B be the statement “ $x^2 < 4$ ”. “Not A ” is “ $x \leq 0$ or $x \geq 2$ ”. Meanwhile, “not B ” is “ $x^2 \geq 4$ ”.

- Its **converse** ($A \Leftarrow B$) is $(x^2 < 4) \Rightarrow (0 < x < 2)$, which is false (counterexample: $x = -1$).
- Its **inverse** (not $A \Rightarrow$ not B) is $(x \leq 0 \text{ or } x \geq 2) \Rightarrow (x^2 \geq 4)$, which is false.
- Its **contrapositive** (not $A \Leftarrow$ not B) is $(x^2 \geq 4) \Rightarrow (x \leq 0 \text{ or } x \geq 2)$, which is true.

Because the contrapositive of a true statement is always true, we can prove a statement by proving its contrapositive. This method is called **proof by contrapositive**.

Quiz

Q12. For each of the following statements, write its converse, inverse, and contrapositive. Then state whether each of them is true or false.

- | | |
|-----------------------------------|--------------------------------------|
| (1) If $x > 3$, then $x > 0$. | (2) If $x = 2$, then $x^2 = 4$. |
| (3) If $ x = 0$, then $x = 0$. | (4) If $x^2 = 4$, then $x \neq 3$. |

Q13. Prove the following statements by *proof by contrapositive*. Namely, first write the contrapositive, then prove it.

- (1) If $x^2 > 1$, then $|x| > 1$. (2) If $x^3 + x^2 + x < 0$, then $x < 0$.
 (3) If $xy \neq 0$, then $x \neq 0$. (4) If n^2 is not integer, then n is not integer.

Another powerful method is **proof by contradiction** (deductio ad absurdum). Imagine you prove “if A , then B ”. You assume “ A ” and “not B ” are both true, and derive a contradiction. If you reach a contradiction, it means “ A and not B ” is false, i.e., “not (A and not B)” is true, which means $A \Rightarrow B$.

Quiz

Q14. Prove the following statements by *proof by contradiction*.

- (1) If x^2 is an even integer, then x cannot be an odd integer.
 (2) If a , b , and c are integers and abc is even, at least one of a , b , or c is even.

Remark: The validity of “proof by contrapositive” and “proof by contradiction” was already discussed on page 29, Eq. (3.2).

Advanced note: You learned another useful method for proof, **mathematical induction**, in high school. In university, you may learn more advanced methods, such as **infinite descent** and **transfinite induction**.

3.5 Facts, Definitions, Assumptions, and Conclusions

One of the main reasons students find university physics difficult is that they treat all statements in the same way. In physics, every statement has its own role:

- experimental fact
- definition (in mathematics: definition)
- assumption / approximation (in mathematics: axiom or assumption)
- derived statements (in mathematics: theorem, corollary, or proposition)

You should always know which role each statement plays.

Remark: Mathematics does not have experimental facts because it is not based on experiments; it is purely based on definitions.

Example 3.5 (Types of statements in a calculation)

A particle moves as $x(t) = kt^2 + x_0$, where k and x_0 are constants. Find the averaged velocity between $t = 0$ and $t = t_0$.

Solution

We should first clarify assumptions and definitions:

- Assumptions (explicitly written): $x(t) = kt^2 + x_0$. k and x_0 are constants.
- Assumptions (implicit): k , x_0 , and t_0 are real numbers, and t means time.

- Definition: averaged velocity is *defined* by $v_{\text{avg}} := \frac{x(t_0) - x(0)}{t_0 - 0}$.

Now, we can calculate $\frac{x(t_0) - x(0)}{t_0 - 0} = \frac{(kt_0^2 + x_0) - x_0}{t_0} = kt_0$ to give a derived statement:

- Derived statement (conclusion): the averaged velocity between $t = 0$ and t_0 is $v_{\text{avg}} = kt_0$.

Quiz

Q15. Carry out a similar discussion for “find the instantaneous velocity at $t = t_1$ ”.

Consider an **experimental fact**, such as “gravity is always attractive”. We have to accept it. We need to remember it and understand what it says. “Protons have charge $+|e|$ and electrons have charge $-|e|$ ” is another experimental fact, but we can understand it as the definition of the elementary charge $|e|$.

Definitions often appear as mathematical equations, such as “instantaneous velocity is defined by $v(t) := dx/dt$ ”. Another example is “uniform circular motion is a motion with constant speed along a circle”; here, the word “uniform circular motion” is defined.

Meanwhile, **assumption** is what we can impose freely. For example, you can decide whether you ignore air resistance or not. However, every conclusion is valid only under its assumptions. If you impose bad assumptions, your conclusion will be meaningless. For example, if you want to analyze the free fall of a sheet of paper but you ignore air resistance, your result should be invalid and meaningless. You must be very careful when you make assumptions.

Remark: We often use “ $A := B$ ” to say “define A by B ”. For example, $v(t) := dx/dt$ is the definition of instantaneous velocity, and $T := 1/f$ is the definition of period. Meanwhile, “ $A = B$ ” just means “ A is equal to B ”.

Some physicists use $A \equiv B$ to say “ A is defined by B ”, but Sho does not recommend it because mathematicians use the symbol $\equiv$ for other purposes.

Problems

★3.12 Prove the following statements by contrapositive or contradiction.

(1) $x^2 + y^2 = 0 \implies x = 0$.

(2) $ab = 0 \implies a = 0$ or $b = 0$.

(3) $ab = 0 \iff a = 0$ or $b = 0$.

★3.13 With de Morgan’s theorem, rewrite the following statements in a form without “not”.

(1) not ($x > 0$ and $y > 0$)

(2) not ($x = 0$ or $y = 0$)

(3) not (if $x > 1$, then $x^2 > 1$)

(4) not ($x > 0$ iff $x^2 > 0$)

★3.14 Find the logical error: Let $a = b = 1$. Then $a^2 = ab$, so $a^2 - b^2 = ab - b^2$. So, $(a - b)(a + b) = b(a - b)$, which gives $a + b = b$, so $2 = 1$.

***3.15 For each of the following statements, write its converse, inverse, and contrapositive. State T or F for the original statement, its converse, inverse, and contrapositive.

- (1) If $x^2 - 5x + 6 = 0$, then $x = 2$ or $x = 3$.
- (2) If $x > 0$ and $y > 0$, then $x + y > 0$.
- (3) If x and y are both even integers, then xy is divisible by 4.

***3.16 A student claims: "I proved $\neg B \Rightarrow \neg A$, so I have proved $B \Rightarrow A$." Is this correct? Explain, and state what the student has actually proved.

***3.17 The following two statements look similar but are logically different. For each, state whether it is T or F and explain.

- (1) For all real x : if $x > 0$, then $x^2 > 0$.
- (2) For all real x : if $x^2 > 0$, then $x > 0$.

***3.18 True or false? Give a reason.

- (1) The contrapositive of a false statement is also false.
- (2) If $A \Rightarrow B$ is true and $B \Rightarrow A$ is false, then A and B are not equivalent.
- (3) If A is false, then $A \Rightarrow B$ is always true, for any B .

*3.19 (1) What is a **proposition**?

(2) Prove the following statements on propositions P , Q , and R .

- (a) $(P \Rightarrow Q \wedge Q \Rightarrow R) \implies P \Rightarrow R$. (**syllogism**)
- (b) $(P \Rightarrow Q) \wedge (P \Rightarrow R) \iff P \Rightarrow (Q \wedge R)$.

Chapter 4

Power, Exponential, and Logarithm

4.1 Powers

Let's begin with a review: can you recall all the rules for powers...?

Quiz

Q1. Calculate without a calculator.

(1) 2^4 , 3^5 , 4^2 , 10^3 , 1^{25} , 2^{10}

(2) 5^0 , 3^0 , 2^0 , 1^0

(3) 2^{-1} , 12^{-1} , 3^{-3} , 2^{-4} , 5^{-2} , 2^{-10}

(4) $\left(\frac{1}{2}\right)^{-1}$, $\left(\frac{1}{3}\right)^{-2}$, $\left(\frac{1}{12}\right)^{-1}$, $\left(\frac{1}{5}\right)^{-2}$, $\left(\frac{1}{2}\right)^{-10}$, $\left(\frac{1}{3}\right)^0$, $\left(\frac{1}{10}\right)^0$

(5) $\left(\frac{1}{2}\right)^4$, $\left(\frac{2}{3}\right)^3$, $\left(\frac{5}{2}\right)^2$, $\left(\frac{2}{5}\right)^{-2}$, $\left(\frac{7}{2}\right)^2$, $\left(\frac{7}{2}\right)^{-2}$, $\left(\frac{1}{3}\right)^2$, $\left(\frac{1}{3}\right)^{-2}$.

(6) $4^{1/2}$, $81^{1/2}$, $3^{1/2}$, $27^{1/3}$, $\left(\frac{1}{4}\right)^{1/2}$, $\left(\frac{1}{7}\right)^{1/2}$, $\left(\frac{8}{9}\right)^{1/2}$, $\left(\frac{15}{3}\right)^{1/2}$

(7) $4^{-1/2}$, $81^{-1/2}$, $3^{3/2}$, $27^{2/3}$, $27^{-2/3}$, $\left(\frac{1}{4}\right)^{3/2}$, $\left(\frac{1}{7}\right)^{-1/2}$, $\left(\frac{1}{2}\right)^{-5/2}$

Q2. Fill in the blanks, where $A > 0$, $B > 0$, and x, y are real numbers.

(1) $A^3 \cdot A^3 = A^{\square}$ (2) $(A^3)^3 = A^{\square}$ (3) $(A \cdot B)^4 = \square \cdot \square$

(4) $A^x \cdot A^y = A^{\square}$ (5) $(A^x)^y = A^{\square}$ (6) $(A \cdot B)^x = A^x \cdot \square$

(7) $A^{\square} = \frac{1}{A^x}$ (8) $A^x = \frac{1}{\square}$ (9) $\frac{A^x}{A^y} = A^{\square}$

(10) $A^0 = \square$ (11) $A^{\square} = A$ (12) $A^{\square} = \sqrt{A}$

(13) $A^{\square} = \sqrt[3]{A}$ (14) $\frac{A^x}{B^x} = \square$ (15) $\left(\frac{A}{B}\right)^{-1} = \square$

Q3. With $A > 0$, $B > 0$, $x \in \mathbb{R}$, and $y \in \mathbb{R}$, the following statements are all false. Find a counterexample for each.

(1) $A^x + A^y = A^{x+y}$ (2) $(A + B)^x = A^x + B^x$ (3) $(A \cdot B)^x = A \cdot B^x$

(4) $(A^x)^y = A^{(x^y)}$ (5) $(A^x)^y = A^{x+y}$

Do not worry if you cannot recall all the rules! Learn this section and come back here, and then you will be able to answer all the questions!

Let's review the definitions of **power**, a^x . Here, a is called the **base** and x is called the **exponent**.

Definition 4.1 (Power for positive base, real exponent)

For positive base $a > 0$ and real exponent $x \in \mathbb{R}$, we define the power a^x by the following steps.

First, consider a non-negative integer n . We define a^n and n -th root $\sqrt[n]{a}$, as

- $a^0 := 1$.
- If $n \in \mathbb{N}^+$, then $a^n := a \cdot a^{n-1}$.
- For $n \in \mathbb{N}^+$, we define $\sqrt[n]{a}$ as the **positive** solution X of the following equation: $X^n = a$. We write $\sqrt[2]{a}$ as $\sqrt{a}$.

If $a > 0$, positive solution always exists and is unique. So, $\sqrt[n]{a}$ is “well-defined” for $a > 0$. Then,

- If $x > 0$ and $x \in \mathbb{Q}$, we can write $x = p/q$ with $p, q \in \mathbb{N}^+$. Then we define

$$a^x = a^{p/q} := \left(\sqrt[q]{a}\right)^p = \left(a^{1/q}\right)^p$$

- If $x > 0$ and $x \notin \mathbb{Q}$, we can consider a sequence of rational numbers $x_k \in \mathbb{Q}$ such that $x_k \rightarrow x$. Then we define $a^x := \lim_{k \rightarrow \infty} a^{x_k}$.

Now we have defined a^x for all $x \geq 0$. Finally,

- If $x < 0$, we define $a^x := 1/(a^{|x|})$.

and then we have defined a^x for all $x \in \mathbb{R}$, if $a > 0$.

Example 4.2 (Integer-power of a positive number)

- $2^0 := 1$, because of the first item in the above definition.
- The second item says $2^3 := 2 \times 2^2$. Using it recursively, we get $2^2 := 2 \times 2^1$ and $2^1 = 2 \times 2^0 = 2 \times 1 = 2$. So, $2^3 = 2 \times 2 \times 2 = 8$.
- Similarly, $1.1^2 = 1.1 \times 1.1 = 1.21$ and $1.1^3 = 1.1 \times 1.21 = 1.331$.

Example 4.3 (n-th root of a positive number)

The third item says $\sqrt[3]{8}$ is the positive solution of $X^3 = 8$. So, $\sqrt[3]{8} = 2$. Similarly, $\sqrt[2]{1.21}$ is the positive solution of $X^2 = 1.21$, so $\sqrt{1.21} = 1.1$.

Example 4.4 (positive-power of a positive number)

To consider a^x for rational but non-integer x , we use the fourth item of the above definition.

- $27^{1/3} = \left(\sqrt[3]{27}\right)^1 = 3^1 = 3$. • $1.21^{0.5} = 1.21^{1/2} = \sqrt{1.21} = 1.1$.
- $27^{2/3} = \left(\sqrt[3]{27}\right)^2 = 3^2 = 9$. • $1.21^{1.5} = 1.21^{3/2} = \left(\sqrt{1.21}\right)^3 = 1.1^3 = 1.331$.

If x is not rational, we need to use the fifth item. For example, if $x = \sqrt{2} = 1.4142\dots$, we consider a sequence $1, 14/10, 141/100, 1414/1000, \dots$ which converges to $\sqrt{2}$. Then,

- $2^{\sqrt{2}}$ is the limit of $2^1, 2^{14/10}, 2^{141/100}, 2^{1414/1000}, \dots \rightarrow 2^{\sqrt{2}} = 2.665\dots$
- $3^\pi = 3^{3.14159\dots}$ is the limit of $3^3, 3^{31/10}, 3^{314/100}, 3^{3141/1000}, \dots \rightarrow 3^\pi = 31.544\dots$

Example 4.5 (negative-power of a positive number)

For a^x with $x < 0$, we use the last item in the above definition.

$$1.1^{-2} = \frac{1}{1.1^2} = \frac{1}{1.21} \quad 1.21^{-1.5} = \frac{1}{1.21^{1.5}} = \frac{1}{1.331} \quad 3^{-\pi} = \frac{1}{3^\pi} = \frac{1}{31.544\dots}$$

We have defined a^x for $a > 0$ and $x \in \mathbb{R}$. Other cases are more complicated, such as:

- a^x with $a > 0$ and $x \in \mathbb{C}$ is discussed in [♣TODO♣chapter](#) it needs a careful treatment.
- a^x with $a \leq 0$ is more complicated and requires the discussion in [♣TODO♣chapter](#). However, if x is an integer, we can safely discuss it, as follows.

Definition 4.6 (Power for negative base)

For a negative base $a < 0$, we only consider a^x for integer x .

- $a^0 := 1$.
- $a^x := a \cdot a^{x-1}$ for positive integer x .
- $a^x := 1/a^{|x|}$ for negative integer x .

We do not consider a^x if $a < 0$ and x is not an integer.

Definition 4.7 (Power for zero base)

We define $0^x := 0$ for $x > 0$, and $\sqrt[n]{0} := 0$ for $n \in \mathbb{N}^+$. (We do not consider 0^x for $x \leq 0$.)

Be careful: Notice that we have not defined $\sqrt{x}$ for $x < 0$. Sho recommends **not** to use $\sqrt[n]{x}$ for $x < 0$ because it is not useful for physics; it will just make you confused.

Quiz

Q4. Some of the following expressions are not defined. Find all the undefined ones.

$$\begin{array}{ccccccc} \pi^{-3} & \pi^{-1/\pi} & (-2)^0 & (-2)^{-3} & (-2)^{3/8} & -2^{-\pi} & \sqrt{-4} \\ \sqrt[3]{3.5} & \sqrt[3.5]{3} & \sqrt[4]{-3} & \sqrt[3]{0} & 0^{2.5} & 0^{-2.5} & 0.1^{0.1} \end{array}$$

In the following sections, we will only consider **powers of positive base**, i.e., a^x for $a > 0$.


Advanced note: Technically, even if $x < 0$, we can define $\sqrt[n]{x}$ for an odd integer n and then we can define a^x for $x \in \mathbb{Q}$ if x can be written by $x = p/q$ with an integer p and an **odd** integer q .

Advanced note: Many mathematicians define $0^0 := 1$ for some technical reasons, but we do not need it.

4.2 Exponential functions and Logarithmic functions

Powers with positive bases, such as 2^x and 0.7^x , are considered as an **exponential function** defined for $x \in \mathbb{R}$. The figure to the right shows the graphs of $y = 2^x$ and $y = 0.7^x$.

Obviously, $f(x) = a^x$ is **strictly increasing** if $a > 1$; **strictly decreasing** if $0 < a < 1$. In Section ♣TODO♣, we will check this property by calculating $f'(x)$. It is also important that a^x can take any positive real value. We will use these facts to define $\log_a x$ below in this section.



Quiz

Q5. Draw the graphs of $y = 2^x$, $y = 0.5^x$, and $y = 1^x$ by hand, without using calculators or computers.

Theorem 4.8 (Properties of Exponential Functions)

For $a > 0$, $b > 0$, and $x, y \in \mathbb{R}$,

$$a^0 = 1, \quad \frac{1}{a^x} = a^{-x}, \quad \sqrt[n]{a^x} = a^{x/n} \quad (n \in \mathbb{N}, n \geq 2); \quad (4.1)$$

$$a^x a^y = a^{x+y}, \quad \frac{a^x}{a^y} = a^{x-y}, \quad (a^x)^y = a^{xy}, \quad (ab)^x = a^x b^x, \quad \left(\frac{a}{b}\right)^x = \frac{a^x}{b^x}. \quad (4.2)$$

Remark: a^{x^y} is usually interpreted as $a^{(x^y)}$; it is in general not equal to $(a^x)^y = a^{xy}$.

Quiz

Q6. Transform the following numbers to the form of $2^{\square}$.

- (1) 2 (2) 1024 (3) $\sqrt{2}$ (4) $\sqrt[3]{4}$ (5) 0.25 (6) $\sqrt[5]{0.25}$
 (7) 4^x (8) $(\sqrt{2})^x$ (9) $4 \cdot 2^x$ (10) $2^x/16$ (11) $2^x 4^y$ (12) $2^x/4^y$

Q7. Without a calculator, arrange the following in order from smallest to largest.

$$2, \quad 1024, \quad 2^5, \quad 8^2, \quad \sqrt{2}, \quad \sqrt[3]{4}, \quad 0.25, \quad 0.5^{0.3}, \quad \sqrt[5]{0.25}, \quad 0, \quad 1$$

Consider the equation $p = a^x$, where $a > 0$, $a \neq 1$ and $p \in \mathbb{R}$. For a given p , how many solutions x does it have? Because $a \neq 1$, a^x is strictly increasing or decreasing and it takes any value $0 < a^x < +\infty$. Therefore, $p = a^x$ has a unique real solution x for any $p > 0$. We call the solution $x = \log_a p$.

Definition 4.9 (Logarithm)

For $a > 0$, $a \neq 1$, and $p > 0$, we define the **logarithm** $\log_a p$ as the unique solution x of $a^x = p$.

$$\text{For } a > 0, a \neq 1, \text{ and } p > 0, \quad a^x = p \iff x = \log_a p. \quad (4.3)$$

Notice that $\log_a p$ is not defined for $p \leq 0$ or $a \leq 0$ or $a = 1$.

Quiz

Q8. Based only on the above discussion, explain why $k = \log_a(a^k)$ and $a^{\log_a k} = k$.

We here review basic properties of $\log_a x$. Drill problems are available later in this section.

Theorem 4.10 (Properties of Logarithmic Functions)

For $a > 0$ but $a \neq 1$, $b > 0$ but $b \neq 1$, $A > 0$, and $B > 0$,

$$\log_a 1 = 0, \quad \log_a a = 1, \quad \log_a A^k = k \log_a A \quad (k \in \mathbb{R}), \quad (4.4)$$

$$\log_a(AB) = \log_a A + \log_a B, \quad \log_a\left(\frac{A}{B}\right) = \log_a A - \log_a B, \quad (4.5)$$

$$\log_a A = \frac{\log_b A}{\log_b a} \quad [\text{changing the base}]. \quad (4.6)$$

4.3 Napier's number and Natural logarithm

There is a special number for the base, called **Napier's number** e :

$$e = 2.718281828\dots = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n = \sum_{k=0}^{\infty} \frac{1}{k!} = 1 + 1 + \frac{1}{2} + \frac{1}{6} + \frac{1}{24} + \frac{1}{120} + \dots \quad (4.7)$$

It is special because of the following theorem:

Theorem 4.11 (Napier's number)

$$\frac{d}{dx} e^x = e^x, \quad \frac{d}{dx} \ln x = \frac{1}{x}, \quad (4.8)$$

where the **natural logarithm** ($\ln x$) is defined by $\ln x := \log_e x$. Also, we often write e^x as $\exp(x)$.

Quiz

Q9. Draw the graphs of $y = \exp(x)$ and $y = \ln x$, using computers or calculators.

Q10. Calculate the following, where $a > 0$, $b > 0$, and $b \neq 1$.

$$(1) \frac{d}{dx} \exp(x) \quad (2) \frac{d}{dx} e^{-3x} \quad (3) \frac{d}{dx} a^x \quad (4) \frac{d}{dx} \ln(3x) \quad (5) \frac{d}{dx} \log_b x$$

Fail safe note: Review [Quiz 4.8](#) for (3). Review Eq. (4.6) for (5). Recall e is just a number.

Problems

You may skip these drill problems, as (Sho expects) you **already have learned** them.

 **4.1** Differentiate them, where a is a positive constant such that $a \neq 1$.

- | | | | | |
|----------------------|-------------------|------------------|-------------------|------------------------|
| (1) $\exp(5x)$ | (2) e^{-x^2} | (3) $3 \exp(2x)$ | (4) $e^{\sin x}$ | (5) $x(e^x)^2$ |
| (6) $\sqrt{e^x + 1}$ | (7) $e^x e^{x^2}$ | (8) $\ln(2x)$ | (9) $\ln 3x $ | (10) $\ln(x^5 + x)$ |
| (11) $\ln \cos x $ | (12) $x \ln x$ | (13) $\ln(1/x)$ | (14) $e^x \ln x$ | (15) $\ln 2x - 1 $ |
| (16) a^{4x} | (17) a^{x^2} | (18) $(2a)^{2x}$ | (19) $\log_a x^5$ | (20) $\log_a(x^5 + x)$ |

 **4.2** Simplify each expression, where a is a positive constant such that $a \neq 1$.

- | | | | |
|-------------------------|-------------------------|------------------------|-------------------------------|
| (1) $9^{-1/2}$ | (2) $16^{3/4}$ | (3) $8^{-2/3}$ | (4) $e^{3 \ln 2}$ |
| (5) $e^{\ln 3 + \ln 4}$ | (6) $e^{\ln(6+7)}$ | (7) $\ln e^{6+7}$ | (8) $e^3(e^4)^4$ |
| (9) $\log_4 8$ | (10) $\log_{10} 0.01$ | (11) $\log_{0.5} 2$ | (12) $\log_3(1/27)$ |
| (13) $\log_6 6^{10}$ | (14) $\ln 6 + \ln(1/6)$ | (15) $2 \ln 3 - \ln 9$ | (16) $\ln 12 - \ln 4 + \ln 2$ |
| (17) 1^{-a} | (18) $\log_a 1$ | (19) $\log_a(2a^2)^4$ | (20) $\log_a(\ln(e^{5a}))$ |

 **4.3** Some of the following expressions are not defined. Find all the undefined ones.

- | | | | | | | |
|-----------------|----------------|-----------------|-----------------|-----------------|-----------------|--------------------------|
| π^0 | π^{-3} | $\pi^{2.5}$ | $\pi^{-2.5}$ | π^π | $\pi^{-\pi}$ | $\pi^{-1/\pi}$ |
| 0^0 | 0^{-3} | $0^{2.5}$ | $0^{-2.5}$ | 0^π | $0^{-\pi}$ | $0^{-1/\pi}$ |
| $(-2)^0$ | $(-2)^{-3}$ | $(-2)^{2.5}$ | $(-2)^{-2.5}$ | $(-2)^\pi$ | $-2^{-\pi}$ | $-2^{-1/\pi}$ |
| $\sqrt{4.5}$ | $\sqrt{-4}$ | $\sqrt[3]{3.5}$ | $\sqrt[3.5]{3}$ | $\sqrt[4]{3}$ | $\sqrt[4]{-3}$ | $\sqrt[4]{1 + \sqrt{2}}$ |
| $\sqrt[1.5]{1}$ | $\sqrt[4]{-8}$ | $\sqrt[3]{8}$ | $\sqrt[0]{8}$ | $\sqrt[3]{\pi}$ | $\sqrt[3]{\pi}$ | $\sqrt[4]{1 - \sqrt{2}}$ |

Problems

★4.4 Solve these equations.

- | | | |
|----------------------|--------------------------|-----------------------------|
| (1) $2^{x+1} = 64$ | (2) $e^{3x-1} = 4$ | (3) $e^x = 5$ |
| (4) $\ln x - 1 = 0$ | (5) $\ln x = 4$ | (6) $\log_5(2x + 3) = 2$ |
| (7) $2^{-2x} = 7$ | (8) $\log_3(2x + 1) = 3$ | (9) $e^{2x} - 3e^x + 2 = 0$ |

*****4.5** Draw the graphs of the following functions.

- | | | | | |
|---------------|-----------------|--------------------------------------|-----------------------|-------------------------|
| (1) $y = 2^x$ | (2) $y = 0.5^x$ | (3) $y = \left(\sqrt[3]{2}\right)^x$ | (4) $y = 2 \cdot 2^x$ | (5) $y = \frac{2^x}{2}$ |
|---------------|-----------------|--------------------------------------|-----------------------|-------------------------|

Fail safe note: Draw $y = 2^x$ first. Rewrite the other functions in the form of $y = 2^\square$.

****4.6** Solve these equations.

$$(1) \ln(x^2 - 3) = \ln(2x)$$

$$(2) \ln x + \ln(x - 1) = \ln 6$$

$$(3) \log_2 x + \log_2(x + 2) = 3$$

$$(4) \ln(x + 1) + \ln(x - 1) = \ln 3$$

$$(5) \log_2 x - \log_2(x - 2) = 3$$

$$(6) \ln(x + 1) - \ln(x - 1) = \ln 4$$

$$(7) \log_2|x| + \log_2|x + 2| = 3$$

$$(8) \ln|x + 1| + \ln|x - 1| = \ln 4$$

****4.7** Calculate the derivative of x^x .

****4.8 Hyperbolic functions** are defined by

$$\cosh x := \frac{e^x + e^{-x}}{2}, \quad \sinh x := \frac{e^x - e^{-x}}{2}, \quad \tanh x := \frac{\sinh x}{\cosh x}, \quad \text{etc.} \quad (4.9)$$

(1) Calculate the derivative of $\cosh x$, $\sinh x$, and $\tanh x$.

(2) Draw the graphs of $\cosh x$, $\sinh x$, and $\tanh x$.

(3) Show the following properties:

$$(a) \cosh^2 x - \sinh^2 x = 1.$$

$$(b) \cosh x \geq 1 \text{ for all } x.$$

$$(c) \cosh x \text{ is an even function.}$$

$$(d) \sinh x \text{ and } \tanh x \text{ are odd functions.}$$

4.4 Exponential Growth and Decay

Consider the function $f(t) = N \exp(At)$, where $N > 0$ and $A \in \mathbb{R}$.

$f(t) = N \exp(At)$ is **strictly increasing** if $A > 0$, while **strictly decreasing** if $A < 0$. (4.10)

Therefore, $f(t) = N \exp(At)$ is called **exponential growth** if $A > 0$ and **exponential decay** if $A < 0$. We learn the properties of this function in the next quiz:

Quiz

Q11. Consider the above function $f(t) = N \exp(At)$, where $N > 0$ and $A \in \mathbb{R}$.

(1) What does “strictly increasing” mean?

(2) Calculate $f'(t)$. Confirm the two statements in Eq. (4.10).

(3) Show that $f(0) = N$ and $f'(t) = Af(t)$.

Remark: We will discuss these equations in **TODO** section.

Consider an exponential decay, where the number at $t = 0$ is N . We usually express such decays by

$$\begin{aligned} f(t) &= N \exp(-\Gamma t) \quad (\Gamma > 0) \\ &= N \exp\left(-\frac{t}{\tau}\right) \quad (\text{we define } \tau := 1/\Gamma). \end{aligned} \quad (4.11)$$

Then we define the **half-life** $T_{1/2}$ as the time t such that $f(T_{1/2}) = f(0)/2$.

Quiz

- Q12.** (1) Show that $f'(t) = -\Gamma f(t)$.
(2) Show that $T_{1/2} = \tau \ln 2$.
(3) Assume $f(t_1) = 40$. Find t_2 and t_3 such that $f(t_2) = 20$ and $f(t_3) = 10$.
(4) Show that, for any t , $\frac{f(t + T_{1/2})}{f(t)} = \frac{1}{2}$.

So, each time one half-life passes, the quantity is reduced by half. If $2T_{1/2}$ has passed, the number will be 1/4 of the original number.

Problems

*****4.9** An RC circuit has charge $Q(t) = Q_0 e^{-t/\tau}$, where $\tau = RC$.

- (1) Verify that $Q(\tau) = Q_0/e$.
(2) Assume $R = 2.0 \text{ k}\Omega$ and $C = 50 \text{ }\mu\text{F}$. Calculate τ . Then, find the time t at which Q dropped to 10% of Q_0 .