

1 Spin determination at colliders

Lepton colliders

Spin measurements in electron colliders are studied in [Feng–Peskin+ \[1\]](#), [Battaglia–Datta+ \[5\]](#), [Bhattacharyya–Dey+ \[4\]](#), [Bhattacharjee–Kundu \[8\]](#), [Riemann \[7\]](#), and more.

LHC, q – l – l chain

- [Barr \[2\]](#) considers SUSY $\tilde{q} \rightarrow \tilde{\chi}_2^0 \rightarrow \tilde{l} \rightarrow \tilde{\chi}_1^0$ cascade decay to find jet–lepton angular correlation. NLO correction is found in [Horsky–Kramer+ \[22\]](#). As applications,
 - [Smillie–Webber \[6\]](#) and [Datta–Kong–Matchev \[9\]](#) to discriminate SUSY and UED by the method (the power is not very strong); the former mentions the production cross section can be used as a criterion; the latter utilizes the second KK mode production.
 - [Goto–Kawagoe–Nojiri \[3\]](#) to study L–R mixing of sleptons.
 - [Choi–Drees+ \[23\]](#) to study whether the gaugino is Majorana or Dirac.
 - [Hisano–Nojiri–Sreethawong \[28\]](#) to determine electroweakino hierarchy.
 - [Gedalia–Lee–Perez \[29\]](#), with third-generation initial squark, to discriminate SUSY/UED.
- [Alves–Eboli–Plehn \[13\]](#) provides a technique to assert \tilde{g} is Majorana fermion, based on the chain $\tilde{g} \rightarrow \tilde{b}(\tilde{b}^*) \rightarrow \tilde{\chi}_2^0 \rightarrow \tilde{l} \rightarrow \tilde{\chi}_1^0$, where we can determine whether \tilde{b} or \tilde{b}^* is produced by measuring the daughter b .
- [Wang–Yavin \[16\]](#) is for chargino/neutralino spin determination in $\tilde{q} \rightarrow qW\tilde{\chi}$ decay channel. It especially provides for $X \rightarrow Y \rightarrow Z$ -chain with $p_1 p_2 Z$ final states

$$\frac{d\Gamma}{dt_{12}} = a_0 + \cdots + a_{2s}(t_{12})^{2s} \quad \text{where} \quad t_{12} = (p_1 + p_2)^2. \quad (1)$$

This channel was studied later in [Smillie \[17\]](#).

- [Kilic–Wang–Yavin \[18\]](#) extended the work to scenarios in which primary and intermediate particles with any spins.
- A general framework for the spin determination in $D \rightarrow Cq \rightarrow Bql \rightarrow Aqll$ chain is found in [Athanasίου–Lester+ \[14\]](#) (and [Athanasίου–Lester+ \[15\]](#)) with plenty of analytic formulae. More practical analyses are found in [Burns–Kong+ \[24\]](#).

LHC, other chains

- [Barr \[11\]](#) focuses on $\tilde{l}\tilde{l}^*$ DY production (short-lived) to discriminate SUSY/UED. They propose a variable $\tanh(\eta_{l^+} - \eta_{l^-})/2$, which is boost-invariant and thus can be useful.
- [Alwall–Rainwater–Plehn \[20\]](#) considers $\tilde{W}^+ \tilde{W}^+ jj$ VBF production, mediated by Majorana neutralino, and determines the spin of \tilde{W}^+ (long-lived) by m_{jj} of the forward jets.
- [Alves–Eboli \[19\]](#) is the same analysis for $\tilde{b}\tilde{b}^*$ DY production.
- [Csaki–Heinonen–Perelstein \[21\]](#) uses m_{jj} distribution of $\tilde{g} \rightarrow jj\tilde{\chi}_1^0$ process to discriminate SUSY/UED.
- [Ehrenfeld–Freitas+ \[30\]](#) uses chains with γ , i.e., $\tilde{\chi}_2^0 \rightarrow \tilde{l} \rightarrow \tilde{\chi}_1^0 \rightarrow \tilde{G} (l\gamma)$ and $\tilde{q} \rightarrow \tilde{\chi}_1^0 \rightarrow \tilde{G} (q\gamma)$ to discriminate spins, but the power is not significant.
- [Kim \[32\]](#) discusses mass and spin measurement in $\tilde{q}_L \rightarrow qZ(\rightarrow ll)\tilde{\chi}_1^0$ chain, where intermediate $\tilde{\chi}_2^0$ can be polarized which results in the angular distribution of Z .

LHC, indirect spin measurement

- [Datta–Kane–Toharia \[10\]](#) focuses on same-sign di-lepton events and utilizes its cross section for SUSY/UED discrimination.
- [Meade–Reece \[12\]](#) discusses t' pair-production, where t' is a particle decaying into t + missing, and its mass and spin determination. The mass is determined by cross section in a two-fold way (lower for scalar, higher for fermion), and then, as an indirect spin determination, the angular distribution *in the lab frame* is utilized to distinguish the two-fold. A detector simulation is considered in [Hallenbeck–Perelstein+ \[27\]](#).
- [Kane–Petrov+ \[31\]](#) discusses crosssection-based spin measurement in colored pair production.

Others

- [Alves–Eboli+ \[25\]](#) determines the spin of a resonance V^+ in $pp \rightarrow V^+ jj$ (VBF) $\rightarrow WZjj$ process, possible in Higgsless models, where $\cos \theta_{ll}^*$ is utilized.
- [Graesser–Shelton \[26\]](#) collects possible correlations in $b\text{--}\tau$, $b\text{--}l$, and $\tau\text{--}l$, including possible mixings, but merely theoretical calculation.

References

- [1] J. L. Feng, M. E. Peskin, H. Murayama and X. R. Tata, *Testing supersymmetry at the next linear collider*, [Phys. Rev. D52 \(1995\) 1418–1432](#) [[hep-ph/9502260](#)].
- [2] A. J. Barr, *Determining the spin of supersymmetric particles at the LHC using lepton charge asymmetry*, [Phys. Lett. B596 \(2004\) 205–212](#) [[hep-ph/0405052](#)].
- [3] T. Goto, K. Kawagoe and M. M. Nojiri, *Study of the slepton non-universality at the CERN large hadron collider*, [Phys. Rev. D70 \(2004\) 075016](#) [[hep-ph/0406317](#)].
- [4] G. Bhattacharyya, P. Dey, A. Kundu and A. Raychaudhuri, *Probing universal extra dimension at the international linear collider*, [Phys. Lett. B628 \(2005\) 141–147](#) [[hep-ph/0502031](#)].
- [5] M. Battaglia, A. Datta, A. De Roeck, K. Kong and K. T. Matchev, *Contrasting supersymmetry and universal extra dimensions at the clic multi-TeV e^+e^- collider*, [JHEP 0507 \(2005\) 033](#) [[hep-ph/0502041](#)].
- [6] J. M. Smillie and B. R. Webber, *Distinguishing spins in supersymmetric and universal extra dimension models at the large hadron collider*, [JHEP 0510 \(2005\) 069](#) [[hep-ph/0507170](#)].
- [7] S. Riemann, *Z' signals from Kaluza-Klein dark matter*, eConf [C050318 \(2005\) 0303](#) [[hep-ph/0508136](#)].
- [8] B. Bhattacharjee and A. Kundu, *The International linear collider as a Kaluza-Klein factory*, [Phys. Lett. B627 \(2005\) 137–144](#) [[hep-ph/0508170](#)].
- [9] A. Datta, K. Kong and K. T. Matchev, *Discrimination of supersymmetry and universal extra dimensions at hadron colliders*, [Phys. Rev. D72 \(2005\) 096006](#) [[hep-ph/0509246](#)].
- [10] A. Datta, G. L. Kane and M. Toharia, *Is it SUSY?*, [hep-ph/0510204 \(2005\)](#).
- [11] A. J. Barr, *Measuring slepton spin at the LHC*, [JHEP 0602 \(2006\) 042](#) [[hep-ph/0511115](#)].
- [12] P. Meade and M. Reece, *Top partners at the LHC: Spin and mass measurement*, [Phys. Rev. D74 \(2006\) 015010](#) [[hep-ph/0601124](#)].

- [13] A. Alves, O. Eboli and T. Plehn, *It's a gluino*, *Phys. Rev.* **D74** (2006) 095010 [[hep-ph/0605067](#)].
- [14] C. Athanasiou, C. G. Lester, J. M. Smillie and B. R. Webber, *Distinguishing Spins in Decay Chains at the Large Hadron Collider*, *JHEP* **0608** (2006) 055 [[hep-ph/0605286](#)].
- [15] C. Athanasiou, C. G. Lester, J. M. Smillie and B. R. Webber, *Addendum to Distinguishing Spins in Decay Chains at the Large Hadron Collider*, [hep-ph/0606212](#) (2006).
- [16] L.-T. Wang and I. Yavin, *Spin measurements in cascade decays at the LHC*, *JHEP* **0704** (2007) 032 [[hep-ph/0605296](#)].
- [17] J. M. Smillie, *Spin correlations in decay chains involving W bosons*, *Eur. Phys. J.* **C51** (2007) 933–943 [[hep-ph/0609296](#)].
- [18] C. Kilic, L.-T. Wang and I. Yavin, *On the existence of angular correlations in decays with heavy matter partners*, *JHEP* **0705** (2007) 052 [[hep-ph/0703085](#)].
- [19] A. Alves and O. Eboli, *Unravelling the sbottom spin at the CERN LHC*, *Phys. Rev.* **D75** (2007) 115013 [[arXiv:0704.0254](#) (hep-ph)].
- [20] J. Alwall, D. Rainwater and T. Plehn, *Same-Sign Charginos and Majorana Neutralinos at the CERN LHC*, *Phys. Rev.* **D76** (2007) 055006 [[arXiv:0706.0536](#) (hep-ph)].
- [21] C. Csaki, J. Heinonen and M. Perelstein, *Testing gluino spin with three-body decays*, *JHEP* **0710** (2007) 107 [[arXiv:0707.0014](#) (hep-ph)].
- [22] R. Horsky, . Kramer, Michael, A. Muck and P. M. Zerwas, *Squark Cascade Decays to Charginos/Neutralinos: Gluon Radiation*, *Phys. Rev.* **D78** (2008) 035004 [[arXiv:0803.2603](#) (hep-ph)].
- [23] S. Y. Choi, M. Drees, A. Freitas and P. M. Zerwas, *Testing the Majorana Nature of Gluinos and Neutralinos*, *Phys. Rev.* **D78** (2008) 095007 [[arXiv:0808.2410](#) (hep-ph)].
- [24] M. Burns, K. Kong, K. T. Matchev and M. Park, *A General Method for Model-Independent Measurements of Particle Spins, Couplings and Mixing Angles in Cascade Decays with Missing Energy at Hadron Colliders*, *JHEP* **0810** (2008) 081 [[arXiv:0808.2472](#) (hep-ph)].
- [25] A. Alves, O. J. P. Eboli, M. C. Gonzalez-Garcia and J. K. Mizukoshi, *Deciphering the spin of new resonances in Higgsless models*, *Phys. Rev.* **D79** (2009) 035009 [[arXiv:0810.1952](#) (hep-ph)].
- [26] M. Graesser and J. Shelton, *Probing Supersymmetry With Third-Generation Cascade Decays*, *JHEP* **0906** (2009) 039 [[arXiv:0811.4445](#) (hep-ph)].
- [27] G. Hallenbeck, M. Perelstein, C. Spethmann, J. Thom and J. Vaughan, *Model Discrimination with the CMS Detector: A Case Study*, *Phys. Rev.* **D79** (2009) 075024 [[arXiv:0812.3135](#) (hep-ph)].
- [28] J. Hisano, M. M. Nojiri and W. Sreethawong, *Discriminating Electroweak-ino Parameter Ordering at the LHC and Its Impact on LFV Studies*, *JHEP* **0906** (2009) 044 [[arXiv:0812.4496](#) (hep-ph)].
- [29] O. Gedalia, S. J. Lee and G. Perez, *Spin Determination via Third Generation Cascade Decays*, *Phys. Rev.* **D80** (2009) 035012 [[arXiv:0901.4438](#) (hep-ph)].
- [30] W. Ehrenfeld, A. Freitas, A. Landwehr and D. Wyler, *Distinguishing spins in decay chains with photons at the Large Hadron Collider*, *JHEP* **0907** (2009) 056 [[arXiv:0904.1293](#) (hep-ph)].

- [31] G. L. Kane, A. A. Petrov, J. Shao and L.-T. Wang, *Initial determination of the spins of the gluino and squarks at LHC*, *J. Phys.* **G37** (2010) 045004 [[arXiv:0805.1397](#) (hep-ph)].
- [32] Y.-G. Kim, *Mass and spin measurement in squark cascade decay $\tilde{q}_L \rightarrow q\tilde{\chi}_2^0 \rightarrow qZ\tilde{\chi}_1^0 \rightarrow qll\tilde{\chi}_1^0$* , *Mod. Phys. Lett.* **A26** (2011) 1407–1411.