



Contents lists available at ScienceDirect

Cytokine

journal homepage: www.elsevier.com/locate/cytokine

Review article

Interleukin-15 in autoimmunity

Hugues Allard-Chamard^{a,b,*}, Hemant K. Mishra^c, Madhuparna Nandi^d, Marian Mayhue^d,
 Alfredo Menendez^{b,e}, Subburaj Ilangumaran^{b,d}, Sheela Ramanathan^{b,d,*}

^a Division of Rheumatology, Department of Medicine, Faculty of Medicine and Health Sciences, Université de Sherbrooke, Sherbrooke, QC, Canada

^b Centre de Recherche Clinique, Centre Hospitalier d'Université de Sherbrooke, Sherbrooke, QC, Canada

^c Vet & Biomedical Sciences, University of Minnesota, Minneapolis, MN, USA

^d Department of Immunology and Cell Biology, Faculty of Medicine and Health Sciences, Université de Sherbrooke, Sherbrooke, QC, Canada

^e Department of Microbiology and Infectious Diseases, Faculty of Medicine and Health Sciences, Université de Sherbrooke, Sherbrooke, QC, Canada

ARTICLE INFO

Keywords:

IL-15

Autoimmunity

Rheumatoid arthritis

Type 1 diabetes

Immunotherapy

ABSTRACT

Interleukin-15 (IL-15) is a member of the IL-2 family of cytokines, which use receptor complexes containing the common gamma (γ_c) chain for signaling. IL-15 plays important roles in innate and adaptive immune responses and is implicated in the pathogenesis of several immune diseases. The IL-15 receptor consists of 3 subunits namely, the ligand-binding IL-15R α chain, the β chain (also used by IL-2) and the γ_c chain. IL-15 uses a unique signaling pathway whereby IL-15 associates with IL-15R α during biosynthesis, and this complex is 'trans-presented' to responder cells that expresses the IL-2/15R $\beta\gamma_c$ receptor complex. IL-15 is subject to post-transcriptional and post-translational regulation, and evidence also suggests that IL-15 cis-signaling can occur under certain conditions.

IL-15 has been implicated in the pathology of various autoimmune diseases such as rheumatoid arthritis, autoimmune diabetes, inflammatory bowel disease, coeliac disease and psoriasis. Studies with pre-clinical models have shown the beneficial effects of targeting IL-15 signaling in autoimmunity. Unlike therapies targeting other cytokines, anti-IL-15 therapies have not yet been successful in humans. We discuss the complexities of IL-15 signaling in autoimmunity and explore potential immunotherapeutic approaches to target the IL-15 signaling pathway.