

Math 318 Homework 3

Xander Naumenko

03/02/23

Question 1a. Clearly $P(X = x) = 0 \forall x < 4$ since it takes at least 4 games for the game to end. For the remaining games, fix the number of games x and consider the games as repeated Bernoulli trials. There are two possible cases: A won or B won (where the event A is A winning and the event B is B winning). Regardless of which one, the winner must have won the last game (otherwise the game would have ended earlier). The remaining games are distributed in a binomial distribution. Therefore:

$$P(X = x \cap A) = \binom{x-1}{3} p^4 (1-p)^{x-4}.$$

$$P(X = x \cap B) = \binom{x-1}{3} p^{x-4} (1-p)^4.$$

Since these are independent events their probability can be summed, so

$$P(X = x) = \binom{x-1}{3} p^4 (1-p)^{x-4} + \binom{x-1}{3} p^{x-4} (1-p)^4.$$

Question 1b. If $X = 4$ then clearly A or B won all four games, A winning occurs with probability

$$P = \frac{p^4}{p^4 + (1-p)^4}.$$

Question 1c. To get to the point that 7 games are played it must have been tied 3-3 previously or else the game would already have been over. The probability that A wins from that point is just the probability that A wins the last game, which is just $P = p$.

Question 2a. Let y be the number of unique results. There are $\binom{6}{y}$ combinations of y numbers that are unique, while there are 6^4 possible combinations of rolls. The number of unique ways of ordering the rolls depend on what y is.

If $y = 1$ then there's only one way of arranging them, and if $y = 4$ then there's $4!$ orderings of rolls. If $y = 3$ then $\binom{4}{2} 3!$ ways of arranging the rolls, since one pair of rolls must result in the same number and there are $3!$ ways of choosing which number has the pair. Finally if $y = 2$ then either the rolls are distributed with 2 in each repeated number or there are 3 in 1 and 1 in the other. If it's 2-2 then there are $\binom{4}{2}$ ways of arranging the rolls, with 4 choices for the first pair and the remaining in the second. If it's arrange 3-1 there are $2 \cdot \binom{4}{1} = 8$ ways of this occurring. Putting this together:

$$P(Y = 1) = \frac{\binom{6}{1} 1^4}{6^4} = \frac{1}{216}.$$

$$P(Y = 2) = \frac{\binom{6}{2} (\binom{4}{2} + 2 \cdot 4)}{6^4} = \frac{35}{216}.$$

$$P(Y = 3) = \frac{\binom{6}{3} \binom{4}{2} 3!}{6^4} = \frac{5}{9}.$$

$$P(Y = 4) = \frac{\binom{6}{4} 4!}{6^4} = \frac{5}{18}.$$

The expected value is just the sum of the mass function:

$$EY = \sum_{i=1}^4 iP(Y = i) = \frac{1}{216} + \frac{70}{216} + \frac{15}{9} + \frac{20}{18} \approx 3.1.$$

Question 2b. Consider that the probability that the minimum is z is the probability that all the numbers are z or greater minus the probability that they are all $z + 1$ or greater. Thus we have:

$$P(Z = z) = 1 - \frac{(6 - z)^4}{6^4} - \left(1 - \frac{(7 - z)^4}{6^4}\right) = \frac{(7 - z)^4 - (6 - z)^4}{6^4}.$$

For the expected value, again we can just sum over the possibilities:

$$EZ = \sum_{i=1}^4 iP(Z = i) = \sum_{i=1}^4 \frac{(7 - i)^4 - (6 - i)^4}{6^4} = \frac{2275}{1296} \approx 1.76.$$

Question 3a. Summing over the possible outcomes:

$$P(Y > m) = \sum_{i=m+1}^{\infty} p(1 - p)^{i-i} = p(1 - p)^m \sum_{i=0}^{\infty} (1 - p)^i = (1 - p)^m.$$

Question 3b. Computing the limit:

$$\lim_{\delta \rightarrow 0} (1 - \lambda\delta)^{\frac{t}{\delta}}.$$

Let $\delta' = -\delta\lambda$. Then:

$$\lim_{\delta' \rightarrow 0} (1 + \delta')^{-\lambda \frac{t}{\delta'}}.$$

This is a well known identity for e , so it converges to $e^{-\lambda t}$

Question 3c. From the previous parts, we know that $P(Y > m)$ is approximately an exponential random variable. We also have:

$$P(Y > m) = \int_m^{\infty} P(Y = y) dy \implies \frac{d}{dt}(1 - e^{-\lambda t}) = P(Y = y) \implies P(Y = y) = \lambda e^{-\lambda t}.$$

Question 4. Suppose without loss of generality that a 0 is being transmitted (the same argument in reverse works in that case). The probability that the message is received correctly is the probability that the normal distribution is less than $\frac{1}{2}$. Plugging this into a calculator (since the CDF of the normal distribution isn't elementary):

$$P = P(N(0, 0.04) < 0.5) = 0.9938.$$

Question 5a. The probability mass function of the poisson distribution is $\frac{\lambda e^{-\lambda}}{k!}$. Plugging in $k = 2$ and maximizing:

$$\frac{d}{d\lambda} \left(\frac{\lambda^2 e^{-\lambda}}{2} \right) = \lambda e^{-\lambda} - \frac{1}{2} \lambda^2 e^{-\lambda} = 0 \implies \lambda = 2.$$

Question 5b. λ is the mean time between murders, and the average over each week in this case was 1.5. Thus the λ that maximizes the probability of this sequence occurring is $\lambda = 1.5$.

Question 5c. Again, λ is the mean time between murders. This time if we calculate the average time between murders we have to average over the sum of the sequence, so we get:

$$\lambda = \frac{1}{k} \sum_{i=1}^k a_k.$$

Question 6a.

```
from math import comb
```

```
p = 1/50
seats = 420
sold = 430
```

```
Prob = 0
for i in range(seats+1,sold+1):
    Prob += p**(sold-i)*(1-p)**i*comb(sold,i)
```

$$\implies P = 0.6405.$$

Question 6b.

```
from math import exp,factorial
```

```
p = 1/50
seats = 420
sold = 430
```

```
lam = p*sold
Prob = 0
for i in range(sold-seats):
    Prob += lam**i*exp(-lam)/factorial(i)
```

```
print(Prob)
```

$$\implies P = 0.63995.$$

Question 6c.

```

import matplotlib.pyplot as plt
import numpy as np
from random import randint
from scipy.stats import poisson

p = 1/50
seats = 420
sold = 430
n = 50000

no_shows = []

X = np.arange(1, 30, 1)
Y = poisson.pmf(X, mu=sold*p)

for _ in range(n):
    no_shows.append(sum(randint(1,50)==1 for _ in range(sold)))

plt.hist(no_shows, density=True, bins=24)
plt.plot(X,Y)
plt.show()

```




Figure 1: Graph for question c

Question 6d. As can be seen by figure 2, the series eventually stabilizes to approximately 0.65. This means that the airline in the long run expect around 65% of flights to be overbooked.

```

import matplotlib.pyplot as plt
import numpy as np
from random import randint
from scipy.stats import poisson

p = 1/50
seats = 420
sold = 430
n = 50000

no_shows = []

```

```

for _ in range(n):
    no_shows.append(sum(randint(1,50)==1 for _ in range(sold)))

N = np.arange(1,n+1,1)
Xn = []
tot = 0
for i in range(len(no_shows)):
    tot += no_shows[i] < sold-seats
    Xn.append(tot / (i+1))

print(Xn[-1])
plt.plot(N,Xn)
plt.show()

```




Figure 2: Graph for part d