MATH 443 Homework 4

Xander Naumenko

13/02/23

Question 1. Suppose by way of contradiction that the root of a DRT T had in-degree nonzero. Let r be the root of T and v be a vertex such that $vr \in E(G)$. Then vr is a path from v to r in the underlying tree of T. Since T is a tree it is the only path between r and v, so there doesn't exist a directed path from r to v, which is a contradiction of our assumption that T is a DRT. Therefore r has in-degree zero since no vertices can lead into it.

To see why each other vertex $v \neq r$ must have in-degree 1, first note that they clearly can't have in-degree 0 since there exists a directed path from r to v, and the last edge in this path will contribute 1 to the in-degree of that vertex. To see why the number of in vertices can't be more than one, supposed by contradiction that it was, i.e. suppose there exist $u_1, u_2 \in V(T)$ s.t. $u_1v \in E(T), u_2v \in E(T)$. Since $u_1, u_2 \in V(T)$ there exists ordered paths P_1, P_2 such that the first vertex of both is r and the last is u_1 and u_2 respectively. Let w be the last vertex of P_1 that is also in P_2 . Let Q_1, Q_2 be the sub paths of P_1, P_2 that go from w to v. $Q_1 \cap Q_2 = \emptyset$ since w was chosen to be the last shared vertex. However then wQ_1vQ_2w is a cycle which is impossible in a tree. Therefore the in-degree number can't be 0 and can't be 2 or greater, so it must be 1. \square

Question 2. Let T_1, T_2 be disjoint DRTs and let e be a directed edge with one endpoint in T_1 and the other in T_2 .

- (\Rightarrow) Assume $(T_1 \cup T_2) + e$ is a DRT, we will prove that the second vertex of e is the root of T_1 or T_2 , call them r_1, r_2 . To see why suppose by contradiction that that the second vertex of e = uv is not r_1, r_2 , and WLOG assume $u \in E(T_1), v \in E(T_2), v \neq r_2$. Let P_1 be a directed path from r_1 to u and let P_2 be a directed path from r_2 to v. Then $P_1 + e$ is a directed path from r_1 to v in the new graph. However this implies that in the new graph v has an in-degree of at least 2 (since the second last vertex of P_1 is in T_1 and the second last vertex of P_2 is in T_2). However problem 1 in this homework proved that no vertex on a DRT has in-degree 2 or greater, so it must be that e was the root of T_1 or T_2 .
- (⇐) Assume the endpoint of e is the root of T_1 or T_2 . WLOG assume $e = ur_2$ where r_2 is the root of T_2 , and let r_1 be the root of T_1 . Let $T = (T_1 \cup T_2) + e$ and let $x \in V(T)$. If $x \in T_1$ then since T_1 is a DRT there exists a directed path from r_1 to x. If $x \in T_2$ then let P_1 be a directed path from r_1 to u and u and u be a directed path from u to u. Then u and u are directed path from u to u and u are directed path from u are directed path from u and u are directed path from u and u are directed path from u and u are directed path from u are directed path from u are directed path from u and u are directed path from u ar

Question 3. The statement is true. Let P be a longest path of a connected graph G, and let u, v be its two endpoints. The statement will be shown for u, although since u, v were arbitrary it also holds for v. By way of contradiction let $x, y \in G - u$ such that x, y are disconnected in G - u.

G is connected so there exists a path $Q \subset G$ from x to y. Given that x, y became disconnected after removing u it must be that $u \in V(Q)$. $x \neq u, y \neq u$, so there exist paths Q_1 between x and u_1 and Q_2 between u_2 and y, where $u_1, u_2 \in N(u)$. We proved in class that $N(u) \in V(P)$, since otherwise you could extend P by including a vertex of N(u) not already in P. Let $P_1 \subset P$ be the portion of P between u_1, u_2 . Note that $u \notin P_1$ since u is an endpoint of P, so it's not a midpoint of any subpath. But then $xQ_1u_1P_1u_2Q_2y$ is a path of G-u between x and y which contradicts our assumption that x, y disconnected. Therefore u couldn't have been a cut vertex, and by the symmetry of the argument v couldn't have been either. \square

Question 4. The flaw is that the distance between the endpoints of a longest path of a graph G are not necessarily the farthest from each other. To see why consider C_4 . Then the longest path includes all 4 vertices, but the distance between two endpoints of such a path is just 1 whereas the actual farthest vertices are distance 2. Therefore it's not valid to assume that the endpoints of a longest path are the vertices that are the farthest from one another.

Question 5. The statement is true. Let C be a component of G. Clearly C is connected, since otherwise it wouldn't be a component. Let V be a set of vertices of size smaller than 1. The only possibility is that $|V| = 0 \implies V = \emptyset$. However $C - V = C - \emptyset = C$ is connected, so C is 1-connected. \square

Question 6. Let v be the vertex added to G and V be a minimal separating set of G'. We will consider two cases: $v \in V$ and $v \notin V$.

Case 1 ($v \in V$): Note that V - v must be a separating set for G (since $V - v \subset G$ and V separates G'). Therefore $|V - v| \ge k \implies |G| \ge k + 1 > k$ as required.

Case 2 ($v \notin V$): If v is a trivial component in G' - V, $N(v) \subset V$ (in fact N(v) = V by V's minimality but this isn't required) and $|V| \geq |N(v)| \geq k$ which is what we're trying to show. If v isn't its own component, it is part of a component $C \subset G', |C| \geq 2$ and G' - V is disconnected. Then $C - v \neq \emptyset$ so G - V = G' - V - v separates G, so $|V| \geq k$. Either way we've shown that $|V| \geq k$, which is the definition of being k-connected. \square

Question 7. Consider the number of edges that must be removed to generate three components. Let x be the number of vertices of the first component G_1 , and y be the number of vertices of the second component G_2 . Then the last vertex has 3n-x-y vertices, call it G_3 . The number of edges to disconnect G_1 is x(3n-x), since each of the x vertices is attached to 3n-x vertices. G_2 has y vertices attached to 3n-y vertices, but x of the edges attached to the second of those vertices was already removed in the first step. Therefore you must remove y(3n-x-y) vertices to separate the second component. After removing both of these the last will be disconnected, since neither of the G_1, G_2 are connected to it by design. Thus the total number of edges required to disconnect is:

$$f(x,y) = x(3n-x) + y(3n-x-y) = 3n(x+y) - (x^2 + xy + y^2).$$

Fix x and consider optimizing y. The optimum occurs either on the edges (y = 1, y = 3n - 2) or where the derivative is zero. Note however that this is a inverted parabola with its vertex at $y = \frac{-b}{2a} = \frac{3n-x}{2}$ which is its maximum point, so if we were to take the derivative and set it to zero we would find a maximum, not am minimum. Thus y = 1 or y = 3n - 2.

If y = 3n - 2, then the only option for x would be x = 1 and the total number of edges is f(1, 3n - 2) = 6n - 3. If instead y = 1, then using the same logic as we used to optimize y, x = 3n - 2 or x = 1. Since the remaining component will be the opposite of whatever choice of x we use (and switching them doesn't change f), WLOG assume x = 1 as well. Then the total number of removed edges is still f(1,1) = 6n - 3. We've covered all possible cases, so the minimum number of edges required is |X| = 6n - 3.