

PHYS 403 Homework 3

Xander Naumenko

18/03/24

Question I. Taking the derivative with respect to a specific p_k :

$$\begin{aligned} \frac{\partial}{\partial p_k} \left(S[p_n] - \alpha \left(\sum_n p_n - 1 \right) - \beta \left(\sum_n p_n - \bar{E} \right) - \gamma \left(\sum_n p_n N_n - \bar{N} \right) \right) \\ = -\log p_k - 1 - \alpha - \beta E_n - \gamma N_n = 0 \implies p_k = e^{-(1+\alpha+\beta E_n+\gamma N_n)}. \end{aligned}$$

By comparing this expression to what we derived in class ($p_k = \frac{1}{Z} e^{-\beta(E_n - \mu N_n)}$), we conclude that $e^{-1-\alpha} = \frac{1}{Z}$, $\beta = \frac{1}{k_B T}$ and $\gamma = -\frac{\mu}{k_B T}$.

Question II. Assume the balloon initially at a radius R , and consider a small change dR in the radius. Let P_i be the pressure inside the balloon and P_o to be that outside. Then we have:

$$P_o dA = P_i dA - \sigma dV \implies P_i - P_o = \sigma \frac{dA}{dV}.$$

We have that $V = \frac{4}{3}\pi R^3$, so $dV = \frac{4}{3}\pi((R + dR)^3 - R^3) = \frac{4}{3}\pi(3R^2 dR + 3R dR^2 + dR^3) \approx 4\pi R^2 dR$. Similarly, we have $dA = 4\pi((R + dR)^2 - R^2) \approx 8\pi R dR$. Plugging this into the above expression, we get $P_i - P_o = \frac{2\sigma}{R}$ as required.

Question III.1.

$$\begin{aligned} Z_{G.C.}^{(d)}(\mu, T) &= \sum_{N=0}^{\infty} \frac{1}{N!} \int \frac{d^3 r_1 \cdots d^3 r_N d^3 p_1 \cdots d^3 p_N}{(2\pi\hbar)^{dN}} e^{-\beta \left(\sum_{i=1}^N \frac{p_i^2}{2m} - (\mu - \epsilon_d)N \right)} \\ &= \sum_{N=0}^{\infty} e^{\beta(\mu - \epsilon_d)N} \frac{1}{N!} \left(\frac{R^{dN}}{\lambda^d} \right)^N = \exp \left(\frac{e^{\beta(\mu - \epsilon_d)} R^{dN}}{\lambda^d} \right). \end{aligned}$$

Question III.2. Using the derivative of the partition function:

$$\begin{aligned} \langle N \rangle &= \frac{1}{\mu} \frac{\partial}{\partial \beta} \log Z = \frac{e^{\beta\mu} R^{3N}}{\lambda^3} = \frac{e^{\beta\mu} V}{\lambda^3} \\ \implies n_V &= \frac{\langle N \rangle}{V} = \frac{e^{\beta\mu}}{\lambda^3}. \end{aligned}$$

Question III.3. Using same technique as before:

$$\begin{aligned} \langle N \rangle &= \frac{1}{\mu} \frac{\partial}{\partial \beta} \log Z = \frac{A}{\lambda^2} e^{\beta(\mu + \epsilon_0)}. \\ \implies n_S &= \frac{e^{\beta(\mu + \epsilon_0)}}{\lambda^2}. \end{aligned}$$

Question III.4. The total change to surface tension, as stated in the question of part 3, is $\Delta\sigma = -\epsilon_0 n_s$. Since n_s is strictly decreasing with temperature (since $\beta = \frac{1}{k_B T}$), this means that increasing temperature causes the surfactant to mix oil and water less effectively.