

Math 406 Homework 4

Xander Naumenko

05/11/23

Question 1a.

Question 2a. Integrating:

$$\begin{aligned}\int_0^1 v L u dx &= \int_0^1 v (x^2 u'' + 3x u' - u) = [v x^2 u' + 3v x]_0^1 - \int_0^1 (2x^2 v)' u' + (3x v)' u + u v dx \\ &= [2x^2 v u' + 3x v u + v' u]_0^1 + \int_0^1 u ((2x^2 v)'' - (3x v)' - v) dx.\end{aligned}$$

Thus the adjoint operator is $L_s^* v = (2s^2 v)'' - (3s v)' - v = 2s^2 v'' + 5s v' \implies L^* = 2s^2 \frac{d^2}{ds^2} + 5s \frac{d}{ds}$. We want to find $v(s, x)$ with $v(0, x) < \infty$ and $v(1, x) = 0$ so we can write $u(x) = v_s(1, x) + \int_0^1 v(s, x) f(s) ds$. We want to find v such that $L^* v(s, x) = \delta(x - s)$. Try $v = s^r$ in the homogeneous equation:

$$L^* v = 2s^2 v_{ss} + 5s v_s = 2r(r-1)s^r + 5r s^r = 0 \implies 2r(r-1) + 5r = r(2r+3) = 0 \implies r = 0 \text{ or } -\frac{3}{2}.$$

For non-homogeneous $L^* v = \delta(s - x)$, we can solve to the right and left of $x = s$:

$$v(s, x) = \begin{cases} A_- + B_- s^{-\frac{3}{2}} & 0 < s < x \\ A_+ + B_+ s^{-\frac{3}{2}} & x < s < 1 \end{cases}.$$

The regularity condition implies that $B_- = 0$, and the $s = 1$ condition imposes $B_+ = -A_+$. We also need continuity, so $A_- = A_+(1 - x^{-\frac{3}{2}})$. Finally, the jump condition:

$$1 = \int_{x-\epsilon}^{x+\epsilon} 2s^2 v_{ss} + 5s v_s ds = (2s^2 v)_s \Big|_{x-\epsilon}^{x+\epsilon} = 2s^2 v_s \Big|_{x-\epsilon}^{x+\epsilon} = 2x^2 \left(\frac{3}{2} A_+ x^{-\frac{5}{2}} - 0 \right) \implies A_+ = \frac{1}{3} \sqrt{x}.$$

Putting this all together, we get

$$v(s, x) = \begin{cases} \frac{1}{3} \left(x^{\frac{1}{2}} - x^{-1} \right) & 0 < s < x \\ \frac{1}{3} \sqrt{x} \left(1 - s^{-\frac{3}{2}} \right) & x < s < 1 \end{cases}.$$

Finally, plugging this into the solution form given above:

$$u(x) = 2v_s(1, x)u(1) + \int_0^1 v(s, x) f(s) ds = \int_0^x \frac{1}{3} \left(x^{\frac{1}{2}} - x^{-1} \right) f(s) ds + \frac{1}{3} \sqrt{x} \int_x^1 \left(1 - s^{-\frac{3}{2}} \right) f(s) ds.$$

The original question asks for G , but of course here $G(s, x) = v(s, x)$ since they represent the same thing.

Question 2b. From class, the factor to multiply the equation by is:

$$F = e^{\int \frac{a_1}{a_0} dx} \frac{1}{a_0} = e^{\int \frac{3}{2x} dx} \frac{1}{2x^2} = \frac{1}{2\sqrt{x}}.$$

Multiplying this, we get:

$$FLu = x^{\frac{3}{2}}u'' + \frac{3}{2}x^{\frac{1}{2}}u' - \frac{1}{2}x^{-\frac{1}{2}}u = \frac{1}{2}x^{-\frac{1}{2}}f.$$

Call this new self adjoint operator L' . Running through the same process as for part a again, we first find the boundary terms for our expression of u . We know that the new operator is self adjoint though, so we can immediately write (choosing $v(1, x) = 0$ and $v(0, x) < \infty$):

$$\int_0^1 vL'udx = \left[vx^{\frac{3}{2}}u' + \frac{3}{2}x^{\frac{1}{2}}v - \frac{3}{2}x^{\frac{1}{2}}vu - x^{\frac{3}{2}}v'u \right]_0^1 + \int_0^1 uL'vdx.$$

From the boundary terms we get that $v(1) = 0$ and $v(0) < \infty$ gives enough information for all of the terms, so the operator L' is also essentially self adjoint. To solve the homogeneous case try $v = s^r$:

$$L's^r = 0 \implies r(r-1) + \frac{3}{2}r - \frac{1}{2} = 0 \implies r = -1 \text{ or } \frac{1}{2}.$$

Applying the boundary conditions $v(0, x) < \infty$ and $v(1, x) = 0$, we can write the solution to $L'v = \delta(x - s)$ as:

$$v(s, x) = \begin{cases} A_- s^{\frac{1}{2}} & 0 < s < x \\ A_+ \left(s^{-1} - s^{\frac{1}{2}} \right) & x < s < 1 \end{cases}.$$

Continuity gives $A_- x^{\frac{1}{2}} = A_+ \left(x^{-1} - x^{\frac{1}{2}} \right) \implies A_- = A_+ \left(x^{-\frac{3}{2}} - 1 \right)$. Finally, the jump condition results in

$$\begin{aligned} s^{\frac{3}{2}}v_s|_{x=s}^{x+\epsilon} = 1 &\implies x^{\frac{3}{2}} \left(A_+ \left(-x^{-2} - \frac{1}{2}x^{-\frac{1}{2}} \right) - A_+ \left(x^{-\frac{3}{2}} - 1 \right) \frac{1}{2}x^{-\frac{1}{2}} \right) = 1. \\ &\implies A_+ = -\frac{2}{3}x^{\frac{1}{2}}. \end{aligned}$$

Thus our expression for the Green's function $v(s, x) = G(s, x)$ is

$$v(s, x) = \begin{cases} -\frac{2}{3} \left(x^{-1} - x^{\frac{1}{2}} \right) s^{\frac{1}{2}} & 0 < s < x \\ -\frac{2}{3} x^{\frac{1}{2}} \left(s^{-1} - s^{\frac{1}{2}} \right) & x < s < 1 \end{cases}.$$

Using this to solve for u :

$$u(x) = \frac{1}{3} \int_0^x \left(1 - x^{-\frac{3}{2}} \right) s^{\frac{1}{2}} f(s) ds + \frac{1}{3} \int_x^1 \left(s^{\frac{1}{2}} - s^{-1} \right) f(s) ds.$$

Question 3. Because $a'_0 = 0 = a_1$, the operator L is self adjoint. This is a special case of the form $(pu')' + qu = f$ which in class we showed can be expressed as

$$u(x) = [vu' - v'u]_0^\infty + \int_0^\infty v(s, x) f(s) ds$$

for $Lv = \delta(s, x)$. The required boundary conditions on v to make each term knowable are $v \rightarrow 0, v' \rightarrow 0$ as $x \rightarrow \infty$. First solving the homogeneous equation, we have

$$Lv = v'' + v = 0 \implies v(s, x) = A \sin(s) + B \cos(s).$$

Applying this to the non-homogeneous equation $Lv = \delta(s - x)$, we have

$$v(s, x) = \begin{cases} A_- \sin(s) + B_- \cos(s) & 0 < s < x \\ A_+ \sin(s) + B_+ \cos(s) & s > x \end{cases}.$$

The boundary conditions on v at infinity force $A_+ = B_+ = 0$. Continuity forces $A_- \sin(x) + B_- \cos(x) = 0 \implies B_- = -\tan(x)A_-$. Finally, the jump condition gives:

$$\int_{x-\epsilon}^{x+\epsilon} v_{ss} + v ds = 1 \implies v_s(x^+, x) - v_s(x^-, x) = (0 - A_- (\cos(x) + \tan(x) \sin(x))) = 1 \implies A_- = -\cos(x).$$

Thus our final expression for v is:

$$v(s, x) = \begin{cases} -\cos(x) \sin(s) + \sin(x) \cos(s) & 0 < s < x \\ 0 & s > x \end{cases}.$$

For the boundary terms seen previous, we then have $v(0, x) = -\sin(x)$ and $v_s(0, x) = \cos(x)$. Expressing u in terms of these Green's functions:

$$u(x) = [vu' - v'u]_0^\infty + \int_0^\infty v(s, x) f(s) ds = \sin(x)v_0 + \cos(x)u_0 - \int_0^x (\cos x \sin s - \sin x \cos s) f(s) ds$$

Question 4a. The question states to solve it assuming that solutions are in the form r^i , but as far as I can tell solutions aren't in this form. Instead, I claim that $G_{ij} = ki + c$:

$$k(i+1) - 2ki + k(i-1) = 0.$$

For the non-homogeneous equation, the boundary conditions and continuity enforce what constants are allowed. Thus solutions are in the form:

$$G_{ij} = \begin{cases} \frac{k}{j}i & 0 \leq i < j \\ k & i = j \\ \frac{k}{N-j}(N-i) & j < i \leq N \end{cases}.$$

The final condition is that $G_{j+1j} - 2G_{jj} + G_{j-1j} = 1$, so $1 = \frac{k}{j}(j-1) - 2k + \frac{k}{N-j}(N-j-1) \implies k = \frac{j(j-N)}{N}$. Thus the explicit solution to (3) is

$$G_{ij} = \begin{cases} \frac{j-N}{N}i & 0 \leq i < j \\ 1 & i = j \\ -\frac{j}{N}(N-i) & j < i \leq N \end{cases}.$$