

MATH318 Homework 4

Xander Naumenko

09/02/23

Question 1a. The cumulative distribution function is the probability that T is less than a certain value t :

$$P(T > t) = P\left(\frac{1}{2}X^2 > t\right) = P(X > \sqrt{2t}) = \begin{cases} 0 & \text{if } t \leq \frac{9}{2} \\ \frac{\sqrt{2t}-3}{5} & \text{if } \frac{9}{2} < t < 32 \\ 1 & \text{if } t \geq 32 \end{cases}.$$

Question 1b. The probability density function is the derivative of the CDF :

$$f(t) = \frac{d}{dt}CDF(t) = \frac{d}{dt} \left(\frac{\sqrt{2t}-3}{5} \right) = \frac{1}{5\sqrt{2t}}.$$

Question 1c. Taking the integral:

$$ET = \int_{\frac{9}{2}}^{32} \frac{t}{5\sqrt{2t}} dt = \left(\frac{\sqrt{2t}^{\frac{3}{2}}}{15} \right) \Big|_{\frac{9}{2}}^{32} = 16.17.$$

Question 2a. Clearly the first coupon will be obtained with probability 1. Once that event has happened, the probability that the next one is unique is $\frac{n-1}{n}$, which is a geometric random variable. This process repeats until all the unique coupons have been picked. The parameter of each random variable is the probability, which for the i th coupon is $p = \frac{n+1-i}{n}$. Consider $G(p)$ to be a geometric random variable with parameter p , then

$$T = \sum_{i=1}^n G\left(\frac{n+1-i}{n}\right).$$

Question 2b. Using the hint:

$$ET = \sum_{i=1}^n EG\left(\frac{n+1-i}{n}\right) = n \sum_{i=1}^n \frac{1}{n+1-i} = n \sum_{i=1}^n \frac{1}{i}.$$

Question 3a. Since T, B are independent their joint PDF is simply $f(t, b) = 6e^{-6t}4e^{-4b} = 24e^{-6t-4b}$

Question 3b. Taking an integral over the part of the plane where $t < b$:

$$p = \int_0^\infty \int_0^b 24e^{-6t-4b} dt db = \int_0^\infty -4e^{-10b} + 4e^{-4b} db = 1 - \frac{4}{10} = 0.6.$$

Question 3c. As the hint suggests let $X = \min(T, B)$. Then we have

$$P(X > y) = P(T > y)P(B > y) = e^{-6y}e^{-4y} = e^{-10y}.$$

Taking the distribution to get the PMF:

$$f(y) = \frac{d}{dy}P(X < y) = \frac{d}{dy}(1 - e^{-10y}) = 10e^{-10y}.$$

This is just an exponential distribution with parameter 10.

Question 4. X, Y are independent (since they are both just $N(0, 1)$), so the joint probability distribution is:

$$P(R) = \left(\frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}x^2} \right) \left(\frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}y^2} \right) = \frac{e^{-\frac{1}{2}(x^2+y^2)}}{2\pi}.$$

To find the cumulative distribution function of R we can integrate over the part of the plane where $X^2 + Y^2 < R^2$:

$$\begin{aligned} P(R < r) &= \int_0^r \int_0^{\sqrt{r^2-x^2}} \frac{1}{2\pi} e^{-\frac{1}{2}(x^2+y^2)} dy dx. \\ &= \frac{1}{2\pi} \int_0^{2\pi} \int_0^r e^{-\frac{1}{2}\rho^2} \rho d\rho d\theta = e^{-\frac{1}{2}r^2}. \end{aligned}$$

The probability density is the derivative of the cumulative probability function:

$$f(r) = \frac{d}{dr}P(R < r) = re^{-\frac{1}{2}r^2}.$$

This is called the Rayleigh distribution.

Question 5a. Since X, Y are uniform, the probability that R is less than r is just the ratio of the area of the circle of radius less than r to the whole circle. In formula:

$$P(R < r) = \frac{\pi r^2}{\pi 1^2} = r^2.$$

The PDF is just the derivative:

$$f(r) = \frac{d}{dr}r^2 = 2r.$$

Question 5b. Doing the integral over the possible values:

$$ER = \int_0^1 r 2r dr = \frac{2}{3} r^3 \Big|_0^1 = \frac{2}{3}.$$

Question 5c. Since the distribution is uniform, the joint probability distribution of X and Y is $f(x, y) = \frac{1}{\text{area}} = \frac{1}{\pi 1^2} = \frac{1}{\pi}$. For a fixed value of $X = x$, the probability of it occurring can be found by integrating this distribution over y :

$$f_X(x) = \int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}} \frac{1}{\pi} dx = \frac{2}{\pi} \sqrt{1-x^2}.$$

Since this argument is symmetric between X and Y , we have

$$f_Y(y) = \frac{2}{\pi} \sqrt{1-y^2}.$$

Question 5d. Taking the product of their PDFs:

$$f_X(x)f_Y(y) = \frac{4}{\pi}\sqrt{(1-x^2)(1-y^2)}.$$

Since this does not equal $f(x, y) = \frac{1}{\pi}$, X and Y aren't independent.

Question 5e. See figure 1. Code:

```
from random import uniform
import matplotlib.pyplot as plt

n = 10000

pts = [(uniform(-1,1),uniform(-1,1)) for _ in range(n)]
pts_inside = [t for t in pts if t[0]**2 + t[1]**2 < 1]
l = list(zip(*pts_inside))

plt.scatter(l[0], l[1])
plt.plot()
plt.show()
```

Question 5f. See figure 2. The distribution does not appear to be random, as there is a bunching up for this one around 0. This is because the probability distribution function for R is not uniform as we saw in part a. Code:

```
from random import uniform
import matplotlib.pyplot as plt
import math

n = 10000

rtheta = [(uniform(0,1),uniform(0, 2*math.pi)) for _ in range(n)]
pts = [(r*math.cos(t), r*math.sin(t)) for (r,t) in rtheta]
l = list(zip(*pts))

plt.scatter(l[0], l[1])
plt.plot()
plt.show()
```

Question 5g. From multivariate calculus, we have that

$$\text{area}(A) = \iint_A r dr d\theta.$$

Since this is identical to what we're trying to find except for a factor of $\frac{1}{\pi}$, we have that

$$f(r, \theta) = \frac{r}{\pi}.$$

Question 6a,b,c. The graphs can be seen in figures 3, 4 and 5. The code:




Figure 1: 5e




Figure 2: 5f

```

import numpy as np
import numpy.random as rn
import matplotlib.pyplot as plt

num = 10000

normals = rn.normal(np.zeros(num))
exps = rn.exponential(np.ones(num)/2)
cauchys = rn.standard_cauchy(num)
for dist in (normals, exps, cauchys):
    averages = []
    s = 0
    ns = np.arange(num)
    for n in ns:
        s += dist[n]
        averages.append(s/(n+1))

plt.plot(ns, averages)
plt.show()

```




Figure 3: 6a




Figure 4: 6b

Question 6d. The plot from part a converges to 0 and the plot from part b converges to 0.5. These make sense as the mean for the normal and exponential distribution are 0 and $\frac{1}{\lambda}$ respectively.




Figure 5: 6c

The plot from part c doesn't seem to converge, and each time the simulation is run the graph seems to change.