Hw 5: STAT: 4520. Yubing Li 7.1. (1). Non-info prior: p(M, 52) 00 52 (-00 < M <00) (0002000) 1 (y/m, 02) = # = 1 (y-m)2 202 ∝ (02) = e - 1/202 × (41-11)2 The propost. : P(M, 02/4) & P(M, 02). L(y 1M, 02) = 02 x (0) 2 · e - 102 [1/2 - 4) 2 + n (y-1) 2] = 152) not e - 202 [(n-1)52+n(y-4)2] Normalize: $p(u, \sigma^2 | y) = \frac{(h-1)^2}{r^2} \frac{1}{(\sigma^2)^{\frac{n+1}{2}}} e^{-\frac{(n-1)^2}{2\sigma^2}} \frac{1}{\sqrt{n}} e^{-\frac{n(y-u)}{2\sigma^2}}$ P(M/y) = 500 P(M, 02/y) do2. (+n-1=tig>)= P(=) P(=) (n-1) (n-1) (1+ n-1 (2/5n)=)= Scale: 52 = 4.2772 = 0,918 P(9.5) J19T . 42797 . [1+ q+ (42797/50)2]10 post mean = = = 50. => 9-70 CS for 11: [147.997, [2.003) (R. 10+9t(cc.ors, .975),19>* sqit (0.9158))

$$|27| P(0^{2}|y) \propto \int_{-\infty}^{\infty} \frac{1}{(0^{2})^{\frac{n}{2}+1}} e^{-\frac{1}{20^{2}}} \frac{1}{(10^{-1})^{\frac{n}{2}+n}} \frac{1}{(9^{-10})^{\frac{n}{2}}} dn$$

$$|27| \frac{1}{(0^{2})^{\frac{n}{2}+1}} e^{-\frac{1}{20^{2}}} \int_{-\infty}^{\infty} \frac{1}{0^{-\frac{n}{2}}} e^{-\frac{n(y-m)^{\frac{n}{2}}}{30^{2}}} dm$$

$$|28| \frac{1}{(0^{2})^{\frac{n}{2}+1}} e^{-\frac{1}{20^{2}}} \int_{-\infty}^{\infty} \frac{1}{0^{-\frac{n}{2}}} e^{-\frac{n(y-m)^{\frac{n}{2}}}{30^{2}}} dm$$

$$|28| \frac{1}{(0^{2})^{\frac{n}{2}+1}} e^{-\frac{1}{20^{2}}} e^{-\frac{n(y-m)^{\frac{n}{2}}}{30^{2}}} e^{-\frac{n(y-m)^{\frac{n}{2}}}{30^{2}}} dm$$

$$|29| \frac{1}{(0^{2})^{\frac{n}{2}+1}} e^{-\frac{1}{20^{2}}} e^{-\frac{n(y-m)^{\frac{n}{2}}}{30^{2}}} e^{-\frac{n(y-m)^{\frac{n}{2}}}{30^{2}}} e^{-\frac{n(y-m)^{\frac{n}{2}}}{30^{2}}} dm$$

$$|29| \frac{1}{(0^{2})^{\frac{n}{2}+1}} e^{-\frac{n(y-m)^{\frac{n}{2}}}{30^{2}}} e^{-\frac{n(y-m)^{\frac{n}{2}$$

Scanned by CamScanner

1), Cong. Prior: P(M, 02) = P(02) P(M/02) = 26/(0, p) × N(Mo, x) = 76, (38, ayy) × N(51, 52) = 444 20 1 102)39 e - 444 110 e - 10(M-51)2 P(M, 0 / 4) x 1/8 e - K(M-M) 7 n (M-y)2 or Tre-exin) (M- KNO+ny) = 110 e-130) (M-510+1000)2 > 11,02/4 ~ 16(\(\alpha + \frac{n}{2}\), \(\beta + \frac{(n-1) \(\sigma^2\)}{2} + \frac{\kn(\sigma' - mo)^2}{2} \times A(\kuo + mg) \(\sigma^2\)\\
\times \(\lambda + \frac{n}{2}\), \(\kappa + \frac{n =76 138+ 20, 444+ 348 + 10×20×150-513) × N (10×51+20×50 02) = IG (48, 64, 533) × N (50.333, 52) D) Penty) is t density w/ mean: KNO+ NY = 10×51 + 20×50 = 10,333 gade parameter No 502 + (n-1)52 + knly-10)2 (kin) (noth) = (19×46-7368+348+ 10×20 (50-51))/(30×19+20) $df = \frac{0.4315}{1000}$ 95% CS: 11 ± t39, 10.95) | Scale para. = (48,2488,52,4179)

Scanned by CamScanner

	137. P(02/4) ~261 (= +a, Z(yi-1)2+B)
	$=26(\frac{20}{2}+38,\frac{368}{2}+444)=26(48,621.333)$
	mean = $\frac{B'}{\sigma^2-1} = \frac{628}{48-1} = 13,361 + E(\sigma^2/4)$
	95% is for 52: (10.04799, 17.74442)
Extra Credit:	
7.3	$P(M, \sigma^2) = \frac{\beta^2}{P(M)} \frac{1}{(\sigma^2)^{\alpha+1}} e^{-\frac{\beta}{\alpha^2}} \frac{TK}{J_{2m\sigma^2}} e^{-\frac{K(M-M_{\sigma})^2}{2\sigma^2}}$
	where $\alpha = \frac{n_0}{2}$, $\beta = \frac{n_0 \sigma_0^2}{2}$
	$(10^{10}) \times 10^{2} $
	$\Rightarrow \frac{1}{5^2} \text{(as } n_0 \Rightarrow -1, k_0 \circ \sigma_0^2 \Rightarrow 0$
	7 62 (NS 110 - 1)
Extra Credit:	$1 = (h-1)^{2} + h(M/N)^{2} \qquad (2.17)$
7.4.	$\int_{0}^{\infty} \frac{1}{(5^{2})^{\frac{n}{2}+1}} e^{\frac{-(n-1)s^{2}+n(m-\bar{y})^{2}}{2\sigma^{2}}} d\sigma^{2} = \frac{\Gamma(\frac{n}{2})}{\frac{[n-1)s^{2}+(n-\bar{y})^{2}}{2}} \frac{1}{2}$
	$= P(\frac{n}{2}) \qquad por [(n-1)s^2 + n(m-y)^2 - \frac{n}{2}] \qquad -(n-1)s^2 + n(y-m)$
	$=\frac{P(\frac{n}{2})}{[n-1)s^{2}+[m-\bar{y})^{2}}\int_{0}^{\infty}\frac{[(n-1)s^{2}+n(m-\bar{y})^{2}]^{\frac{n}{2}}}{[(n-1)s^{2}+(m-\bar{y})^{\frac{n}{2}+1}e^{\frac{n}{2}}]}\frac{1}{2\sigma^{2}}\frac{-(n-1)s^{2}+n(y-m)}{2\sigma^{2}}$
	$= \sqrt{\left(\frac{n}{2}, \frac{(n-1)s^2 + n(y-u)^2}{2}\right)} d\sigma^2$
	$=\frac{\Gamma^{2}(\frac{h}{2})}{\Gamma(h-1)S^{2}+(M-\frac{1}{2})^{2}}$
	Scanned by CamScanner