

## Regional Environmental Change

# PREDICTING MACROALGAL SHIFTS: CLIMATE AND ENVIRONMENTAL DRIVERS IN THE CANARY ISLANDS (NE ATLANTIC OCEAN) OVER 28 YEARS

--Manuscript Draft--

|                           |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|---------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Manuscript Number:</b> | REEC-D-25-00040                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| <b>Full Title:</b>        | PREDICTING MACROALGAL SHIFTS: CLIMATE AND ENVIRONMENTAL DRIVERS IN THE CANARY ISLANDS (NE ATLANTIC OCEAN) OVER 28 YEARS                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
| <b>Article Type:</b>      | Original Article                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| <b>Keywords:</b>          | Macroalgae distribution; environmental variability; Climate change; predictive modeling; oceanic islands                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| <b>Abstract:</b>          | <p>Macroalgal communities are sensitive to environmental changes, making them key indicators for assessing climate change impacts on marine biodiversity. This study examines the relationship between environmental variables and the relative abundance of macroalgal orders Bryopsidales, Ceramiales, Dictyotales, and Fucales across the Canary Islands over 28 years. Significant multi-variable correlations were found, reflecting temporal environmental changes. While some models explained substantial variance, high environmental heterogeneity across the archipelago reduced model precision. Grouping islands into eastern and western regions and applying Generalized Additive Models (GAMs) improved predictive accuracy, emphasizing the value of region-specific modeling approaches. Principal Component Analysis (PCA) revealed an environmental gradient influencing macroalgal distributions, with nutrient-rich eastern islands showing stronger correlations for certain macroalgae. Environmental factors such as sea surface temperature, salinity, and nutrient levels were key drivers of distribution, varying by region and macroalgal order. For example, Bryopsidales and Ceramiales showed stable responses in nutrient-rich eastern waters, while temperate species in western islands were more sensitive to warming, highlighting vulnerability to climate change. The findings highlight the need for predictive models integrating environmental variables and ecological interactions, particularly in heterogeneous regions like the Canary Islands. Incorporating fine-scale biogeochemical data into future models will enhance our understanding of species shifts under climate change, identifying areas at risk of biodiversity loss and tropicalization. Long-term satellite data and ecological records are essential for devising effective conservation strategies to safeguard marine biodiversity in response to climate-driven changes.</p> |

Word count: 7620

Nº of figures: 5

Nº of tables: 4

Nº of references: 91

## **PREDICTING MACROALGAL SHIFTS: CLIMATE AND ENVIRONMENTAL DRIVERS IN THE CANARY ISLANDS (NE ATLANTIC OCEAN) OVER 28 YEARS**

**Eros F. Geppi<sup>1</sup>, Daniel González-Aragón<sup>2</sup> & Rodrigo Riera<sup>1\*</sup>**

<sup>1</sup>BIOCON, IU-ECOQUA, University of Las Palmas de Gran Canaria, Gran Canaria, Canary Is., Spain

\*corresponding author: [rodrigo.riera@ulpgc.es](mailto:rodrigo.riera@ulpgc.es)

<sup>2</sup>Departamento de Ecología, Facultad de Ciencias, Universidad Católica de la Santísima Concepción, Chile

### **ABSTRACT**

Macroalgal communities are sensitive to environmental changes, making them key indicators for assessing climate change impacts on marine biodiversity. This study examines the relationship between environmental variables and the relative abundance of macroalgal orders Bryopsidales, Ceramiales, Dictyotales, and Fucales across the Canary Islands over 28 years. Significant multi-variable correlations were found, reflecting temporal environmental changes. While some models explained substantial variance, high environmental heterogeneity across the archipelago reduced model precision. Grouping islands into eastern and western regions and applying Generalized Additive Models (GAMs) improved predictive accuracy, emphasizing the value of region-specific modeling approaches. Principal Component Analysis (PCA) revealed an environmental gradient

1  
2  
3  
4 influencing macroalgal distributions, with nutrient-rich eastern islands showing stronger  
5  
6 correlations for certain macroalgae. Environmental factors such as sea surface temperature,  
7  
8 salinity, and nutrient levels were key drivers of distribution, varying by region and macroalgal  
9  
10 order. For example, Bryopsidales and Ceramiales showed stable responses in nutrient-rich eastern  
11  
12 waters, while temperate species in western islands were more sensitive to warming, highlighting  
13  
14 vulnerability to climate change. The findings highlight the need for predictive models integrating  
15  
16 environmental variables and ecological interactions, particularly in heterogeneous regions like the  
17  
18 Canary Islands. Incorporating fine-scale biogeochemical data into future models will enhance our  
19  
20 understanding of species shifts under climate change, identifying areas at risk of biodiversity loss  
21  
22 and tropicalization. Long-term satellite data and ecological records are essential for devising  
23  
24 effective conservation strategies to safeguard marine biodiversity in response to climate-driven  
25  
26 changes.  
27  
28  
29  
30  
31  
32

33  
34 **Keywords:** Macroalgae distribution, environmental variability, climate change, predictive  
35  
36 modeling, oceanic islands.  
37  
38  
39  
40  
41

## 42 INTRODUCTION

43  
44 Climate change has introduced simultaneous shifts in temperature, ocean circulation, and  
45  
46 stratification, producing significant and potentially widespread negative effects on marine  
47  
48 ecosystems ([Waycott et al., 2009](#); [Brothers et al., 2018](#); [Turschwell et al., 2021](#)). Rising ocean  
49  
50 temperatures contribute to sea level rise ([Jevrejeva et al., 2008](#); [Alcántara et al., 2022](#)), reduced  
51  
52 sea-ice extent ([Hunt et al., 2022](#)), and changes in ocean circulation ([Noisette et al., 2022](#)), all of  
53  
54 which drive accelerated degradation of marine habitats. Coastal systems, in particular, are  
55  
56 experiencing severe loss, with global declines including 50% of salt flats, 35% of mangroves, 30%  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 of coral reefs, and 29% of seagrasses (Jackson, 2010). These environmental changes force marine  
5  
6 species to adapt, migrate, or face local extinction if conditions exceed their tolerance (Parmesan,  
7  
8 2006; Somero, 2010, 2012). In oceanic ecosystems, shifts in species distributions due to these  
9  
10 changes are altering community structures and biodiversity (Baselga, 2013), with notable impacts  
11  
12 on foundational species like ecosystem-forming seaweeds (Martínez et al., 2017), which are  
13  
14 critical to the stability of coastal ecosystems.  
15  
16  
17  
18  
19

20 Predicting how habitat-forming macroalgae will respond to climate change is essential due  
21  
22 to their role in providing food, shelter, and stabilizing ecosystem dynamics (Halpern et al., 2007;  
23  
24 Thomsen et al., 2010). Shifts in these macroalgae populations, such as potential acclimatization or  
25  
26 declines, have already been documented in several ecologically significant species (Takolander et  
27  
28 al., 2017; Riera et al., 2015; Franco et al., 2016). Long-term datasets, such as herbarium records  
29  
30 and remote sensing data, are valuable in studying these changes over decades, helping distinguish  
31  
32 climate-driven trends from natural variability (Ducklow et al., 2009; Beck, 2017; McClain, 2009).  
33  
34 Integrating these biological data with environmental metrics enables the application of remote  
35  
36 sensing techniques to monitor and predict shifts in habitat-forming species distributions under  
37  
38 changing climate conditions (Martínez et al. 2018; Kanavaugh et al. 2021).  
39  
40  
41  
42  
43  
44

45 Generalized Additive Models (GAMs) are particularly useful for analyzing long-term  
46  
47 datasets, as they allow for the exploration of non-linear relationships between environmental  
48  
49 variables and species distributions (Hastie et al. 2017). GAMs can integrate multiple predictor  
50  
51 variables to model species responses to environmental factors, providing insights into the  
52  
53 conditions that influence their distribution and abundance (Yee & Mitchell, 1991; Stoner et al.,  
54  
55 2011). These models have proven effective in studying complex ecosystems, but their accuracy  
56  
57 often depends on the availability of extensive, region-specific data, which can be challenging to  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 acquire ([Grim, 1994](#); [Drexler & Ainsworth, 2013](#)). The Canary Islands, with their distinct  
5  
6 oceanographic and ecological characteristics, offer an ideal location for applying GAMs to study  
7  
8 marine and terrestrial ecosystems. The pronounced environmental gradient across the archipelago,  
9  
10 shaped by the variability in Canary currents ([Barton et al. 1998](#); [Noisette et al., 2022](#)), creates a  
11  
12 natural laboratory for exploring non-linear species-environment relationships. From the nutrient-  
13  
14 rich waters of the eastern islands like Lanzarote and Fuerteventura to the warmer, more stable  
15  
16 ecosystems of western islands such as La Palma and El Hierro, the region provides diverse and  
17  
18 contrasting conditions that can be modeled effectively using GAMs. Additionally, the islands'  
19  
20 geographic, climatic, and geological gradients ([Hanz et al., 2022](#)) enable the study of species with  
21  
22 restricted geographic distributions. Their wealth of unique habitats and long-term environmental  
23  
24 datasets enhances our understanding of how local and regional ecological factors influence  
25  
26 biodiversity, despite challenges posed by anthropogenic pressures in this archipelago ([Riera et al.](#)  
27  
28 [2014](#)), considering the spatial scale in the studied regions ([MacArthur et al. 1992](#))  
29  
30  
31  
32  
33  
34  
35  
36

37 In this study, we use [herbarium records](#) of four macroalgae orders (Bryopsidales,  
38  
39 Ceramiales, Dictyotales, and Fucales) from the Canary Islands to explore environmental  
40  
41 interactions affecting macroalgal populations over the past three decades (1993–2020). As  
42  
43 ecosystem engineers, these macroalgae support biodiversity and stabilize coastal habitats in the  
44  
45 Canary Islands (e.g., [Valdazo et al. 2017](#); [Casado-Abellán & Riera, 2024](#); [González-Aragón et al.](#)  
46  
47 [2024](#)). They enhance ecosystem resilience by providing habitat and contributing to nutrient  
48  
49 cycling, which sustains marine species and supports local fisheries ([Mineur et al. 2015](#)). By  
50  
51 analyzing species abundance against temporal records of sea surface temperature, salinity, and  
52  
53 height for the entire archipelago, individual islands, and regional groupings (Western and Eastern  
54  
55 islands), we aim to model the dependence of these macroalgae orders on climate-sensitive  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 environmental factors using GAMs. This approach will provide insights into keystone  
5  
6 macroalgae's vulnerability and adaptive responses in the Canarian coastal ecosystems.  
7  
8  
9

## 10 11 12 MATERIAL AND METHODS 13 14 15 16 17

### 18 *Study area* 19 20

21 The study area defined for this study is the Canary Archipelago (NE Atlantic Ocean). The data  
22  
23 used in this study were collected separately for each island, allowing for distinct information on  
24  
25 each island to be analyzed independently. In addition, considering the oceanographic  
26  
27 characteristics and the environmental gradient that the archipelago presents (Barton et al., 1998;  
28  
29 Riera & Delgado, 2019), the area was classified into two large groups of islands. The western  
30  
31 islands, conformed by El Hierro, La Palma, La Gomera and Tenerife, with tropical affinities in  
32  
33 their marine flora (Tuya et al. 2004) (Fig. 1). Eastern islands conformed by Gran Canaria,  
34  
35 Lanzarote and Fuerteventura characterized by a cold-temperate flora (Tuya et al. 2004).  
36  
37  
38  
39  
40

### 41 *Macroalgae records* 42 43 44

45 A total of 11,304 records of four macroalgae orders were used. Namely, 2,157 records of  
46  
47 Bryopsidales; 4,081 records of Ceramiales; 1,784 records of Dictyotales; and 3,282 records of  
48  
49 Fucales. These specimens were collected in 5,437 locations throughout the seven islands of the  
50  
51 Canary archipelago in a 28-year time interval (from 1992 to 2020) (Fig 2). The samples used were  
52  
53 obtained from the records of the Canary Island Biodiversity Data Bank (Government of the Canary  
54  
55 Islands, 2022) and some records deposited in the Herbarium BCM database from the Marine  
56  
57 Science Faculty (University of Las Palmas de Gran Canaria, 2021).  
58  
59  
60  
61  
62  
63  
64  
65

## ***Environmental data***

Environmental variables from the Copernicus Marine Service ([Dréville et al. 2021](#)) were utilized in this study. The variables chosen were the Sea Surface Temperature (°C) (SST), Sea Surface Salinity (PSU), Significant Wave Height (m)(SWH), Sea Surface Radiation ( $\text{J m}^{-2}$ )(SSR), nitrate ( $\mu\text{M/L}$ )( $\text{NO}_3$ ), and phosphate ( $\mu\text{M/L}$ )( $\text{PO}_4$ ), due to the relevant effects produced on macroalgae communities in the ocean ([Sunny, 2017](#)). These data were collected in netCDF.nc format with a spatial resolution of  $1/12^\circ$  per pixel (*ca.*  $8\text{km}^2/\text{pixel}$ ), a monthly average of data as temporal resolution ([Dréville et al. 2021](#)) and a temporal range of 28 years according to the macroalgae records.

Two sets of environmental variables were prepared: one specific to each island, and another distinguishing between western (El Hierro, La Palma, La Gomera, and Tenerife) and eastern islands (Gran Canaria, Lanzarote, and Fuerteventura) to account for the environmental gradient across the region ([Tuya et al. 2004](#)) (Fig. 3).

## ***Data processing***

To minimize spatial and temporal biases due to heterogeneity in macroalgae records, specimens were filtered by species, sampling location, and year. This process retained only one record per species at each sampling point per year (Fig. 2), preventing multiple samples of the same macroalgae at the same location and time from artificially inflating the total record.

Once the macroalgae data was filtered, a count matrix was created by each order, island, and year. From this matrix, only orders with more than 1,500 records across the 28-year study period (1992-2020) were included in the analysis (Table 1). These count values of selected orders

per year were treated as relative abundances for each island, enabling the assessment of macroalgal abundance trends throughout the archipelago over time (Fig. 2). Additionally, the total macroalgae records for each year were represented as frequencies to evaluate the distribution patterns of the response variable for each order (Bryopsidales, Ceramiales, Dictyotales, and Fucales) (Fig. 3).

With the data visualization program NASA GISS: Panoply netCDF, HDF, and GRIB Data Viewer the environmental information was treated to recapitulate area averages and convert satellite image records into 28 year time series. A filtering treatment was applied to these time series according to the sampling frequency using the function `decompose {stats::decompose()}` from the package STATS V3.6.2 in RStudio (R Core Team, 2022). This filtering had the purpose of eliminating the residual components of the time series, obtaining the trend and stationarity of each variable over time more clearly. Once the time series were treated, the information was assigned to the macroalgae counting records by statistical model selection. In this way, the macroalgae counts will have added the environmental measurements for spatial region and year.

### ***Statistical analysis***

For the statistical analysis, we first establish relationships between environmental variables alongside the time series of macroalgal relative abundance to visually assess potential relationships by Principal Component Analysis (PCA) (Fig. 4). Given the temporal variability and environmental heterogeneity across the archipelago, linear correlations between the dependent variable (relative abundance) and the independent variables (environmental drivers) were not apparent. Therefore, a Generalized Additive Model (GAM) was selected for the analysis to account for non-linear relationships and temporal trends (Wood et al. 2017).

GAM models were applied to examine the environmental effects on relative abundance across the entire archipelago (Table 2). Second, separate GAM models were developed for each island (Table 3). Finally, GAM models were created for the two island groups: Western and Eastern islands (Table 4). The GAM analysis was performed in R Studio version 2023.9.1.494 (R Core Team 2022), using cubic regression splines (“cr”) with a smoothing parameter (k) of 30 for each environmental variable. The Tweedie family ( $p = 1.5$ ) with a logarithmic link function was used, as it is well-suited for data comprising zeros and positive values, particularly when dealing with highly dispersed distributions. The model was fitted using the REML method for stable smoothing parameter estimation, and adaptive penalization allowed the removal of non-informative predictors. In addition to the aforementioned environmental variables, *Year* was included as a categorical factor to capture inter-annual variability.

The analysis was conducted in R Studio using the mgcv package (Wood et al. 2017, Pedersen et al. 2019). The different models were chosen by ANOVA comparison between several model designs, selecting the greatest explanatory capacity and the least defined complexity.

## RESULTS

Analysis of the correlations between environmental variables and macroalgal relative abundance records across the Canary Islands landscape revealed significant multi-variable relationships for each macroalgal order, corresponding to temporal variability in environmental factors. Temporal changes in environmental components accounted for 54.9% to 66.1% of the variation in macroalgal relative abundance over the 28-year time series. Despite these significant relationships, model fits did not show consistently strong alignment, likely due to high

1  
2  
3  
4 environmental variability in the Canary Islands, which affects model precision even with  
5  
6 significant correlations between predictor and response variables (Table 2).  
7  
8  
9

10 For Bryopsidales across the archipelago, environmental factors explained 66.8% of the  
11  
12 variance with four key variables, though the correlation adjustment remained low at 35%.  
13  
14 Similarly, Ceramiales variance was explained at 61.5% with five variables but yielded a moderate  
15  
16 correlation of 43.9%. Dictyotales exhibited 64.5% data variance explained by five significant  
17  
18 variables and a correlation adjustment of 30.4%, while Fucales records showed 54.9% variance  
19  
20 explained by two environmental variables with a 28.9% correlation.  
21  
22  
23  
24  
25

26 PCA indicated regional differentiation, with a gradient from positive PC1 values in the  
27  
28 western islands (e.g., La Palma and El Hierro) to negative PC1 values in the eastern islands (Fig.  
29  
30 4). This regional gradient suggests that temporal series should be analyzed on an island-by-island  
31  
32 basis to account for region-specific environmental variables affecting macroalgal distributions.  
33  
34 The model selection process led to the establishment of Generalized Additive Models (GAMs) for  
35  
36 each macroalgal order per region. Despite using the same variables for all models, regional  
37  
38 variability caused significant differences in fit and explained variance, likely due to both sample  
39  
40 heterogeneity and environmental variation across islands. The observed differences in  $R^2$  values  
41  
42 and explanatory deviation highlight the influence of regional environmental heterogeneity on  
43  
44 model accuracy (Table 3).  
45  
46  
47  
48  
49  
50

51 When records were grouped into two broad regions—western islands (El Hierro, La Palma,  
52  
53 La Gomera, and Tenerife) and eastern islands (Lanzarote, Fuerteventura, and Gran Canaria)—  
54  
55 environmental variables showed greater model fit and explanatory power than models-based on  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

individual islands or the entire archipelago (Table 4). This grouping revealed that specific environmental variables impacted macroalgal orders differently in each region.

For Bryopsidales in the western islands, three environmental variables, i.e., Sea Surface Temperature (SST), Sea Surface Radiation (SSR), and Salinity explained 85% of the variance with a correlation of 0.763. In the eastern islands, significant variables included SSR, Salinity, and Significant Wave Height (SWH), achieving a stronger fit with a 92.5% variance explained and a correlation of 0.852. For Ceramiales, in the western region, four environmental factors (SST, SWH, Nitrate ( $\text{NO}_3$ ), and Phosphate ( $\text{PO}_4$ )) explained 84.9% of the variance with a 0.77 correlation. In the eastern islands, only three variables were necessary (excluding  $\text{PO}_4$ ), yielding 90.5% variance explained and a 0.782 correlation. For Dictyotales, variance explained reached 77.5% in the west with three variables (SST, Salinity,  $\text{PO}_4$ ) but showed a lower correlation of 0.511, likely reflecting fewer samples. In the eastern region, five variables (SSR, SWH, Salinity,  $\text{NO}_3$ , and  $\text{PO}_4$ ) improved the variance explained to 82.7% with a 0.733 correlation. Finally, for Fucales, two environmental variables (SST and SSR) in the western region explained 81.6% of the variance with a correlation of 0.79. In the east, the addition of SWH and  $\text{PO}_4$  resulted in a slightly lower variance explained (73.5%) and a correlation of 0.653.

Using island-specific models and grouping islands into two main regions—western (El Hierro, La Palma, La Gomera, and Tenerife) and eastern islands (Lanzarote, Fuerteventura, and Gran Canaria)—improved model fit and accuracy, as environmental factors influenced macroalgal orders more predictably within each region. In the western islands, Bryopsidales variance was explained at 85% by SST, Sea Surface Radiation (SSR), and Salinity, with a correlation of 0.763. In the eastern islands, the inclusion of SSR, Salinity, and Significant Wave Height (SWH) yielded a stronger fit, with 92.5% variance explained and a correlation of 0.852. For Ceramiales, four

variables (SST, SWH, NO<sub>3</sub>, and PO<sub>4</sub>) explained 84.9% of the variance in the western region, while in the eastern region, three factors excluding PO<sub>4</sub> improved the variance explained to 90.5%, with a correlation of 0.782.

## DISCUSSION

The main results of this study illustrate a clear decline in macroalgal populations across the Canary Islands, which is primarily driven by human-driven activities and environmental factors showing considerable spatial and temporal variability. Significant differences exist between the western and eastern islands, including SST, nutrient concentrations, and salinity levels, each of which influences macroalgal distributions differently (Tuya et al. 2006; Barton et al. 1998). The environmental characteristics of the Canary archipelago—observed through satellite data—emphasize that environmental components distinguish regions and contribute to contrasting macroalgal dynamics at a regional level across time.

Analyses of environmental variables against long-term macroalgal relative abundance records indicate significant multi-variable relationships for each macroalgal order across each island, reflecting temporal changes in environmental conditions. Temporal variability accounted for 54.9% to 66.1% of the variance in macroalgal relative abundance over a 28-year time series. However, high environmental variability across the archipelago limits model accuracy, reflected in moderate correlation adjustments despite significant predictor-response relationships. For instance, the Bryopsidales model explained 66.8% of the variance with four significant variables but had only a 35% correlation, revealing challenges due to regional heterogeneity.

1  
2  
3  
4 The present results align with previous research showing that environmental stressors  
5  
6 disproportionately impact macroalgal populations based on local tolerance thresholds for SST,  
7  
8 salinity, and nutrient availability (Zou and Gao, 2013; Wernberg et al. 2012; Staehr and Wernberg,  
9  
10 2009; Pekker et al. 1996). Former studies revealed that environmental heterogeneity in the Canary  
11  
12 Islands influences the connectivity and dispersal of temperate-water macroalgae, such as Fucales,  
13  
14 which face declines due to increased SST and SSH in warmer waters of the western islands (Duarte  
15  
16 et al. 2013; Riera et al. 2015). In contrast, lower temperatures and higher nutrient input in the  
17  
18 eastern islands provide a temporary buffer against warming impacts, favoring specific taxa  
19  
20 (Afonso et al. 2013; Ribeiro et al. 2019).  
21  
22  
23  
24  
25  
26

27 GAMs emphasized the significance of regional environmental variability in shaping  
28  
29 macroalgal distributions, with island grouping models outperforming those for individual islands  
30  
31 or the entire archipelago by capturing localized heterogeneity. This aligns with PCA, which  
32  
33 revealed an environmental gradient: positive PC1 values in the western islands and negative values  
34  
35 in the eastern islands, driven by distinct regional factors. Bryopsidales and Ceramiales, for  
36  
37 instance, showed stronger associations with nutrient-rich, stable eastern waters. While these  
38  
39 models provide insights into environmental influences on macroalgal abundance, they do not fully  
40  
41 predict species metacommunity dynamics due to the complex interplay of species traits and  
42  
43 ecological interactions. Reliable predictions require integrating biological characteristics with  
44  
45 environmental factors through frameworks like species distribution modeling. Spatial  
46  
47 heterogeneity in the Canary Islands further complicates predictions, with unique temporal  
48  
49 fluctuations at each site limiting model performance (Riera et al. 2015). Broader data collection  
50  
51 and localized assessments are essential to complement large-scale models, addressing fine-scale  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

variability and accurately reflecting species dynamics in heterogeneous landscapes (MacArthur, et al. 1992; Zou & Gao, 2013).

As climate change progresses, the variability in SST, SSH, and Salinity could exacerbate regional declines in macroalgal diversity, particularly in western islands, where populations already display sensitivity to warming (Valdazo et al. 2024). These shifts may allow species with tropical affinities to expand, while temperate species decline (Perry et al. 2005; Martínez et al. 2017; Franco et al. 2018). If warming continues as projected by the IPCC (2020), this shift could trigger cascading effects on ecosystem function and biodiversity (Smale and Wernberg, 2013; Bracken et al. 2007), potentially leading to local extinctions due to limited adaptation capacity linked to genetic diversity loss (Pearson et al. 2009). As observed in other studies, a decrease in habitat-specific species could favor tropical species suited to warmer conditions, reshaping community composition (Vergés et al. 2016; Valdazo et al. 2017; Ribeiro et al. 2019).

The present study reinforces the importance of developing statistical models that incorporate both physical and biogeochemical interactions to better predict long-term impacts on marine ecosystems. Enhanced regional models integrating SST, salinity, nutrient levels, and other variables could provide clearer insights into climate change effects and support targeted conservation strategies (Martínez et al. 2012; Bennett et al. 2018). These models should also identify climate refugia—areas that are less affected by climate change and can serve as safe havens for species under stress. Preserving these refugia is vital for sustaining native macroalgal species in the Canary Islands and ensuring their role in maintaining marine biodiversity (Riera & Delgado, 2019). By prioritizing the protection of regions with stable environmental conditions and high resilience to climate shifts, conservation efforts can help mitigate the adverse effects of climate change. This approach aligns with calls for adaptive management strategies to safeguard

1  
2  
3  
4 marine biodiversity under the growing climate crisis ([Wernberg et al. 2011](#); [Riera et al. 2015](#);  
5  
6 [Vergés et al. 2014, 2016](#)). Identifying and preserving such critical areas will enhance ecosystem  
7  
8 stability and offer a foundation for the recovery of vulnerable species.  
9

## 10 11 12 13 14 **CONCLUSIONS**

15  
16  
17 Integrating satellite data with long-term biological records has provided valuable insights into how  
18  
19 the unique environmental conditions of the Canary archipelago shape macroalgal tolerance and  
20  
21 distribution. This research underscores the importance of incorporating diverse environmental  
22  
23 variables into population studies, particularly in complex marine ecosystems like the Canaries.  
24  
25 Scaling up these findings to other regions requires the development of spatially consistent and  
26  
27 temporally continuous sampling frameworks, enabling more precise and globally relevant  
28  
29 predictions for conservation.  
30  
31  
32

33  
34  
35 Future research should prioritize identifying areas most vulnerable to climate impacts or  
36  
37 those that could serve as refugia for native species. Incorporating detailed biogeochemical  
38  
39 variables, high-resolution datasets, and refined ecological models will be crucial for accurately  
40  
41 predicting species dynamics and pinpointing regions susceptible to biodiversity shifts, including  
42  
43 the establishment of invasive tropical species. Conservation strategies must focus on mitigating  
44  
45 the loss of macroalgal populations by safeguarding critical habitats essential for their persistence.  
46  
47  
48  
49  
50

51  
52 This study highlights the broader applicability of its methods and findings, providing a  
53  
54 foundation for predictive modeling in other biodiverse and environmentally heterogeneous  
55  
56 regions. Proactive measures are essential to preserve biodiversity in the face of ecosystem shifts  
57  
58 driven by climate change. Escalating these predictive models can inform adaptive management  
59  
60  
61  
62  
63  
64  
65

plans to diverse marine landscapes, ensuring the protection of both macroalgal communities and the broader ecosystems they support. By refining these approaches, researchers and policymakers can develop robust conservation frameworks to address the global challenges posed by climate-driven changes in marine biodiversity.

## ACKNOWLEDGEMENTS

This contribution is based on the MSc thesis of the lead author (EG). We extend our gratitude to Prof. Antonio Juan González Ramos (ULPGC) for his valuable guidance on the use of satellite data.

## REFERENCES

1. **Adebowale, M. (2021).** Rate and amplitudes of climate change: Are there tipping points in marine ecosystems? *University of Bristol*. Retrieved from <http://research-information.bristol.ac.uk>
2. **Alcántara, L., Creencia, L., Madarcos, K., Roderick-Madarcos, J., Beth-Jontila, J., & Culhane, F. (2022).** Perceptions of climate change, sea level rise, and factors affecting the coastal marine ecosystem of Palawan, Philippines. *UCL Open Environment*. <https://doi.org/10.14324/111.444/000150.v1>
3. **Barton, E. D., Arístegui, J., Tett, P., Cantón, M., & García-Braun, J., et al. (1998).** The transition zone of the Canary Current upwelling region. *Progress in Oceanography*, 41(4–5), 455–504. [https://doi.org/10.1016/S0079-6611\(98\)00023-8](https://doi.org/10.1016/S0079-6611(98)00023-8)
4. **Baselga, A. (2010).** Partitioning the turnover and nestedness components of beta diversity. *Global Ecology and Biogeography*, 19(1), 134–143. <https://doi.org/10.1111/j.1466-8238.2009.00490.x>
5. **Baselga, A. (2013).** Separating the two components of abundance-based dissimilarity: Balanced changes in abundance vs. abundance gradients. *Methods in Ecology and Evolution*, 4(6), 552–557. <https://doi.org/10.1111/2041-210X.12029>
6. **Beck, H. J., Feary, D. A., Nakamura, Y., & Booth, D. J. (2017).** Temperate macroalgae impacts tropical fish recruitment at the forefronts of range expansion. *Coral Reefs*, 36(2), 639–651. <https://doi.org/10.1007/s00338-017-1553-1>
7. **Bernal-Ibáñez, A., Cacabelos, E., Melo, R., & Gestoso, I. (2021).** The role of sea-urchins in marine forests from Azores, Webbnesia, and Cabo Verde: Human pressures, climate-change effects and restoration opportunities. *Frontiers in Marine Science*, 8, Article 649873. <https://doi.org/10.3389/fmars.2021.649873>
8. **Brothers, C. J., Van der Pol, W. J., Morrow, C. D., Hakim, J. A., Koo, H., & McClintock, J. B. (2018).** Ocean warming alters predicted microbiome functionality in a common sea urchin.

- Proceedings of the Royal Society B: Biological Sciences*, 285, Article 20180340. <https://doi.org/10.1098/rspb.2018.0340>
9. **Buonomo, R., Chefaoui, R. M., Lacida, R. B., Engelen, A. H., Serrão, E. A., & Airoidi, L. (2018).** Predicted extinction of unique genetic diversity in marine forests of *Cystoseira* spp. *Marine Environmental Research*, 138, 119–128. <https://doi.org/10.1016/j.marenvres.2018.04.013>
  10. **Government of the Canary Islands. (2022).** Canary Islands Biodiversity Database. Retrieved from <https://www.biodiversidadcanarias.es/biota>
  11. **Casado-Abellán, I., & Riera, R. (2024).** Epifaunal community associated with *Cystoseira humilis* in tide pools: Assessing the significance of depth and size as pivotal factors. *Cahiers de Biologie Marine*, 65(3), 173–184. <https://doi.org/10.21411/CBM.A.B63CA70E>
  12. **Clarke, A. (2003).** Costs and consequences of evolutionary temperature adaptation. *Trends in Ecology and Evolution*, 18(11), 573–581. <https://doi.org/10.1016/j.tree.2003.08.007>
  13. **Constable, A. J., Melbourne-Thomas, J., Corney, S. P., Arrigo, K. R., Barbraud, C., et al. (2014).** Climate change and Southern Ocean ecosystems I: How changes in physical habitats directly affect marine biota. *Global Change Biology*, 20(10), 3004–3025. <https://doi.org/10.1111/gcb.12623>
  14. **Dré villon, M., Greiner, E., Hernandez, F., Rautenhaus, M., Levier, B., Rémy, E., & De Nicola, C. (2021).** Environmental variables from the Copernicus Marine Service. *Copernicus Marine Environment Monitoring Service (CMEMS)*. Retrieved from <https://marine.copernicus.eu>
  15. **Diehl, N., Karsten, U., & Bischof, K. (2020).** Impacts of combined temperature and salinity stress on the endemic Arctic brown seaweed *Laminaria solidungula* J. Agardh. *Polar Biology*, 43(6), 647–656. <https://doi.org/10.1007/s00300-020-02668-5>
  16. **Doney, S. C., Ruckelshaus, M., Duffy, E. J., Barry, J. P., Chan, F., et al. (2012).** Climate change impacts on marine ecosystems. *Annual Review of Marine Science*, 4, 11–37. <https://doi.org/10.1146/annurev-marine-041911-111611>
  17. **Drexler, M., & Ainsworth, C. H. (2013).** Generalized additive models used to predict species abundance in the Gulf of Mexico: An ecosystem modeling tool. *PLoS ONE*, 8(5), e64458. <https://doi.org/10.1371/journal.pone.0064458>
  18. **Duarte, C. M., Losada, I. J., Hendriks, I. E., Mazarrasa, I., & Marbà, N. (2013).** The role of coastal plant communities for climate change mitigation and adaptation. *Nature Climate Change*, 3(11), 961–968. <https://doi.org/10.1038/nclimate1970>
  19. **Duarte, B., Martins, I., Rosa, R., Matos, A. R., Roleda, R., et al. (2018).** Climate change impacts on seagrass meadows and macroalgal forests: An integrative perspective on acclimation and adaptation potential. *Frontiers in Marine Science*, 5, Article 190. <https://doi.org/10.3389/fmars.2018.00190>
  20. **Ducklow, H. W., Doney, S. C., & Steinberg, D. K. (2009).** Contributions to long-term research and time-series observations of marine ecology and biogeochemistry. *Annual Review of Marine Science*, 1, 279–302. <https://doi.org/10.1146/annurev.marine.010908.163801>
  21. **Eggert, A., Visser, R. J. W., van Hasselt, P. R., & Breeman, A. M. (2006).** Differences in acclimation potential of photosynthesis in seven isolates of the tropical to warm temperate macrophyte *Valonia utricularis* (Chlorophyta). *Phycologia*, 45(5), 546–556. <https://doi.org/10.2216/05-03.1>
  22. **Fahrig, L. (2003).** Effects of habitat fragmentation on biodiversity. *Annual Review of Ecology, Evolution, and Systematics*, 34(1), 487–515. <https://doi.org/10.1146/annurev.ecolsys.34.011802.132419>
  23. **Franco, J. N., Tuya, F., Bertocci, I., Rodríguez, L., Martínez, B., et al. (2018).** The ‘golden kelp’ *Laminaria ochroleuca* under global change: Integrating multiple eco-physiological responses with species distribution models. *Journal of Ecology*, 106(1), 47–58. <https://doi.org/10.1111/1365-2745.12810>

24. **Geppi, E. F., & Riera, R. (2022).** Responses of intertidal seaweeds to warming: A 38-year time series shows differences in sizes. *Estuarine, Coastal and Shelf Science*, 270, Article 107841. <https://doi.org/10.1016/j.ecss.2022.107841>
25. **González-Aragón, D., Núñez, J., & Riera, R. (2024).** Epifauna-associated community of tide pools dominated by *Cystoseira humilis* from the NE Atlantic Ocean. *Cahiers de Biologie Marine*, 65, 115–123. <https://doi.org/10.21411/CBM.A.FD71E6D9>
26. **Grimm, V. (1994).** Mathematical models and understanding in ecology. *Ecological Modelling*, 75/76, 641–651. [https://doi.org/10.1016/0304-3800\(94\)00026-E](https://doi.org/10.1016/0304-3800(94)00026-E)
27. **Guisan, A., Edwards, T. C., & Hastie, T. (2002).** Generalized linear and generalized additive models in studies of species distributions: Setting the scene. *Ecological Modelling*, 157, 89–100. [https://doi.org/10.1016/S0304-3800\(02\)00204-1](https://doi.org/10.1016/S0304-3800(02)00204-1)
28. **Halpern, B. S., Silliman, B. R., Olden, J. D., Bruno, J. P., & Bertness, M. D. (2007).** Incorporating positive interactions in aquatic restoration and conservation. *Frontiers in Ecology and the Environment*, 5(3), 153–160. [https://doi.org/10.1890/1540-9295\(2007\)5\[153:IPIIAR\]2.0.CO;2](https://doi.org/10.1890/1540-9295(2007)5[153:IPIIAR]2.0.CO;2)
29. **Hanz, D. M., Cutts, V., Barajas-Barbosa, M. P., Algar, A. C., Beierkuhnlein, C., Fernández-Palacios, J.-M., Field, R., Kreft, H., Steinbauer, M. J., Weigelt, P., & Irl, S. D. H. (2022).** Climatic and biogeographical drivers of functional diversity in the flora of the Canary Islands. *Global Ecology and Biogeography*, 31, 1313–1331. <https://doi.org/10.1111/geb.13507>
30. **Harley, C. D. G., Anderson, K. M., Demes, K. W., & Kordas, J. P., et al. (2012).** Effects of climate change on global seaweed communities. *Journal of Phycology*, 48(5), 1064–1078. <https://doi.org/10.1111/j.1529-8817.2012.01224.x>
31. **Harley, C. D. G., Hughes, A. R., Hultgren, K. M., Miner, B. G., & Sorte, C. J. B. (2006).** The impacts of climate change in coastal marine systems. *Ecology Letters*, 9(2), 228–241. <https://doi.org/10.1111/j.1461-0248.2005.00871.x>
32. **Hastie, T., Tibshirani, R., & Friedman, J. (2017).** *The elements of statistical learning: Data mining, inference, and prediction*. Springer. <https://doi.org/10.1007/978-0-387-84858-7>
33. **Hastie, T., Tibshirani, R., & Tibshirani, R. J. (2017).** Extended comparisons of best subset selection, forward stepwise selection, and the lasso. *arXiv preprint, arXiv:1707.08692*. <https://doi.org/10.48550/arXiv.1707.08692>
34. **Hilbe, J. M. (2011).** *Negative binomial regression* (Illustrated, revised ed.). Cambridge University Press. <https://doi.org/10.1017/CBO9780511973420>
35. **Hooper, D. U., Chapin, F. S., Ewel, J. J., Hector, A., Inchausti, P., et al. (2005).** Effects of biodiversity on ecosystem functioning: A consensus of current knowledge. *Ecological Monographs*, 75(1), 3–35. <https://doi.org/10.1890/04-0922>
36. **Hunt, G. L., Yasumiishi, E. M., Eisner, L. B., Stabeno, P. J., & Decker, M. B. (2022).** Climate warming and the loss of sea ice: The impact of sea-ice variability on the southeastern Bering Sea pelagic ecosystem. *ICES Journal of Marine Science*, 79(3), 937–953. <https://doi.org/10.1093/icesjms/fsaa206>
37. **Jackson, J. B. C. (2010).** The future of the ocean's past. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 365(1558), 3765–3778. <https://doi.org/10.1098/rstb.2010.0278>
38. **Jevrejeva, S., Moore, J. C., Grinsted, A., & Woodworth, P. L. (2008).** Recent global sea level acceleration started over 200 years ago? *Geophysical Research Letters*, 35, L08715. <https://doi.org/10.1029/2008GL033611>
39. **Kavanaugh, M. T., Bell, T., Catlett, D., Cimino, M. A., Doney, S. C., et al. (2021).** Satellite remote sensing and the marine biodiversity observation network. *Oceanography*, 34(2), 62–79. <https://doi.org/10.5670/oceanog.2021.215>
40. **Katsanevakis, S., & Maravelias, C. D. (2009).** Bathymetric distribution of demersal fish in the Aegean and Ionian Seas based on generalized additive modelling. *Fisheries Science*, 75(1), 13–23. <https://doi.org/10.1007/s12562-008-0033-5>

41. Kirby, R. R., Beaugrand, G., & Lindley, J. A. (2009). Synergistic effects of climate and fishing in a marine ecosystem. *Ecosystems*, 12(4), 548–561. <https://doi.org/10.1007/s10021-009-9241-9>
42. López-González, E., & Ruiz-Soler, M. (2011). Análisis de datos con el modelo lineal generalizado: Una aplicación con R. *Revista Española de Pedagogía*, 69(248), 59–80. Retrieved from <http://www.jstor.org/stable/23766383>
43. MacArthur, R. H., Levin, S. A., & MacArthur, S. A. L. (1992). The problem of pattern and scale in ecology. *Ecology*, 73(6), 1943–1967. <https://doi.org/10.2307/1941447>
44. Martín García, L., Rancel, N., Sangil, C., Reyes, J., Benito, B., Orellana, S., & Sansón, M. (2022). Environmental and human factors drive subtropical marine forests. *Marine Environmental Research*, 181, Article 105759. <https://doi.org/10.1016/j.marenvres.2022.105759>
45. Martínez, B., Radford, B., Thomsen, M. S., Connell, S. D., Carreño, F., et al. (2018). Distribution models predict large contractions of habitat-forming seaweeds in response to ocean warming. *Diversity and Distributions*, 24(10), 1350–1366. <https://doi.org/10.1111/ddi.12767>
46. Masson-Delmotte, V., Zhai, P., Pörtner, H.-O., Roberts, D., Skea, J., Calvo, E., Priyadarshi, B., Shukla, R., Slade, R., Connors, S., van Diemen, R., Ferrat, M., Haughey, E., Luz, S., Neogi, S., Pathak, M., Petzold, J., Pereira, J. P., Vyas, P., ... Malley, J. (2020). El cambio climático y la tierra: Resumen para responsables de políticas. *Intergovernmental Panel on Climate Change (IPCC)*. Retrieved from <https://www.ipcc.ch>
47. McClain, C. R. (2009). A decade of satellite ocean color observations. *Annual Review of Marine Science*, 1, 19–42. <https://doi.org/10.1146/annurev.marine.010908.163650>
48. Medina, M., & Haroun, H. (2004). Preliminary study on the dynamics of *Cystoseira abies-marina* populations in Tenerife. *Courier Forschungsinstitut Senckenberg*. Retrieved from <http://hdl.handle.net/10553/76923>
49. Mineur, F., Arenas, F., Assis, J., Davies, A. J., Engelen, A. H., Fernandes, F., Malta, E.-J., Thibaut, T., Van Nguyen, T., Vaz-Pinto, F., Vranken, S., Serrão, E. A., & De Clerck, O. (2015). European seaweeds under pressure: Consequences for communities and ecosystem functioning. *Journal of Sea Research*, 98, 91–108. <https://doi.org/10.1016/j.seares.2014.11.004>
50. Moisen, G. G., & Frescino, T. S. (2002). Comparing five modeling techniques for predicting forest characteristics (special issue on advances in GLM/GAM modeling). *Ecological Modelling*, 157(2–3), 209–225. [https://doi.org/10.1016/S0304-3800\(02\)00197-7](https://doi.org/10.1016/S0304-3800(02)00197-7)
51. Noisette, F., Pansch, C., Wall, M., Wahl, M., & Hurd, C. L. (2022). Role of hydrodynamics in shaping chemical habitats and modulating the responses of coastal benthic systems to ocean global change. *Global Change Biology*, 28(12), 3812–3829. <https://doi.org/10.1111/gcb.16165>
52. National Oceanic and Atmospheric Administration (NOAA). (2009). ETOPO1 1 arc-minute global relief model. *National Centers for Environmental Information*. <https://doi.org/10.7289/V5C8276M>
53. Parmesan, C. (2006). Ecological and evolutionary responses to recent climate change. *Annual Review of Ecology, Evolution, and Systematics*, 37, 637–669. <https://doi.org/10.1146/annurev.ecolsys.37.091305.30000024>
54. Pecl, G. T., Araújo, M. B., Bell, J. D., Blanchard, J., Bonebrake, T. C., Chen, I.-C., Clark, T. D., Colwell, R. K., Danielsen, F., Evengård, B., Falconi, L., Ferrier, S., Frusher, S., Garcia, R. A., Griffis, R. B., Hobday, A. J., Janion-Scheepers, C., Jarzyna, M. A., Jennings, S., Lenoir, J., Linnetved, H. I., Martin, V. Y., McCormack, P. C., McDonald, J., Mitchell, N. J., Mustonen, T., Pandolfi, J. M., Pettorelli, N., Popova, E., Robinson, S. A., Scheffers, B. R., Shaw, J. D., Sorte, C. J. B., Strugnell, J. M., Sunday, J. M., Tuanmu, M.-N., Vergés, A., Villanueva, C., Wernberg, T., Wapstra, E., & Williams, S. E. (2017). Biodiversity redistribution under climate change: Impacts on ecosystems and human well-being. *Science*, 355(6332), 1–11. <https://doi.org/10.1126/science.aai9214>
55. Pedersen, E. J., Miller, D. L., Simpson, G. L., & Ross, N. (2019). Hierarchical generalized additive models in ecology: An introduction with *mgcv*. *PeerJ*, 7, e6876. <https://doi.org/10.7717/peerj.6876>

56. **Perry, A. L., Low, P. J., Ellis, J. R., & Reynolds, J. D. (2005).** Climate change and distribution shifts in marine fishes. *Science*, 38(5730), 1912–1915. <https://doi.org/10.1126/science.1111322>
57. **Pimm, S. L., Jenkins, C. N., Abell, R., Brooks, T. M., Gittleman, J. L., Joppa, L. N., Raven, P. H., Roberts, C. M., & Sexton, J. O. (2014).** The biodiversity of species and their rates of extinction, distribution, and protection: Background rates of species extinction. *Science*, 344(6187), 1246752. <https://doi.org/10.1126/science.1246752>
58. **Ribeiro, C., Neto, A. I., Moreau, I., Haroun, R., & Neves, P. (2019).** A new signal of marine tropicalization in the Macaronesia region: First record of the mesophotic macroalga *Avrainvillea canariensis* A. Gepp & E.S. Gepp in the Madeira archipelago. *Aquatic Botany*, 153, 40–43. <https://doi.org/10.1016/j.aquabot.2018.11.008>
59. **R Core Team. (2022).** *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>
60. **Riera, R., Becerro, M. A., Stuart-Smith, R. D., Delgado, J. D., & Edgar, G. J. (2014).** Out of sight, out of mind: Threats to the marine biodiversity of the Canary Islands (NE Atlantic Ocean). *Marine Pollution Bulletin*, 86(1–2), 9–18. <https://doi.org/10.1016/j.marpolbul.2014.07.014>
61. **Riera, R., & Delgado, J. D. (2019).** Canary Islands. In *World Seas: An Environmental Evaluation* (pp. 483–500). Academic Press. <https://doi.org/10.1016/B978-0-12-805068-2.00024-3>
62. **Riera, R., Sangil, C., & Sansón, M. (2015).** Long-term herbarium data reveal the decline of a temperate-water algae at its southern range. *Estuarine, Coastal and Shelf Science*, 165, 159–165. <https://doi.org/10.1016/j.ecss.2015.05.008>
63. **Solan, M., Bennett, E. M., Mumby, P. J., Leyland, J., & Godbold, J. A. (2020).** Benthic-based contributions to climate change mitigation and adaptation. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 375, 20190107. <https://doi.org/10.1098/rstb.2019.0107>
64. **Somero, G. N. (2010).** The physiology of climate change: How potentials for acclimatization and genetic adaptation will determine “winners” and “losers.” *Journal of Experimental Biology*, 213(6), 912–920. <https://doi.org/10.1242/jeb.037473>
65. **Somero, G. N. (2012).** The physiology of global change: Linking patterns to mechanisms. *Annual Review of Marine Science*, 4, 39–61. <https://doi.org/10.1146/annurev-marine-120710-10093>
66. **SPA/RAC. (2018).** *SPA-BD Protocol - Annex II: List of endangered or threatened species*. Regional Activity Centre for Specially Protected Areas. <https://www.rac-spa.org>
67. **Steneck, R. S., & Dethier, M. N. (1994).** A functional group approach to the structure of algal-dominated communities. *Oikos*, 69(3), 476–498. <https://doi.org/10.2307/3545860>
68. **Stoner, A. W., Manderson, J. P., & Pessutti, J. P. (2001).** Spatially explicit analysis of estuarine habitat for juvenile winter flounder: Combining generalized additive models and geographic information systems. *Marine Ecology Progress Series*, 271, 213–253. <https://doi.org/10.3354/meps213253>
69. **Sunny, A. R. (2017).** A review on effect of global climate change on seaweed and seagrass communities. *International Journal of Fisheries and Aquatic Studies*, 5(6), 19–22. (<https://www.fisheriesjournal.com/archives/2017/vol5issue6/PartA/5-5-49-118.pdf>)
70. **Takolander, A., Cabeza, M., & Leskinen, E. (2017).** Climate change can cause complex responses in Baltic Sea macroalgae: A systematic review. *Journal of Sea Research*, 123, 16–29. <https://doi.org/10.1016/j.seares.2017.03.007>
71. **Thomsen, M. S., Wernberg, T., Altieri, A., Tuya, F., Gulbransen, D., McGlathery, K. J., Holmer, M., & Silliman, B. R. (2010).** Habitat cascades: The conceptual context and global relevance of facilitation cascades via habitat formation and modification. *Integrative and Comparative Biology*, 50(2), 158–175. <https://doi.org/10.1093/icb/icq042>
72. **Turschwell, M. P., Connolly, R. M., Dunic, J. C., Sievers, M., Buelow, C. A., et al. (2021).** Anthropogenic pressures and life history predict trajectories of seagrass meadow extent on a global scale. *Proceedings of the National Academy of Sciences of the United States of America*, 118, e2110802118. <https://doi.org/10.1073/pnas.2110802118>

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
73. Tuya, F., Boyra, A., Sanchez-Jerez, P., Barbera, C., & Haroun, R. J. (2004). Relationships between rocky-reef fish assemblages, the sea urchin *Diadema antillarum* and macroalgae throughout the Canarian Archipelago. *Marine Ecology Progress Series*, 278, 157–169. <https://doi.org/10.3354/meps278157>
74. Tuya, F., & Haroun, R. J. (2006). Spatial patterns and response to wave exposure of shallow water algal assemblages across the Canarian Archipelago: A multi-scaled approach. *Marine Ecology Progress Series*, 311, 15–28. <https://doi.org/10.3354/meps311015>
75. University of Las Palmas de Gran Canaria. (2021). Herbarium BCM. Retrieved from <https://www.geoportal.ulpgc.es/herbariobcm/#/>
76. Valdazo, J., Viera-Rodríguez, M. A., Espino, F., Haroun, R., & Tuya, F. (2017). Regresión aguda de los bosques de *Cystoseira abies-marina* en la isla de Gran Canaria (Islas Canarias, Atlántico este). *Scientia Marina*, 81(4), 499–507. <https://doi.org/10.3989/scimar.04655.23A>
77. Valdazo, J., Coca, J., Haroun, R., Bergasa, O., Viera-Rodríguez, M. A., & Tuya, F. (2024). Local and global stressors as major drivers of the drastic regression of brown macroalgae forests in an oceanic island. *Regional Environmental Change*, 24, 65. <https://doi.org/10.1007/s10113-024-02228-1>
78. Vélez-Rubio, G. M., González-Etchebehere, L., Scarabino, F., Trinchin, R., Manta, G., et al. (2021). Macroalgae morpho-functional groups in Southern marine ecosystems: Rocky intertidal in the Southwestern Atlantic (33°–35° S). *Marine Biology*, 168, 153. <https://doi.org/10.1007/s00227-021-03960-6>
79. Vergés, A., Doropoulos, C., Malcolm, H. A., Skye, M., Garcia-Pizá, M., et al. (2016). Long-term empirical evidence of ocean warming leading to tropicalization of fish communities, increased herbivory, and loss of kelp. *Proceedings of the National Academy of Sciences of the United States of America*, 113(48), 13791–13796. <https://doi.org/10.1073/pnas.1610725113>
80. Vergés, A., Steinberg, P. D., Hay, M. E., Poore, A. G. B., Campbell, A. H., et al. (2014). The tropicalization of temperate marine ecosystems: Climate-mediated changes in herbivory and community phase shifts. *Proceedings of the Royal Society B: Biological Sciences*, 281(1789), 1–12. <https://doi.org/10.1098/rspb.2014.0846>
81. Wählström, I., Hammar, L., Hume, D., Pålsson, J., Almroth-Rosell, E., et al. (2022). Projected climate change impact on a coastal sea—as significant as all current pressures combined. *Global Change Biology*, 28(17), 5310–5319. <https://doi.org/10.1111/gcb.16312>
82. Walsh, W. A., & Kleiber, P. (2001). Generalized additive model and regression tree analyses of blue shark (*Prionace glauca*) catch rates by the Hawaii-based commercial longline fishery. *Fisheries Research*, 53(2), 115–131. [https://doi.org/10.1016/S0165-7836\(00\)00306-4](https://doi.org/10.1016/S0165-7836(00)00306-4)
83. Walther, G.-R., Post, E., Convey, P., Menzel, A., Parmesan, C., et al. (2002). Ecological responses to recent climate change. *Nature*, 416, 389–395. <https://doi.org/10.1038/416389a>
84. Waycott, M., Duarte, C. M., Carruthers, T. J. B., Orth, R. J., Dennison, W. C., et al. (2009). Accelerating loss of seagrasses across the globe threatens coastal ecosystems. *Proceedings of the National Academy of Sciences of the United States of America*, 106(30), 12377–12381. <https://doi.org/10.1073/pnas.0905620106>
85. Wernberg, T., Bennett, S., Babcock, R. C., de Bettignies, T., Cure, K., et al. (2016). Climate-driven regime shift of a temperate marine ecosystem. *Science*, 353(6295), 169–172. <https://doi.org/10.1126/science.aad8745>
86. Wernberg, T., Coleman, M. A., Bennett, S., Thomsen, M. S., Tuya, F., & Kelaher, B. P. (2018). Genetic diversity and kelp forest vulnerability to climatic stress. *Scientific Reports*, 8(1851), 1–8. <https://doi.org/10.1038/s41598-018-20009-9>
87. Wernberg, T., Russell, B. D., Thomsen, M. S., Gurgel, C. F. D., Bradshaw, C. J. A., et al. (2011). Seaweed communities in retreat from ocean warming. *Current Biology*, 21(21), 1828–1832. <https://doi.org/10.1016/j.cub.2011.09.028>
88. Wood, S. N. (2017). *Generalized additive models: An introduction with R* (2nd ed.). Chapman and Hall/CRC. <https://doi.org/10.1201/9781315370279>

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
89. **Wood, S. N., Li, Z., Shaddick, G., & Augustin, N. H. (2017).** Generalized additive models for gigadata: Modeling the U.K. Black Smoke Network daily data. *Journal of the American Statistical Association*, 112(519), 1199–1210. <https://doi.org/10.1080/01621459.2016.1195744>
90. **Yee, T. W., & Mitchell, N. D. (1991).** Generalized additive models in plant ecology. *Journal of Vegetation Science*, 2(5), 587–602. <https://doi.org/10.2307/3236170>
91. **Zou, D., & Gao, K. (2013).** Thermal acclimation of respiration and photosynthesis in the marine macroalga *Gracilaria lemaneiformis* (Gracilariales, Rhodophyta). *Journal of Phycology*, 49(1), 61–68. <https://doi.org/10.1111/jpy.12009>

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

**TABLES AND FIGURES**

**Table 1:** Number of algae samples from the four macroalgae orders (Bryopsidales, Ceramiales, Dictyotales and Fucales) selected for the entire Canary Islands region (Totals) and for each island of the archipelago [El Hierro (EH), La Palma (LP), La Gomera (LG), Tenerife (TF), Gran Canaria (GC), Lanzarote (LZ) and Fuerteventura (FV)].

| Order        | EH  | LP  | LG  | TF   | GC   | LZ   | FV   | Total |
|--------------|-----|-----|-----|------|------|------|------|-------|
| Bryopsidales | 45  | 50  | 140 | 255  | 880  | 503  | 284  | 2157  |
| Ceramiales   | 117 | 364 | 17  | 680  | 885  | 564  | 1454 | 4081  |
| Dictyotales  | 38  | 122 | 11  | 431  | 278  | 763  | 141  | 1784  |
| Fucales      | 42  | 135 | 39  | 430  | 2190 | 291  | 155  | 3282  |
| Total        | 242 | 671 | 207 | 1796 | 4233 | 2121 | 2034 | 11304 |

**Table 2.** Generalized additive model (GAM) results for four macroalgae orders (Bryopsidales, Ceramiales, Dictyotales, and Fucales) in the Canary Islands, assessing their response to environmental factors such as Salinity, Sea Surface Temperature (SST), Significant Wave Height (SWH), Surface Solar Radiation (SSR), Nitrate (NO<sub>3</sub>), and Phosphate (PO<sub>4</sub>).

| Order        | Significant variables                                 | R <sup>2</sup> (Adj) | Deviance explained |
|--------------|-------------------------------------------------------|----------------------|--------------------|
| Bryopsidales | SST, SWH, Salinity                                    | 0.35                 | 66.8%              |
| Ceramiales   | SST, SSR, SWH, NO <sub>3</sub> , PO <sub>4</sub>      | 0.439                | 61.5%              |
| Dictyotales  | SST, SWH, Salinity, NO <sub>3</sub> , PO <sub>4</sub> | 0.304                | 64.5%              |
| Fucales      | SSR, Salinity                                         | 0.289                | 54.9%              |

**Table 3.** Generalized additive model (GAM) results for four macroalgae orders (Bryopsidales, Ceramiales, Dictyotales, and Fucales) in the seven islands of Canary Islands, assessing their response to environmental factors such as Salinity, Sea Surface Temperature (SST), Significant Wave Height (SWH), Surface Solar Radiation (SSR), Nitrate (NO<sub>3</sub>), and Phosphate (PO<sub>4</sub>).


| Island    | Order        | Significant variables     | R <sup>2</sup><br>(Adj) | Deviance<br>explained |
|-----------|--------------|---------------------------|-------------------------|-----------------------|
| El Hierro | Bryopsidales | -                         | 0.037                   | 16.1%                 |
|           | Ceramiales   | SWH, NO <sub>3</sub>      | 0.76                    | 75.3%                 |
|           | Dictyotales  | SSR                       | 0.601                   | 70.2%                 |
|           | Fucales      | SSR, PO <sub>4</sub>      | 0.296                   | 59.1%                 |
| La Palma  | Bryopsidales | Salinity, NO <sub>3</sub> | 0.654                   | 76.7%                 |
|           | Ceramiales   | SST                       | 0.25                    | 47.3%                 |
|           | Dictyotales  | SSR, SWH                  | 0.491                   | 73.5%                 |
|           | Fucales      | SST, NO <sub>3</sub>      | 0.75                    | 78.5%                 |
| La Gomera | Bryopsidales | Salinity                  | 0.878                   | 81.2%                 |
|           | Ceramiales   | NO <sub>3</sub>           | 0.13                    | 41%                   |
|           | Dictyotales  | SSR                       | 0.522                   | 79.2%                 |
|           | Fucales      | SST, SWH, NO <sub>3</sub> | 0.796                   | 86.1%                 |
| Tenerife  | Bryopsidales | SST, SSR                  | 0.678                   | 60.2%                 |

|               |              |                                        |       |       |
|---------------|--------------|----------------------------------------|-------|-------|
| Gran Canaria  | Ceramial     | SST, NO <sub>3</sub>                   | 0.213 | 38.9% |
|               | Dictyotal    | SSR, NO <sub>3</sub> , PO <sub>4</sub> | 0.664 | 53.9% |
|               | Fucal        | Salinity, PO <sub>4</sub> ,            | 0.182 | 35.6% |
|               | Bryopsidales | SSR, SWH                               | 0.6   | 65.3% |
|               | Ceramiales   | SST, SWH, Salinity, NO <sub>3</sub>    | 0.531 | 48.9% |
| Lanzarote     | Dictyotales  | SST, SWH, Salinity, PO <sub>4</sub>    | 0.748 | 73.9% |
|               | Fucales      | SST, PO <sub>4</sub>                   | 0.6   | 70.7% |
|               | Bryopsidales | SST, SWH, NO <sub>3</sub>              | 0.431 | 62.1% |
|               | Ceramiales   | Salinity, PO <sub>4</sub>              | 0.815 | 68.4% |
|               | Dictyotales  | SST, NO <sub>3</sub>                   | 0.359 | 51.8% |
| Fuerteventura | Fucales      | SST, SWH                               | 0.217 | 55%   |
|               | Bryopsidales | SSR, SWH, Salinity                     | 0.667 | 73.8% |
|               | Ceramiales   | NO <sub>3</sub> , PO <sub>4</sub>      | 0.69  | 62.8% |
|               | Dictyotales  | SST                                    | 0.466 | 52.6% |
|               | Fucales      | SST, Salinity                          | 0.6   | 68.7% |

---


**Table 4.** Generalized additive model (GAM) results for four macroalgae orders (Bryopsidales, Ceramiales, Dictyotales, and Fucales) in the western islands of Canary Islands (El Hierro, Las Palmas, La Gomera, and Tenerife) and the eastern islands of Canary Islands (Gran Canaria, Lanzarote, and Fuerteventura), assessing their response to environmental factors such as Salinity, Sea Surface Temperature (SST), Significant Wave Height (SWH), Surface Solar Radiation (SSR), Nitrate (NO<sub>3</sub>), and Phosphate (PO<sub>4</sub>).

| Island group | Order        | Significant variables                                 | R2 (Adj) | Deviance explained |
|--------------|--------------|-------------------------------------------------------|----------|--------------------|
| Western      | Bryopsidales | SST, SSR, Salinity                                    | 0.763    | 85%                |
|              | Ceramiales   | SST, SWH, NO <sub>3</sub> , PO <sub>4</sub>           | 0.77     | 84.9%              |
|              | Dictyotales  | SST, Salinity, PO <sub>4</sub>                        | 0.511    | 77.5%              |
|              | Fucales      | SST, SSR                                              | 0.79     | 81.6%              |
| Eastern      | Bryopsidales | SWH, SSR, Salinity                                    | 0.852    | 92.5%              |
|              | Ceramiales   | SST, SSR, NO <sub>3</sub>                             | 0.782    | 90.5%              |
|              | Dictyotales  | SSR, SWH, Salinity, NO <sub>3</sub> , PO <sub>4</sub> | 0.733    | 82.7%              |
|              | Fucales      | SST, SSR, SWH, PO <sub>4</sub>                        | 0.653    | 73.5%              |




**Figure 1.** Bathymetric map of the Northeast Atlantic Ocean and the Canary Islands: top panel displays the bathymetry of the Northeast Atlantic Ocean, with contour lines highlighting major underwater structures; Bottom panel provides a detailed view of the Canary Islands region, illustrating bathymetric gradients and the main islands of the archipelago. Colors represent depth, ranging from deep areas (dark blue) to emerged land (gray). Labels indicate the names of the main islands (NOAA, 2009).


1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



**Figure 2.** Histograms of algae records: (a) distribution across the Canary Islands (El Hierro, La Palma, La Gomera, Tenerife, Gran Canaria, Lanzarote, and Fuerteventura); (b) grouped by island region (Western and Eastern); and (c) by selected orders (Bryopsidales, Ceramiales, Dictyotales, and Fucales).



**Figure. 3.** Frequency diagrams of annual records for the four studied macroalgae orders over a 28-year period (1992–2020): Bryopsidales (Red), Ceramiales (Blue), Dictyotales (Green), and Fucales (Brown).



**Figure 4.** Principal Component Analysis (PCA) of presence-absence records for algae species (order Bryopsidales, Ceramiales, Dictyotales, and Fuciales) and environmental time series data collected over 18 years (1992–2020). Environmental variables include nutrient surface concentrations ( $\mu\text{g}/\text{km}^2$ ) for  $\text{NH}_4$ ,  $\text{NO}_3$ , and  $\text{PO}_4$ ; as well as environmental parameters such as temperature ( $^{\circ}\text{C}$ ), seawater salinity, sea surface radiation (SSR), sea surface height (SSH), depth, significant wave height (SWH), and sampling year.




Figure 5. Log-transformed coefficient estimates from GAM models fitted to predictor variables influencing the distribution of different orders of marine algae in different perspectives of the Canary Islands, comparing the effects of environmental variables across different algal orders and regions. **Top Panel:** Effects of environmental variables on algal orders (Bryopsidales, Ceramiales, Dictyotales, and Fucales) across parametric and smooth terms. Term types are represented by different shapes (diamonds for parametric, triangles for smooth). **Middle Panel:** Island-level differences in environmental variable effects on algal orders. Each point represents the estimated coefficient for a given environmental variable across different islands. **Bottom Panel:** Regional differences in the effects of environmental variables on algal orders. Lines and points distinguish between estimates for Western Islands and Eastern Islands, with orders differentiated by color and region-order combinations indicated in the legend. Dashed vertical lines at -1 and 1 indicate thresholds for interpretation of the Log10(Coefficient Estimate). Stars indicate the significance level of terms (\* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$ ).