Loud the Sound: The Brazilian Butt Lift Has Come to Stay, But the Epidemiological Evidence Says It May Be the Deadliest Cosmetic Surgery of Our Time

Olumide M. Arigbede, DrPH, MPH
Chair, The Olumide Arigbede Data Clinic
Published: April 2026

"In God we trust; all others must bring data."
W. Edwards Deming

The Paradox of Popularity and Peril

There are procedures in medicine that grow quietly, earning their place through decades of accumulated evidence. The Brazilian Butt Lift is not one of them. Its ascent has been volcanic. In just 12 years, global procedure volume surged from approximately 14,705 in 2012 to over 600,000 in 2024, representing an estimated 4,000% increase, according to data compiled by the International Society of Aesthetic Plastic Surgery (ISAPS).[7] In the United States alone, the American Society of Plastic Surgeons (ASPS) documented a rise from roughly 8,500 procedures in 2012 to more than 28,000 by 2019, with the American Society of Aesthetic Plastic Surgeons (ASAPS) reporting 61,387 buttock augmentations by its members in 2021 (a 37% single year increase).[11] These data capture only cases performed by board-certified plastic surgeons. The actual global volume, accounting for procedures by non-certified practitioners, medical tourism clinics, and unregulated facilities, is almost certainly far higher.

Yet behind this extraordinary demand lies a mortality signal that no epidemiologist can ignore. The BBL has earned a distinction no cosmetic procedure wants: it carries the highest procedure-specific mortality rate of any aesthetic surgery ever tracked.

The Mortality Epidemiology: From Signal to Certainty

The mortality alarm was sounded formally in 2017, when the Aesthetic Surgery Education and Research Foundation (ASERF) published results from a landmark survey of plastic surgeons. Mofid and colleagues, surveying 692 respondents covering 198,857 cases, estimated the BBL mortality rate at approximately 1 in 3,000 (33.3 per 100,000 procedures).[2] To place this in an epidemiological context, the case fatality rate for traditional liposuction ranges from 2.6 to 20.6 per 100,000 procedures. The early BBL estimate was therefore between 1.6 and 12.8 times the mortality rate of liposuction, and roughly 33 times the background mortality rate for cosmetic surgery overall (approximately 1.0 per 100,000). ASPS itself described the BBL mortality rate as three times higher than any other cosmetic procedure and 16 times higher than the average for all procedures performed in accredited surgical centers.[1]

Subsequent studies refined these estimates. Rios and Gupta (2020), using improved survey methodology, reported a revised mortality rate of approximately 1 in 14,952 (6.7 per 100,000),[3] a meaningful reduction attributed partly to updated surgical techniques. A 2022 survey from the World Association of Gluteal Surgeons (WAGS) calculated a rate of 1 in 23,818 among its multispecialty membership. These revised figures, while encouraging, still place the BBL above tummy tuck mortality (estimated at 1 in 13,147 by the same 2020 survey)[3] and far above the baseline for elective cosmetic procedures.

The causal mechanism is well established. Nearly all BBL deaths result from pulmonary fat embolism (PFE), a mechanical obstruction of the pulmonary vasculature caused by adipose tissue entering the venous circulation during gluteal fat injection. PFE is not a gradual complication; it is typically catastrophic, with death occurring within the first 24 hours after surgery. Autopsy studies from South Florida confirmed that in every fatal case examined, fat grafts had been injected into the gluteal musculature at multiple levels, breaching the vascular planes that lie deep within the muscle.[5]

Geographic Clustering: South Florida and Beyond

The geographic distribution of BBL mortality is far from uniform, and this heterogeneity itself tells an epidemiological story. South Florida has emerged as a global outlier. Between 2010 and 2022, the region documented 25 BBL related fat embolism deaths.[5] What is particularly alarming is that 14 of these 25 deaths (56%) occurred *after* publication of the ASERF 2018 safety guidelines and the Florida Board of Medicine's 2019 "subcutaneous only" rule. The expected mortality, applying the most favorable published rates to an estimated 20,000 annual BBLs in the Miami-Dade area, would be one or perhaps two deaths per year. Instead, eight women died from BBL related fat embolism in South Florida in 2021 alone.[5] The observed-to-expected ratio points unmistakably toward systemic failures: high-volume, low-cost clinics operating under intense production pressure, with surgeons sometimes performing procedures in under 90 minutes and working on commission structures that incentivize speed over safety. Some clinics in Florida advertise BBLs for as little as $2,900, compared to a national range of $8,000 to $20,000.

The Medical Tourism Amplifier

The epidemiological picture becomes bleaker when one considers the medical tourism pipeline. A 2024 CDC report published in the *Morbidity and Mortality Weekly Report* documented 93 deaths of U.S. citizens following cosmetic surgery in the Dominican Republic between 2009 and 2022 (Hudson et al., 2024). Of these, 92% involved gluteal fat transfer. The death rate accelerated sharply over the study period, rising from 4.1 deaths per year to 13 per year, peaking at 17 deaths in 2020. All but one decedent was female; the median age was 40 years; and the mean BMI was 32 kg/m^2. Autopsy records revealed that 90% of the deaths were attributable to embolic phenomena. Venous thromboembolism comorbidities were present in 92% of cases.[6]

These numbers represent a single country and capture only American deaths. The true global toll, encompassing patients from the United Kingdom, Europe, and across Latin America who travel to Colombia, Mexico, Turkey, and the Dominican Republic for discounted packages, is almost certainly a substantial multiple of these counts. The

British Association of Aesthetic Plastic Surgeons (BAAPS) has taken the most drastic regulatory position of any national body, advising its members not to perform BBL surgery at all. The U.K. Foreign Office has confirmed awareness of more than 25 British nationals who died in Turkey alone following medical procedures since January 2019. NHS England has publicly warned about the burden of treating returning patients with post-surgical complications, with hospitalization costs for such cases ranging from $26,000 to $154,000 per admission.[9]

A Visual Epidemiological Summary

Figure 1 presents two complementary panels. Panel A displays the observed global trajectory of BBL procedure volume from 2012 through 2024, with moderate growth projections (8% to 12% annual increase) extending through 2030. If current trends persist, global annual volume may approach or exceed 1 million procedures by the end of this decade. Panel B compares mortality rates per 100,000 procedures across the spectrum of cosmetic surgeries and across different BBL injection techniques. The contrast between the early intramuscular injection mortality rate (33.3 per 100,000) and the subcutaneous-only technique (0.0 per 100,000 in 12,800 observed cases; Cuzalina et al., 2024)[2,4,7] illustrates both the scale of the original hazard and the potential of technique standardization to eliminate it.



Figure 1. Global BBL procedure volume (Panel A, observed and projected) and comparative mortality rates across cosmetic procedures (Panel B).

Technique, Regulation, and the Path Toward Harm Reduction

The most consequential finding in the recent literature is the dramatic effect of the injection plane on survival. Cuzalina and colleagues (2024), surveying members of the American Board of Cosmetic Surgeons across two consecutive three-year windows, reported a mortality rate of zero (0 out of 12,800 cases) when 96% of surgeons injected

fat exclusively above the muscle. In contrast, during the earlier window, when 39% of 22,800 cases involved intramuscular injection, the mortality rate was 1 in 11,400. This is the kind of evidence that, in infectious disease epidemiology, would prompt an immediate change in clinical guidelines. In the BBL context, it points to a clear, modifiable risk factor: depth of injection.[4]

Ultrasound-guided injection, which allows real-time visualization of cannula position relative to the gluteal fascia and musculature, has emerged as a promising technological solution. A systematic review published in 2024, analyzing 6,235 patients who received ultrasound-guided gluteal fat grafting, found no reported deaths (0.00 per 100, 95% CI: 0.00 to 0.00) and no fat embolism events. Minor complications occurred at a pooled rate of 6.32 per 100 (95% CI: 3.23 to 10.27), with seroma at 2.94 per 100 and infection at 0.23 per 100. These data are preliminary, drawn from a small number of studies, but they represent the strongest safety signal in the BBL literature to date.[10,12]

Regulatory responses have been uneven. The Florida Board of Medicine issued an emergency order in June 2022 limiting BBL surgeries to three per day per surgeon and mandating intraoperative ultrasound. Yet a survey of 178 board-certified plastic surgeons by Finkelstein and colleagues (published in 2025 in the *Public Health Post*) found that 60% opposed mandatory ultrasound regulations, and more than half significantly underestimated the BBL mortality rate. Only 20% of respondents could accurately estimate the current fatality figures. This knowledge gap among practitioners is itself an epidemiological risk factor, because a surgeon who underestimates the hazard is less likely to adopt protective techniques.[8]

The Population at Risk: Demographics and Disparities

The typical BBL patient is a woman aged 21 to 36, and this group accounts for approximately 70% of all cases globally. African American, Latina, and Afro-Brazilian women have historically driven early demand, reflecting the cultural valorization of curvaceous body ideals. More recently, however, ISAPS data from 2024 show a 50% increase in BBL uptake among white women in the United States and European Union over a five-year span. Male patients now represent a growing share, with a 12% year-on-year increase reported. The age ceiling is also expanding, with patients aged 45 and older increasingly pursuing the procedure.[7]

A 2023 study by the British Association of Aesthetic Plastic Surgeons found that one in three female BBL patients under 25 cited social media as the primary motivator for seeking surgery. The #BBL hashtag generates millions of views daily on Instagram and TikTok. Google Trends data show that searches for “BBL surgery” increased 700% between 2015 and 2022. This social media-driven demand pipeline has fundamentally altered the risk landscape. Patients arrive at clinics with reference photographs from influencers, often with unrealistic expectations, and many travel to unregulated facilities abroad to obtain the procedure at a fraction of the domestic cost.

An Epidemiologist’s Reckoning

We (epidemiologists) are trained to look at numbers and see people. Behind every “1 in 3,000” is a woman who walked into an operating room healthy and never walked out.

Behind every upward revision of the denominator is a silent cohort of unreported cases in countries where surgical registries do not exist or where deaths are attributed to other causes.

The BBL is not going away. Demand projections suggest the procedure could surpass one million cases annually by 2030. What must change is the framework within which those procedures are performed. The evidence base, though imperfect, points toward a clear hierarchy of interventions: mandatory subcutaneous-only injection protocols, ultrasound-guided visualization for all cases, accreditation requirements for surgical facilities, enforceable limits on daily case volume per surgeon, and, perhaps most critically, international harmonization of safety standards for medical tourism destinations.

In epidemiology, we distinguish between hazards that can be eliminated and those that can only be mitigated. The BBL mortality hazard falls into the latter category, but the difference between mortality rates of 33 per 100,000 and 0 per 100,000 (as demonstrated in subcutaneous-only, ultrasound-guided cohorts)[4,12] is not theoretical. It is achievable. The question is whether the regulatory and professional will exist to impose these standards universally, or whether market forces, medical tourism arbitrage, and the algorithmic amplification of body ideals will continue to expose hundreds of thousands of patients to avoidable risk.

Data brought us to this moment of understanding. Only action will carry us beyond it.

Selected References

1. American Society of Plastic Surgeons (ASPS). Plastic surgery societies issue urgent warning about the risks associated with Brazilian butt lifts. Press release. August 6, 2018.

2. Mofid MM, Teitelbaum S, Suissa D, et al. Report on mortality from gluteal fat grafting: recommendations from the ASERF Task Force. Aesthet Surg J. 2017;37(7):796–806.

3. Rios L, Gupta V. Improvement in Brazilian butt lift (BBL) safety with the current recommendations from ASERF, ASAPS, and ISAPS. Aesthet Surg J. 2020;40(8):864–870.

4. Cuzalina A, Mostofi P, Hah W. Gluteal fat grafting technique and mortality update among surveyed ABCS surgeons. Am J Cosmet Surg. 2024;41(1):1–9.

5. Pazmino P, Garcia O. Brazilian butt lift–associated mortality: the South Florida experience. Aesthet Surg J. 2023;43(2):162–178.

6. Hudson M, Matos JA, Alvarez B, et al. Deaths of U.S. citizens undergoing cosmetic surgery, Dominican Republic, 2009–2022. MMWR Morb Mortal Wkly Rep. 2024;73(3):62–65.

7. International Society of Aesthetic Plastic Surgery (ISAPS). Global Survey 2024: Full Report. Published June 19, 2025.

8. Finkelstein ER, et al. If BBL safety measures exist, why are mortality rates so high? Public Health Post. May 21, 2025.

9. NHS England. NHS warns of 'bargain basement' BBLs in run up to Christmas. December 2024.

10. Cansancao AL, Conde Green A, Vidigal RA, et al. Real time ultrasound assisted gluteal fat grafting. Plast Reconstr Surg. 2018;142(2):372–376.

11. American Society of Plastic Surgeons (ASPS). 2024 Plastic Surgery Statistics Report. Published 2025.

12. Delobaux A. Systematic review: safety of ultrasound guided gluteal fat grafting. Analysis of 6,235 patients. Published 2024.

Disclaimer: *This commentary is intended for educational and scholarly purposes. The author holds no financial interest in any cosmetic surgery product, service, or facility. Mortality estimates cited reflect published survey data and may undercount actual fatalities due to underreporting, particularly in unregulated settings and medical tourism destinations.*