

An Algebraic Framework for the Game of Go

0. Acknowledgment

This essay is a revised version of an earlier draft. The author gratefully acknowledges detailed reviews (Go_v0-review_1_Gem.md, Go_v1-review_2_Cla.md, Go_v1-review_3_Gem.md, Go_v1-review_4_Cla.md, Go_v2-review_5_gpt.md, Go_v3-review_6_gpt.md), whose suggestions have shaped the framework presented here: the state space is now modeled as a finite DAG under positional superko (with history made explicit), liberties are defined via the coboundary operator, life and death is formalized through the Knaster-Tarski fixed point of Benson's monotone operator, and endgames are analyzed through combinatorial game theory (with the independence hypothesis explicitly stated).

1. Introduction

The ancient board game of Go, played on a grid of $n \times n$ intersections (typically $n = 19$), has long been recognized as a fertile ground for mathematical inquiry. While much of the existing literature addresses Go from the standpoint of combinatorial game theory, algorithmic complexity, and statistical evaluation, relatively few treatments have explored its underlying algebraic structure. This essay proposes a systematic algebraic formulation of Go that casts the game in terms of posets, chain complexes, Boolean lattices, and the theory of surreal games. We shall demonstrate that the board, the set of legal positions, and the dynamics of play admit natural structures that are amenable to the tools of abstract algebra. The presentation is aimed at readers comfortable with the language of groups, rings, modules, and categories; no prior knowledge of Go is assumed, though familiarity with the basic rules will aid intuition.

2. The State Space of Go

Let $B_n = \{(i,j) : 1 \leq i,j \leq n\}$ denote the set of intersections of an $n \times n$ board. A **state** (or board configuration) is a function

$$s : B_n \longrightarrow \{0, 1, 2\},$$

where 0 signifies an empty intersection, 1 a black stone, and 2 a white stone. The set of all such functions,

$$S_n = \{0, 1, 2\}^{B_n} \cong \mathbb{Z}_3^{n^2},$$

has cardinality 3^{n^2} . A state s is **legal** if every monochromatic connected component (a "group") of stones has at least one adjacent empty intersection (a "liberty") or, equivalently, if the position could arise from the empty board by a sequence of legal moves. Let $P_n \subseteq S_n$ denote the set of legal positions.

A **move** consists of placing a stone on an empty intersection, followed by the removal of any opponent groups that have thereby lost their last liberty (a **capture**). Let M_n denote the set of legal moves, and define the **legal-move relation**

$$R \subseteq P_n \times P_n, \quad (p, q) \in R \iff \text{there exists a legal move } p \rightarrow q.$$

The pair $G_n = (P_n, R)$ is a directed graph whose vertices are legal positions and whose edges are legal single moves.

Go is played under the **Positional Superko rule**: a move from p to q is illegal if the position q has already appeared earlier in the current game. To make this history dependence precise, we define a **game state** as a pair (p, h) , where $p \in P_n$ is the current board position and $h \in P_n^*$ is the finite sequence of positions visited so far (the game history). Let

$$\mathcal{G}_n = \{(p, h) : p \in P_n, h \in P_n^*, p \notin h\}$$

be the set of game states, and define the transition relation T on G_n by

$$(p, h) \xrightarrow{T} (q, h \cdot p) \iff (p, q) \in R \text{ and } q \notin h.$$

Proposition 2.1. *The graph T on G_n is a finite directed acyclic graph (DAG). Therefore the reachability relation $\leq_T$ on G_n (defined by $(p,h) \leq_T (q,h')$ iff (q,h') is reachable from (p,h) by a directed path) is a partial order.*

Proof. Each step of T appends a position to the history, so the length of the history is a strictly increasing function on T -paths. Since there are only finitely many positions, no infinite paths exist, and any cycle would force a repeated history length — impossible. Hence T is acyclic, and reachability in a finite acyclic graph is a partial order. $\square$

For a game state (p,h) , let $D(p,h)$ denote the maximum length of a directed T -path starting at (p,h) . Since T is a finite DAG, D is well-defined and finite.

Corollary 2.2. *Every game of Go starting from a legal position p terminates after at most $D(p,\varepsilon)$ moves, and in particular after at most $3^{n^2} - 1$ moves. More precisely, for any current state (p,h) ,*

$$D(p,h) \leq |\mathcal{P}_n| - |h| - 1 \leq 3^{n^2} - |h| - 1,$$

since each move must visit a position not yet appearing in the history h .

This is the load-bearing finiteness fact: the Superko rule converts the naive state graph into a poset, and the finiteness of the poset forces termination of all maximal chains. It is precisely this property that makes the recursive definitions of later sections well-founded.

3. Topological Go: Liberties and Eyes

Model the board as the graph $G = (V, E)$ where $V = B_n$ and E consists of the $2n(n-1)$ orthogonal adjacencies. Let $C_0(G; \mathbb{Z}_2)$ and $C_1(G; \mathbb{Z}_2)$ be the usual chain groups over $\mathbb{Z}_2$, with boundary operator $\partial_1 : C_1 \rightarrow C_0$. The dual cochain groups C^0, C^1 carry the coboundary $\delta_0 : C^0 \rightarrow C^1$, where

$$\langle \delta_0 \varphi, e \rangle = \varphi(\partial_1 e) = \varphi(u) + \varphi(v) \quad \text{for } e = \{u, v\}.$$

A **stone group** c is a connected component of monochromatic stones on the board. Represent c by its indicator cochain $1_c \in C^0(G; \mathbb{Z}_2)$, which is 1 on vertices of c and 0 elsewhere. Then $\delta_0(1_c) \in C^1(G; \mathbb{Z}_2)$ is the indicator of the **edge cut** separating c from its complement.

Let $V_\square$ denote the set of empty intersections on the board. Define the **liberty set** of c by

$$L(c) = \{v \in V_\square : \exists u \in c \text{ with } \{u, v\} \in \text{supp}(\delta_0(1_c))\}.$$

In words, $L(c)$ is the set of empty vertices incident to edges in the coboundary of c .

Proposition 3.1. *A stone group c is captured (i.e., has zero liberties) if and only if $\delta_0(1_c)$ is supported entirely on edges whose endpoints both lie in $V \setminus V_\square$ (the occupied vertices).*

Proof. By definition, c is captured precisely when every neighbor of every vertex of c is occupied, i.e., when no edge in $\text{supp}(\delta_0(1_c))$ is incident to a vertex of $V_\square$. This is exactly the stated condition on the support of $\delta_0(1_c)$. $\square$

An **eye** is an empty intersection $v \in V_\square$ whose four orthogonal neighbors (when they exist) are all stones of the same color, forming a cycle in the induced subgraph on those neighbors. Equivalently, v is an eye of a stone group c when $v \in V_\square$ and $\partial_1^* 1_{\{v\}} = 0$ in $C_1(G_c; \mathbb{Z}_2)$, where G_c is the subgraph induced by $c \cup \{v\}$. In the language of relative homology, nontrivial classes in $H_1(G_c, c; \mathbb{Z}_2)$ provide a topological certificate of such enclosure.

Remark 3.2 (on false eyes). Orthogonal enclosure of an empty point — detectable purely homologically — is a necessary but not sufficient condition for a true eye: a “false eye” may be surrounded orthogonally yet its controlling diagonal points lie outside c , allowing the opponent to play there and destroy the enclosure. The diagonal/ko-dependent part of eye-trueness is therefore a dynamic property of the game tree, not a static topological invariant of a single position. We accordingly record only the topological *candidate* eyes via H_1 , leaving the true/false distinction to the lattice-theoretic analysis of Section 4.

4. Lattice Theory: Life and Death

Fix a board position and consider the Boolean lattice

$$L = \mathcal{P}(\text{Black stones on the board})$$

ordered by inclusion. Define a monotone map $f : L \rightarrow L$ (Benson's operator) as follows: for $X \in L$, $f(X)$ is the subset of X consisting of those stones that belong to groups whose liberties lie entirely within regions that are *enclosed* by X in the topological sense of Section 3 (i.e., whose incident empty vertices are eyes or interior points of X). Because f is order-preserving on a finite complete lattice, the **Knaster-Tarski theorem** applies:

Theorem 4.1. *The operator f possesses a greatest fixed point*

$$\text{gfp}(f) = \bigvee \{ X \in L : f(X) \supseteq X \},$$

and this greatest fixed point is exactly the set of Black stones that are unconditionally alive (pass-alive) on the board.

Proof. The set of f -invariant elements is closed under arbitrary joins in a complete lattice, so the displayed supremum is itself a fixed point and dominates every other fixed point. By construction, a stone belongs to $\text{gfp}(f)$ iff it lies in a group whose liberties are permanently sealed off from the opponent — the precise definition of unconditionally alive. $\square$

Lemma 4.2 (bridge to topology). *If a Black stone group c possesses two disjoint candidate eyes that remain enclosed throughout Benson's greatest-fixed-point iteration, then $c \subseteq \text{gfp}(f)$; in particular, c is unconditionally alive.*

Proof. Let e_1, e_2 be the two disjoint candidate eyes of c . By definition of candidate eye, each e_i is orthogonally surrounded by stones of c , so any liberty of c incident to e_i lies in a region enclosed by c . Since the eyes are disjoint, any opponent move can fill at most one of them; the other remains an empty point orthogonal-neighbor to c . Consequently, after any opponent move, c still retains at least one liberty, so c is never captured. This means that for every $X \in L$ with $c \subseteq X$, we have $f(X) \supseteq c$. In particular, $f(c) \supseteq c$, so c is a post-fixed point of f . By the Knaster-Tarski theorem, every post-fixed point is contained in the greatest fixed point, i.e., $c \subseteq \text{gfp}(f)$. $\square$

Thus the topological candidate eyes of Section 3 feed directly into the lattice fixed point that decides life and death — the two sections are not merely adjacent but genuinely interlocking.

Remark 4.3. The converse of Lemma 4.2 fails: a group may be unconditionally alive without possessing two disjoint eyes. The lattice formulation is strictly more general than the classical “two eyes” heuristic, which is why we treat the eyes as a sufficient but not necessary certificate of life.

5. Combinatorial Game Theory: Endgames

In the endgame, the board typically fractures into isolated, independent skirmishes. Following Conway, Berlekamp, and Wolfe, each local region R_i defines a game g_i in the sense of combinatorial game theory, and the global position decomposes as a **direct sum**:

$$G = g_1 \oplus g_2 \oplus \cdots \oplus g_k.$$

This decomposition is valid precisely when the regions are **independent**: a move in R_i does not affect the set of legal moves in any R_j ($i \neq j$). In practice, this independence holds for most endgame positions, but one must be careful with regions that interact via ko threats, sente/gote dependencies, or shared liberties — in those cases the sum must be refined to a more general CGT structure (e.g., a sum of games with a “temperature” parameter).

The collection of such sums forms a partially ordered abelian group, where the order $G > 0$ signifies that the next player (Black, by convention) has a winning strategy. The value

of a Go position is not merely an integer but a **surreal number**, possibly enriched with infinitesimal components ("tiny", "switch", "fuzz") that capture subtle endgame nuances. Evaluating a board reduces to computing the sum of these algebraic values — a task that blends algebra with strategy in the most direct way.

6. Conclusion

We have presented an algebraic framework for Go that unifies several branches of abstract algebra: the game tree as a finite DAG (hence a poset) via positional superko, liberties and eyes as coboundary and homology invariants, life/death as the Knaster-Tarski fixed point of Benson's monotone operator, and endgames as direct sums in the group of surreal games. Each structure captures a different facet of the game, and together they form a coherent mathematical portrait of Go. We hope this essay stimulates further work at the intersection of algebra and this venerable game, turning strategic intuition into theorem and computation.

Open problems. (i) Can one strengthen Lemma 4.2 to a full characterization of true eyes as those candidate eyes that survive the greatest-fixed-point iteration? (ii) Does the DAG height $D(p)$ of Section 2 admit a closed form in terms of local topological invariants? (iii) How do the irreducible representations of the board's dihedral symmetry group D_4 decompose the space of endgame values?