

Why column space?

Mitaxi Mehta: Lecture 10

- Three basic vector spaces associated with a matrix are defined as,

- Three basic vector spaces associated with a matrix are defined as,
Column space $C(A)$: contains all linear superpositions of column vectors of A .

- Three basic vector spaces associated with a matrix are defined as,
Column space $C(A)$: contains all linear superpositions of column vectors of A .
Row space $R(A)$: contains all linear superpositions of row vectors of A .

- Three basic vector spaces associated with a matrix are defined as,
Column space $C(A)$: contains all linear superpositions of column vectors of A .
Row space $R(A)$: contains all linear superpositions of row vectors of A .
Null space $N(A)$: contains all linear superpositions of Null vectors of A .

- Three basic vector spaces associated with a matrix are defined as,
Column space $C(A)$: contains all linear superpositions of column vectors of A .
Row space $R(A)$: contains all linear superpositions of row vectors of A .
Null space $N(A)$: contains all linear superpositions of Null vectors of A .
- A natural question is, why should one study $C(A)$?

- Three basic vector spaces associated with a matrix are defined as,
 Column space $C(A)$: contains all linear superpositions of column vectors of A .
 Row space $R(A)$: contains all linear superpositions of row vectors of A .
 Null space $N(A)$: contains all linear superpositions of Null vectors of A .
- A natural question is, why should one study $C(A)$?

$$Ax \uparrow = x_1 C_1 \uparrow + x_2 C_2 \uparrow + \dots x_n C_n \uparrow = b \uparrow$$

(where C_i are column vectors of A and x_j are components of the vector $x \uparrow$) is solvable only if the vector $b \uparrow \in C(A)$.

Make a fat and short person look slim and tall

- How would you make a fat and short person look slim and tall in a photograph, provided you have position vectors for the outline of the person ?

Make a fat and short person look slim and tall

- How would you make a fat and short person look slim and tall in a photograph, provided you have position vectors for the outline of the person ?
- Drashti: Use the matrix,

$$\begin{pmatrix} 100 & 2 \\ 98 & 0 \end{pmatrix}$$

Make a fat and short person look slim and tall

- How would you make a fat and short person look slim and tall in a photograph, provided you have position vectors for the outline of the person ?
- Drashti: Use the matrix,

$$\begin{pmatrix} 100 & 2 \\ 98 & 0 \end{pmatrix}$$

- If the top of the head of the person were at (x, y) after transformation it would be

$$\begin{pmatrix} 100 & 2 \\ 98 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 100x + 2y \\ 98x \end{pmatrix}$$

This would not work.

- Dhruv Parikh: Use the matrix

$$\begin{pmatrix} .5 & 0 \\ 0 & 2 \end{pmatrix}$$

- Dhruv Parikh: Use the matrix

$$\begin{pmatrix} .5 & 0 \\ 0 & 2 \end{pmatrix}$$

- This would make the height double but also shift the location.

- How can one store a black and white image (with graylevels) in a matrix ?

- How can one store a black and white image (with graylevels) in a matrix ?
- Jainam K: To store the image of the line $x = y$, set the values on the matrix diagonal as 1.

- How can one store a black and white image (with graylevels) in a matrix ?
- Jainam K: To store the image of the line $x = y$, set the values on the matrix diagonal as 1.
- What about graylevels?

- How can one store a black and white image (with graylevels) in a matrix ?
- Jainam K: To store the image of the line $x = y$, set the values on the matrix diagonal as 1.
- What about graylevels? use values between 0 and 1 to show different graylevels.

- How can one store a black and white image (with graylevels) in a matrix ?
- Jainam K: To store the image of the line $x = y$, set the values on the matrix diagonal as 1.
- What about graylevels? use values between 0 and 1 to show different graylevels.
- How can one store colored images in a matrix ?, Read up before the next class.