

LU decomposition

Mitaxi Mehta: Lecture 8

- While multiplying a matrix with a number, the number gets multiplied to every element of the matrix. For example,

$$3 \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} = \begin{pmatrix} 3a_{11} & 3a_{12} \\ 3a_{21} & 3a_{22} \end{pmatrix}$$

- While multiplying a matrix with a number, the number gets multiplied to every element of the matrix. For example,

$$3 \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} = \begin{pmatrix} 3a_{11} & 3a_{12} \\ 3a_{21} & 3a_{22} \end{pmatrix}$$

- This is different from multiplying a determinant with a number where the number distributes only to a single row.

- While multiplying a matrix with a number, the number gets multiplied to every element of the matrix. For example,

$$3 \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} = \begin{pmatrix} 3a_{11} & 3a_{12} \\ 3a_{21} & 3a_{22} \end{pmatrix}$$

- This is different from multiplying a determinant with a number where the number distributes only to a single row.

$$3 \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} = \begin{vmatrix} 3a_{11} & 3a_{12} \\ a_{21} & a_{22} \end{vmatrix}$$

- To understand this distribution law, consider a major usage of matrices, solving simultaneous linear equations

$$Ax \uparrow = b \uparrow$$

- To understand this distribution law, consider a major usage of matrices, solving simultaneous linear equations

$$Ax \uparrow = b \uparrow$$

- If one wants to multiply both the sides by a number, in the RHS vector the number will distribute to all the elements.

- To understand this distribution law, consider a major usage of matrices, solving simultaneous linear equations

$$Ax = b$$

- If one wants to multiply both the sides by a number, in the RHS vector the number will distribute to all the elements.
- To keep the corresponding line equations the same each coefficient on the LHS also is required to scaled by the same amount.

- To understand this distribution law, consider a major usage of matrices, solving simultaneous linear equations
 $Ax \uparrow = b \uparrow$
- If one wants to multiply both the sides by a number, in the RHS vector the number will distribute to all the elements.
- To keep the corresponding line equations the same each coefficient on the LHS also is required to scaled by the same amount.
- This explains why the multiplying number gets distributed to all elements in the matrix.

- To understand this distribution law, consider a major usage of matrices, solving simultaneous linear equations
 $Ax \uparrow = b \uparrow$
- If one wants to multiply both the sides by a number, in the RHS vector the number will distribute to all the elements.
- To keep the corresponding line equations the same each coefficient on the LHS also is required to scaled by the same amount.
- This explains why the multiplying number gets distributed to all elements in the matrix.
- One may also consider the column picture of the equation and apply the same reasoning.

Column space

- Students are required to refresh basic matrix property from section 1.4 of the reference book by Strang.

Column space

- Students are required to refresh basic matrix property from section 1.4 of the reference book by Strang.
- Note that interchanging of rows of a matrix is accomplished by matrix multiplication as follows,

$$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} = \begin{pmatrix} a_{21} & a_{22} \\ a_{11} & a_{12} \end{pmatrix}$$

Column space

- Students are required to refresh basic matrix property from section 1.4 of the reference book by Strang.
- Note that interchanging of rows of a matrix is accomplished by matrix multiplication as follows,

$$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} = \begin{pmatrix} a_{21} & a_{22} \\ a_{11} & a_{12} \end{pmatrix}$$

- Work out the matrices that can be used for row exchange in higher dimensions.

Column space

- Students are required to refresh basic matrix property from section 1.4 of the reference book by Strang.
- Note that interchanging of rows of a matrix is accomplished by matrix multiplication as follows,

$$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} = \begin{pmatrix} a_{21} & a_{22} \\ a_{11} & a_{12} \end{pmatrix}$$

- Work out the matrices that can be used for row exchange in higher dimensions.
- Satisfy yourself that matrix products are non-commutative.

- Linear equations are particularly easy to solve if the corresponding matrix is in upper or lower triangular form (entries below or above diagonal are zero). For example, solve,

- Linear equations are particularly easy to solve if the corresponding matrix is in upper or lower triangular form (entries below or above diagonal are zero). For example, solve,

$$\begin{pmatrix} 1 & 2 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 3 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 3 \\ 2 \\ 3 \end{pmatrix}$$

- Linear equations are particularly easy to solve if the corresponding matrix is in upper or lower triangular form (entries below or above diagonal are zero). For example, solve,

$$\begin{pmatrix} 1 & 2 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 3 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 3 \\ 2 \\ 3 \end{pmatrix}$$

- Often, it is of interest to decompose a matrix in upper and lower triangular form like $A = LU$

- We shall do the LU decomposition of a matrix using an example, for more details consult section 1.5 of the reference book.

- We shall do the LU decomposition of a matrix using an example, for more details consult section 1.5 of the reference book.
- Consider the following matrix equation,

$$\begin{pmatrix} 2 & 1 & 1 \\ 4 & -6 & 0 \\ -2 & 7 & 2 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 5 \\ -2 \\ 9 \end{pmatrix}$$

- The first step is to make the (2,1) entry zero. This is accomplished by the row operation $R_2 = R_2 - 2R_1$

- The first step is to make the (2,1) entry zero. This is accomplished by the row operation $R_2 = R_2 - 2R_1$
- The matrix that accomplishes this task on multiplication is,

$$E = \begin{pmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

- The first step is to make the (2,1) entry zero. This is accomplished by the row operation $R_2 = R_2 - 2R_1$
- The matrix that accomplishes this task on multiplication is,

$$E = \begin{pmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

- Notice that this is essentially the identity matrix with the entry (2,1) being the multiplying factor.

- The general rule for the row operation $R_m = R_m - cR_k$, replace the (k, m) entry of the identity matrix with $-c$.

- The general rule for the row operation $R_m = R_m - cR_k$, replace the (k, m) entry of the identity matrix with $-c$.
- Check that the following sequence of matrix multiplication brings the matrix in an upper triangular form. $GFEA = U$

- The general rule for the row operation $R_m = R_m - cR_k$, replace the (k, m) entry of the identity matrix with $-c$.
- Check that the following sequence of matrix multiplication brings the matrix in an upper triangular form. $GFEA = U$

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 2 & 1 & 1 \\ 4 & -6 & 0 \\ -2 & 7 & 2 \end{pmatrix}$$

$$= \begin{pmatrix} 2 & 1 & 1 \\ 0 & -8 & 2 \\ 0 & 0 & 1 \end{pmatrix}$$

- The inversion works as follows,

$$GFEA = U$$

$$\Rightarrow FEA = G^{-1}U$$

$$\Rightarrow EA = F^{-1}G^{-1}U$$

$$\Rightarrow A = E^{-1}F^{-1}G^{-1}U$$

$$\Rightarrow A = LU$$

- The inversion works as follows,

$$GFEA = U$$

$$\Rightarrow FEA = G^{-1}U$$

$$\Rightarrow EA = F^{-1}G^{-1}U$$

$$\Rightarrow A = E^{-1}F^{-1}G^{-1}U$$

$$\Rightarrow A = LU$$

- Notice that the order of the multiplication is important.

- Convince yourself that multiplication of two lower (upper) triangular matrices results in a lower (upper) triangular matrix.

- Convince yourself that multiplication of two lower (upper) triangular matrices results in a lower (upper) triangular matrix.
- Also notice that inverse of an upper triangular matrix is another upper triangular matrix.

- Exercise: find the value of x in the following equation.

- Exercise: find the value of x in the following equation.

$$\begin{pmatrix} 1 & 0 & 0 \\ x & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

- One use of LU decomposition in solving linear equations is as follows.

- One use of LU decomposition in solving linear equations is as follows.

$$Ax \uparrow = b \uparrow$$

$$\Rightarrow LUx \uparrow = b \uparrow$$

$$\Rightarrow Ly \uparrow = b \uparrow \quad \text{using} \quad y \uparrow = ux \uparrow$$

- One use of LU decomposition in solving linear equations is as follows.

$$Ax = b$$

$$\Rightarrow L U x = b$$

$$\Rightarrow L y = b \quad \text{using} \quad y = U x$$

- The above equation is easy to solve for y

- One use of LU decomposition in solving linear equations is as follows.

$$\begin{aligned} Ax &\uparrow = b \uparrow \\ \Rightarrow LUx &\uparrow = b \uparrow \\ \Rightarrow Ly &\uparrow = b \uparrow \quad \text{using } y \uparrow = ux \uparrow \end{aligned}$$

- The above equation is easy to solve for $y \uparrow$
- Plugging in $y \uparrow$ in the definition $ux \uparrow = y \uparrow$ and solving for $x \uparrow$ is also easy.