Linear Algebra, in context

Mitaxi Mehta: Lecture 2

Linear equations and solutions

 In this class we discussed linear equations and their solutions. Here is the mind map created collaboratively by us.

Linear equations and solutions

 In this class we discussed linear equations and their solutions. Here is the mind map created collaboratively by us.




Figure: Mind map: linear

 As pointed out by Rushil, when the word linear comes up most of the people think of straight lines.

- As pointed out by Rushil, when the word linear comes up most of the people think of straight lines.
- Mansi thought up a possible connection with equations of one variable, for example x = 3 which works in this case. However all one variable equations are not linear equations, example: $\sin^2(x) + \cos^2(x) = 1$.

- As pointed out by Rushil, when the word linear comes up most of the people think of straight lines.
- Mansi thought up a possible connection with equations of one variable, for example x = 3 which works in this case. However all one variable equations are not linear equations, example: $\sin^2(x) + \cos^2(x) = 1$.
- In answer to, can there be two variable equations connected to the word linear?, Nisarg came up with the general equation ax + by = c. Note that when b is not zero, one can always divide the equation by b to get the standard equation of line in the form y = mx + c', where m = -a/b and c' = c/b.



discussion

 Shubham came up with the unusual idea that any function with power 1 can be associated with the word linear. Later in the course we shall treat general functions like sin, cosine and Hermite and other polinomials within linear algebra.

 Now Dhruvik came up with the suggestion of connecting "linear" with collection of points. I have changed it to connecting two given points (and the extrapolation), along the shportest path. This is a very interesting thought. Though the course will not address it.

- Now Dhruvik came up with the suggestion of connecting "linear" with collection of points. I have changed it to connecting two given points (and the extrapolation), along the shportest path. This is a very interesting thought. Though the course will not address it.
- Aatre started out with arguing that definition of a line in terms of points along the shortest path between two points, will not work in 3D. I convinced the class that the definition works in 3D too. Aatre was thinking about planes in connection with linear algebra, which is quite a well placed thought, We shall talk a lot more about planes and hyperplanes later in the course.

A tangent

• This is a good place to take off on a tangent, When we say plane, we mean geometric planes and not the aeroplanes.

A tangent

- This is a good place to take off on a tangent, When we say plane, we mean geometric planes and not the aeroplanes.
- The same word may mean different things in different contexts. For example a vector means an n-tuple in mathematics, an n-tuple with specific transformation properties in physics and a virus in Biology.

A tangent

- This is a good place to take off on a tangent, When we say plane, we mean geometric planes and not the aeroplanes.
- The same word may mean different things in different contexts. For example a vector means an n-tuple in mathematics, an n-tuple with specific transformation properties in physics and a virus in Biology.
- That is the reason that you need to know your audience before you talk and companies need to understand their customers before designing their products.

 For an interesting discussion on logic, coding etc. a good read is, "Godel, Escher, Bach an eternal golden braid" by Douglas Hofstadter.

- For an interesting discussion on logic, coding etc. a good read is, "Godel, Escher, Bach an eternal golden braid" by Douglas Hofstadter.
- In the race between Achilles and tortoise where the Tortoise has been given a head start, why does Achiles still win?

- For an interesting discussion on logic, coding etc. a good read is, "Godel, Escher, Bach an eternal golden braid" by Douglas Hofstadter.
- In the race between Achilles and tortoise where the Tortoise has been given a head start, why does Achiles still win?
- The enmity between Newton and Liebnitz, non-standard analysis and its relevance.

- For an interesting discussion on logic, coding etc. a good read is, "Godel, Escher, Bach an eternal golden braid" by Douglas Hofstadter.
- In the race between Achilles and tortoise where the Tortoise has been given a head start, why does Achiles still win?
- The enmity between Newton and Liebnitz, non-standard analysis and its relevance.
- A slide entitled "tangent", like this one, is just an aside and does not form a part of the course

Open notebook tests

 All the tests will be open notebook. Please use one big notebook for their notes.

Open notebook tests

- All the tests will be open notebook. Please use one big notebook for their notes.
- Efforts by all should be on learning because copying does disservice to both parties involved, whereas learning is a win-win game.

Open notebook tests

- All the tests will be open notebook. Please use one big notebook for their notes.
- Efforts by all should be on learning because copying does disservice to both parties involved, whereas learning is a win-win game.
- The furure workplaces will value creativity, orginality, problem solving skills and team work much more than memory.

Mind-map Algebra

Mind map of algebra




Figure: Mind map: Algebra

• Two randomly chosen students were asked to give one equation of line each, The equations suggested were, 2x + 3y = 5 and x = y.

- Two randomly chosen students were asked to give one equation of line each, The equations suggested were, 2x + 3y = 5 and x = y.
- Draw the lines, find the intersection point, find x and y axis intercepts.

- Two randomly chosen students were asked to give one equation of line each, The equations suggested were, 2x + 3y = 5 and x = y.
- Draw the lines, find the intersection point, find x and y axis intercepts.
- Select three random lines by having students suggest line equations.

- Two randomly chosen students were asked to give one equation of line each, The equations suggested were, 2x + 3y = 5 and x = y.
- Draw the lines, find the intersection point, find x and y axis intercepts.
- Select three random lines by having students suggest line equations.
- What can you say about the intersection points for such random choice of 3 lines?

- Two randomly chosen students were asked to give one equation of line each, The equations suggested were, 2x + 3y = 5 and x = y.
- Draw the lines, find the intersection point, find x and y axis intercepts.
- Select three random lines by having students suggest line equations.
- What can you say about the intersection points for such random choice of 3 lines?
- Discuss possible number of intersection points for two randomly selected lines.



A solution can be arrived in many different ways. In this
course any method used to arrive at a solution will be valid,
provided you start from fundamentals to build the solution.

- A solution can be arrived in many different ways. In this
 course any method used to arrive at a solution will be valid,
 provided you start from fundamentals to build the solution.
- Some of the methods used for finding intersection points are, (1) geometric (2) computational (3) substitution (4) elimination (5) Inversion of a matrix.

- A solution can be arrived in many different ways. In this
 course any method used to arrive at a solution will be valid,
 provided you start from fundamentals to build the solution.
- Some of the methods used for finding intersection points are, (1) geometric (2) computational (3) substitution (4) elimination (5) Inversion of a matrix.
- Some thoughts on the computational methods of solution: While computational methods of arriving at solutions are getting more accessible and faster in time, mindless use of such methods is likely to weaken your mental abilities. Do apply your mind before using the computer.

• Finding intersection of 2x + 3y = 5 and x - y = 0

- Finding intersection of 2x + 3y = 5 and x y = 0
- substitution.

- Finding intersection of 2x + 3y = 5 and x y = 0
- substitution. solve for x from the second equation (x = y) and substitute in the first to get 5y = 5. Hence the intersection point is x = 1, y = 1.

- Finding intersection of 2x + 3y = 5 and x y = 0
- substitution. solve for x from the second equation (x = y) and substitute in the first to get 5y = 5. Hence the intersection point is x = 1, y = 1.
- Elimination.

- Finding intersection of 2x + 3y = 5 and x y = 0
- substitution. solve for x from the second equation (x = y) and substitute in the first to get 5y = 5. Hence the intersection point is x = 1, y = 1.
- Elimination. Multiply the second equation by 3 and add to the first equation.

more methods

• Inversion of a matrix. Given the equation of the form $A\vec{x} = \vec{b}$.

$$\left(\begin{array}{cc} 2 & 3 \\ 1 & -1 \end{array}\right) \left(\begin{array}{c} x \\ y \end{array}\right) = \left(\begin{array}{c} 5 \\ 0 \end{array}\right)$$

more methods

• Inversion of a matrix. Given the equation of the form $A\vec{x} = \vec{b}$,

$$\left(\begin{array}{cc}2&3\\1&-1\end{array}\right)\left(\begin{array}{c}x\\y\end{array}\right)=\left(\begin{array}{c}5\\0\end{array}\right)$$

The solution is $\vec{x} = A^{-1}\vec{b}$.

$$\left(\begin{array}{c} x \\ y \end{array}\right) = -\frac{1}{5} \left(\begin{array}{cc} -1 & -3 \\ -1 & 2 \end{array}\right) \left(\begin{array}{c} 5 \\ 0 \end{array}\right) = \left(\begin{array}{c} 1 \\ 1 \end{array}\right)$$

Exercise

 Write the following set of equations in matrix form and solve.

Exercise

 Write the following set of equations in matrix form and solve.

$$x + y = 1$$
$$3x + 2y = 5$$