

# Linear Independence

Mitaxi Mehta: Lectures 16

- Solve the following equation

- Solve the following equation

$$\begin{pmatrix} 1 & 2 \\ 1 & -1 \\ 3 & 2 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 3 \\ 2 \\ -7 \end{pmatrix}$$

- Solve the following equation

$$\begin{pmatrix} 1 & 2 \\ 1 & -1 \\ 3 & 2 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 3 \\ 2 \\ -7 \end{pmatrix}$$

- Dhruval:  $x = 2, y = 5,$

- Solve the following equation

$$\begin{pmatrix} 1 & 2 \\ 1 & -1 \\ 3 & 2 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 3 \\ 2 \\ -7 \end{pmatrix}$$

- Dhruval:  $x = 2, y = 5$ , Aarya: Incorrect, it does not satisfy the 3rd equation,

- Solve the following equation

$$\begin{pmatrix} 1 & 2 \\ 1 & -1 \\ 3 & 2 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 3 \\ 2 \\ -7 \end{pmatrix}$$

- Dhruval:  $x = 2, y = 5$ , Aarya: Incorrect, it does not satisfy the 3rd equation, Milind: Two unknowns and 3 equations overdetermined system.

- Geometrically, three randomly selected lines usually don't intersect at one point. Thus such overdetermined systems usually don't have a solution. However in special circumstances they do, how to verify if a solution exists ?

- Geometrically, three randomly selected lines usually don't intersect at one point. Thus such overdetermined systems usually don't have a solution. However in special circumstances they do, how to verify if a solution exists ?
- Jainam: Find solution for each pair of lines and compare (this works but requires more work than necessary), or plot the three lines and see.



- Geometrically, three randomly selected lines usually don't intersect at one point. Thus such overdetermined systems usually don't have a solution. However in special circumstances they do, how to verify if a solution exists ?
- Jainam: Find solution for each pair of lines and compare (this works but requires more work than necessary), or plot the three lines and see. Aarya: Solve for a pair and plug it into the third (this is a good strategy).

- How about,

- How about,

$$\begin{pmatrix} 1 & 1 & 3 & 4 \\ 2 & -1 & 2 & 1 \\ -1 & -4 & -7 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \\ t \end{pmatrix} = \begin{pmatrix} 3 \\ 2 \\ -7 \end{pmatrix} ?$$

- How about,

$$\begin{pmatrix} 1 & 1 & 3 & 4 \\ 2 & -1 & 2 & 1 \\ -1 & -4 & -7 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \\ t \end{pmatrix} = \begin{pmatrix} 3 \\ 2 \\ -7 \end{pmatrix} ?$$

- Aatre: Infinite number of solutions.

- How about,

$$\begin{pmatrix} 1 & 1 & 3 & 4 \\ 2 & -1 & 2 & 1 \\ -1 & -4 & -7 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \\ t \end{pmatrix} = \begin{pmatrix} 3 \\ 2 \\ -7 \end{pmatrix} ?$$

- Aatre: Infinite number of solutions. Preet: Bring to echelon form.

- How about,

$$\begin{pmatrix} 1 & 1 & 3 & 4 \\ 2 & -1 & 2 & 1 \\ -1 & -4 & -7 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \\ t \end{pmatrix} = \begin{pmatrix} 3 \\ 2 \\ -7 \end{pmatrix} ?$$

- Aatre: Infinite number of solutions. Preet: Bring to echelon form.
- One free variable, the solution is a line.

- If number of variables exceeds number of equations then usually there are infinite number of solutions. Such a system is called under-determined system.

- If number of variables exceeds number of equations then usually there are infinite number of solutions. Such a system is called under-determined system.
- A simple example  $3x + 2y = 0$ , every point on the line is a solution.



- If number of variables exceeds number of equations then usually there are infinite number of solutions. Such a system is called under-determined system.
- A simple example  $3x + 2y = 0$ , every point on the line is a solution.
- Another example,

$$2x + 3y + 2 = 0$$

$$x - y + 2 = 5$$

- Solve

- Solve

$$\begin{pmatrix} 1 & 1 & 3 \\ 2 & -1 & 2 \\ -1 & -4 & -7 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 3 \\ 2 \\ -7 \end{pmatrix}$$

- Solve

$$\begin{pmatrix} 1 & 1 & 3 \\ 2 & -1 & 2 \\ -1 & -4 & -7 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 3 \\ 2 \\ -7 \end{pmatrix}$$

- Aarya: The RHS is in the column space, so there is at-least one solution.

- Solve

$$\begin{pmatrix} 1 & 1 & 3 \\ 2 & -1 & 2 \\ -1 & -4 & -7 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 3 \\ 2 \\ -7 \end{pmatrix}$$

- Aarya: The RHS is in the column space, so there is at-least one solution. Drashti: There are infinite number of solutions since  $z$  is a free variable.

- Linear independence: A set of vectors  $\{v_1, v_2, \dots, v_n\}$  is linearly independent if

$$c_1 v_1 + \dots + c_n v_n = 0$$

implies that the numbers  $c_1 = c_2 = \dots = c_n = 0$ .

- Linear independence: A set of vectors  $\{v_1, v_2, \dots, v_n\}$  is linearly independent if

$$c_1 v_1 + \dots + c_n v_n = 0$$

implies that the numbers  $c_1 = c_2 = \dots = c_n = 0$ .

- Are the following two vectors linearly independent ?

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

- Shubham:

$$c_1 \begin{pmatrix} 1 \\ 0 \end{pmatrix} + c_2 \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \Rightarrow \begin{pmatrix} c_1 \\ c_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$



- Shubham:

$$c_1 \begin{pmatrix} 1 \\ 0 \end{pmatrix} + c_2 \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \Rightarrow \begin{pmatrix} c_1 \\ c_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

- Hence the two vectors are linearly independent.

- Are the following two vectors linearly independent ?

- Are the following two vectors linearly independent ?

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

- Are the following two vectors linearly independent ?

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

- Dhuvik: Not linearly independent.

- Are the following two vectors linearly independent ?

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

- Dhuvik: Not linearly independent. If that is the case you should be able to find  $c_1$  and  $c_2$  both nonzero so that  $c_1 v_1 + c_2 v_2 = 0$ .

- Are the following two vectors linearly independent ?

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

- Dhuvik: Not linearly independent. If that is the case you should be able to find  $c_1$  and  $c_2$  both nonzero so that  $c_1 v_1 + c_2 v_2 = 0$ .
- Shrey, does not agree.

- Are the following two vectors linearly independent ?

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

- Dhuvik: Not linearly independent. If that is the case you should be able to find  $c_1$  and  $c_2$  both nonzero so that  $c_1 v_1 + c_2 v_2 = 0$ .
- Shrey, does not agree. in that case show that only  $c_1 = c_2 = 0$  case is possible.

- Maharshi:

$$c_1 \begin{pmatrix} 1 \\ 0 \end{pmatrix} + c_2 \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \Rightarrow \begin{pmatrix} c_1 + c_2 \\ c_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$



- Maharshi:

$$c_1 \begin{pmatrix} 1 \\ 0 \end{pmatrix} + c_2 \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \Rightarrow \begin{pmatrix} c_1 + c_2 \\ c_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

- Hence  $c_1 = c_2 = 0$ .

- Since  $(1, 0)$  can not be written in terms of  $(1, 1)$ , the two vectors are linearly independent.

- Since  $(1, 0)$  can not be written in terms of  $(1, 1)$ , the two vectors are linearly independent.
- How about linear independence of  $(0, 1)$ ,  $(1, 0)$  and  $(1, 1)$ ?

$$\begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \\ c_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

- Since  $(1, 0)$  can not be written in terms of  $(1, 1)$ , the two vectors are linearly independent.
- How about linear independence of  $(0, 1)$ ,  $(1, 0)$  and  $(1, 1)$ ?

$$\begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \\ c_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

- Thankyou Manasi for reminding me that the zero vector has two components and not three.

- Since  $(1, 0)$  can not be written in terms of  $(1, 1)$ , the two vectors are linearly independent.
- How about linear independence of  $(0, 1)$ ,  $(1, 0)$  and  $(1, 1)$ ?

$$\begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \\ c_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

- Thankyou Manasi for reminding me that the zero vector has two components and not three.
- Nonzero  $c_1, c_2, c_3$  exist such that  $c_1 v_1 + c_2 v_2 + c_3 v_3 = 0$ ,  $c_1 = c_2 = 1, c_3 = -1$ . Thus one of the vectors can be written in terms of the rest two.

- A set of linearly independent vectors constitute a basis set of a vector space. Every vector in the space can be written as a linear superposition of the basis vectors.

- A set of linearly independent vectors constitute a basis set of a vector space. Every vector in the space can be written as a linear superposition of the basis vectors.
- $(1, 0), (0, 1)$  is a basis set of  $R^2$  so is,  $(1, 0), (1, 1)$ .

- A set of linearly independent vectors constitute a basis set of a vector space. Every vector in the space can be written as a linear superposition of the basis vectors.
- $(1, 0), (0, 1)$  is a basis set of  $R^2$  so is,  $(1, 0), (1, 1)$ .
- Express  $(3, 2)$  as a linear combination of  $(1, 0)$  and  $(1, 1)$ .



- A set of linearly independent vectors constitute a basis set of a vector space. Every vector in the space can be written as a linear superposition of the basis vectors.
- $(1, 0), (0, 1)$  is a basis set of  $R^2$  so is,  $(1, 0), (1, 1)$ .
- Express  $(3, 2)$  as a linear combination of  $(1, 0)$  and  $(1, 1)$ .  
Ans:  $(3, 2) = (1, 0) + 2(1, 1)$ .

- A basis set is said to be orthogonal if the dot product between any two distinct basis vectors is zero.

- A basis set is said to be orthogonal if the dot product between any two distinct basis vectors is zero.
- In linear algebra, the dot product between two vectors is given as the matrix multiplication between a  $1 \times n$  matrix and a  $n \times 1$  matrix.

$$\vec{a} \cdot \vec{b} \uparrow = \vec{a}^T \vec{b} = (a_1, a_2) \begin{pmatrix} b_1 \\ b_2 \end{pmatrix} = a_1 b_1 + a_2 b_2$$

- A basis set is said to be orthogonal if the dot product between any two distinct basis vectors is zero.
- In linear algebra, the dot product between two vectors is given as the matrix multiplication between a  $1 \times n$  matrix and a  $n \times 1$  matrix.

$$\vec{a} \cdot b \uparrow = a^T b = (a_1, a_2) \begin{pmatrix} b_1 \\ b_2 \end{pmatrix} = a_1 b_1 + a_2 b_2$$

- $\vec{a}$  is called the dual vector of  $a \uparrow$ .

- A basis set is said to be orthogonal if the dot product between any two distinct basis vectors is zero.
- In linear algebra, the dot product between two vectors is given as the matrix multiplication between a  $1 \times n$  matrix and a  $n \times 1$  matrix.

$$\vec{a} \cdot b \uparrow = a^T b = (a_1, a_2) \begin{pmatrix} b_1 \\ b_2 \end{pmatrix} = a_1 b_1 + a_2 b_2$$

- $\vec{a}$  is called the dual vector of  $a \uparrow$ .
- The standard notation for norm is  $\|\vec{a}\|^2 = a^T a$ . Note that in the dot product the order is not important  $\vec{a}b \uparrow = \vec{b}a \uparrow$ .

- Evaluate the matrix corresponding to the product  $a \uparrow \vec{b}$ .

- Evaluate the matrix corresponding to the product  $a \uparrow \vec{b}$ .
- If  $a^T b = 0$  what can you conclude? Give the four possible conclusions.

- Evaluate the matrix corresponding to the product  $a \uparrow \vec{b}$ .
- If  $a^T b = 0$  what can you conclude? Give the four possible conclusions.
- Either the two vectors are orthogonal or one of them is zero or both are zero.