

Solving Linear Equations

Mitaxi Mehta: Lectures 13and14

- The equation $Ax \uparrow = b \uparrow$ can have no solutions, a unique solution or infinitely many solutions.

- The equation $Ax \uparrow = b \uparrow$ can have no solutions, a unique solution or infinitely many solutions.
- When there are are infinitely many solutions, the general solution can be written as,

$$x^* \uparrow = x_p \uparrow + x_n \uparrow$$

- The equation $Ax \uparrow = b \uparrow$ can have no solutions, a unique solution or infinitely many solutions.
- When there are infinitely many solutions, the general solution can be written as,

$$x^* \uparrow = x_p \uparrow + x_n \uparrow$$

- $x_p \uparrow$ is a vector in the row space that solves the equation and $x_n \uparrow$ is any vector in the null space.

- The equation $Ax \uparrow = b \uparrow$ can have no solutions, a unique solution or infinitely many solutions.
- When there are infinitely many solutions, the general solution can be written as,

$$x^* \uparrow = x_p \uparrow + x_n \uparrow$$

- $x_p \uparrow$ is a vector in the row space that solves the equation and $x_n \uparrow$ is any vector in the null space.
- The transformation A takes the row space vector $x_p \uparrow$ to the column space vector $b \uparrow$ and maps any null space vector x_n to the zero vector.

Eucledean vectors

- Consider two vectors $a \uparrow$ and $b \uparrow$, $a \uparrow$ defines displacement from $(0, 0)$ to $(1, 0)$ and $b \uparrow$ defines displacement from $(1, 1)$ to $(2, 1)$.

Eucledean vectors

- Consider two vectors $a \uparrow$ and $b \uparrow$, $a \uparrow$ defines displacement from $(0, 0)$ to $(1, 0)$ and $b \uparrow$ defines displacement from $(1, 1)$ to $(2, 1)$.
- What is the difference between $a \uparrow$ and $b \uparrow$?

Eucledean vectors

- Consider two vectors $a \uparrow$ and $b \uparrow$, $a \uparrow$ defines displacement from $(0, 0)$ to $(1, 0)$ and $b \uparrow$ defines displacement from $(1, 1)$ to $(2, 1)$.
- What is the difference between $a \uparrow$ and $b \uparrow$?
- Shrey Patel: $a \uparrow = b \uparrow$.

Euclidean vectors

- Consider two vectors $a \uparrow$ and $b \uparrow$, $a \uparrow$ defines displacement from $(0, 0)$ to $(1, 0)$ and $b \uparrow$ defines displacement from $(1, 1)$ to $(2, 1)$.
- What is the difference between $a \uparrow$ and $b \uparrow$?
- Shrey Patel: $a \uparrow = b \uparrow$.
- In Euclidean geometry location of a vector is unimportant. The unit vectors are independent of position.

Vector Spaces

- Read the definition of a vector space from Strang (Ed. 4) , page 74 problem 6.

Vector Spaces

- Read the definition of a vector space from Strang (Ed. 4) , page 74 problem 6.
- To define a vector space, one needs two objects. A collection of vectors V , with $v_j \in V$ and a collection of numbers A with members $a, b, \dots \in A$ and a number of properties satisfied by the elements of V and A .

Vector Spaces

- Read the definition of a vector space from Strang (Ed. 4) , page 74 problem 6.
- To define a vector space, one needs two objects. A collection of vectors V , with $v_j \in V$ and a collection of numbers A with members $a, b, \dots \in A$ and a number of properties satisfied by the elements of V and A .
- Please note that different properties apply to numbers and vectors. In this course $A = \mathbb{R}$, the real line, unless stated otherwise.

Vector Spaces

- Question: Consider $V = \mathbb{Z}^2$ that contains all two dimensional vectors with integer entries and $A = \mathbb{R}$, the set of real numbers, do they define a vector space ?

Vector Spaces

- Question: Consider $V = \mathbb{Z}^2$ that contains all two dimensional vectors with integer entries and $A = \mathbb{R}$, the set of real numbers, do they define a vector space ?
- No, because a member of V , when multiplied by a fraction, ceases to be a member of V .

- Consider V to be a set of all 2×2 matrices,

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

with $a, b, c, d \in R$.

- Consider V to be a set of all 2×2 matrices,

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

with $a, b, c, d \in R$.

- Is V a vector space ?

- Consider V to be a set of all 2×2 matrices,

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

with $a, b, c, d \in R$.

- Is V a vector space ?
- Yes, because all the vector space properties are satisfied.

- For a given matrix A , consider the row space $R(A)$, the column space $C(A)$ and the null space $N(A)$.

- For a given matrix A , consider the row space $R(A)$, the column space $C(A)$ and the null space $N(A)$.
- To prove that each of the above spaces is a vector space, show that any linear combination of two vectors of a space also belongs to that space.

- Consider the plane $2x + 3y + z = 10$ and the set of vectors with components (x, y, z) that satisfy the above equation. Is that a vector space ?

- Consider the plane $2x + 3y + z = 10$ and the set of vectors with components (x, y, z) that satisfy the above equation. Is that a vector space ?
- Shubham Pujara: No because zero vector does not belong to the space, i.e., $x = y = z = 0$ does not satisfy the plane equation.

- Consider the plane $2x + 3y + z = 10$ and the set of vectors with components (x, y, z) that satisfy the above equation. Is that a vector space ?
- Shubham Pujara: No because zero vector does not belong to the space, i.e., $x = y = z = 0$ does not satisfy the plane equation.
- Negative vectors also don't belong to the set, for example $(0, 0, 10)$ satisfies the plane equation but $(0, 0, -10)$ does not.

- Exercise: What is the null space of,

$$A = \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}$$

- Exercise: What is the null space of,

$$A = \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}$$

- Dhruv Parikh: Only the zero vector.

- Exercise: What is the null space of,

$$A = \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}$$

- Dhruv Parikh: Only the zero vector.
- For the given matrix, $N(A)$ contains only one vector $(0, 0)$, it is a zero dimensional space.

Echelon form of a matrix

- Echelon form of a matrix is a special form, brought about using row operations and row exchange.

Echelon form of a matrix

- Echelon form of a matrix is a special form, brought about using row operations and row exchange.
- In this form, in every row, the first leading entry (nonzero entry) is such that all elements in the column below it are zero (An upper diagonal form).

Echelon form: example

- Calculate the Echelon form of the following matrix.

Echelon form: example

- Calculate the Echelon form of the following matrix.

$$A = \begin{pmatrix} 1 & 3 & 3 & 2 \\ 2 & 6 & 9 & 7 \\ -1 & -3 & 3 & 4 \end{pmatrix}$$

Echelon form: example

- The row operation $R_3 = R_3 + R_1$ brings the matrix in the form

$$A = \begin{pmatrix} 1 & 3 & 3 & 2 \\ 2 & 6 & 9 & 7 \\ 0 & 0 & 6 & 6 \end{pmatrix}$$

- Proceeding in a manner similar to Gauss elimination using $R_2 = R_2 - (-2)R_1$ and $R_3 = R_3 + (-2)R_2$, the matrix comes into the following form,

$$B = \begin{pmatrix} 1 & 3 & 3 & 2 \\ 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

- The entries B_{11} and B_{23} are the first nonzero entries in the row, they are called pivots.

- The entries B_{11} and B_{23} are the first nonzero entries in the row, they are called pivots.
- The number of pivots in the Echelon form gives the rank of the matrix.

- The entries B_{11} and B_{23} are the first nonzero entries in the row, they are called pivots.
- The number of pivots in the Echelon form gives the rank of the matrix.
- Rank of the matrix also equals the dimensions of $R(A)$ and $C(A)$.

- The entries B_{11} and B_{23} are the first nonzero entries in the row, they are called pivots.
- The number of pivots in the Echelon form gives the rank of the matrix.
- Rank of the matrix also equals the dimensions of $R(A)$ and $C(A)$.
- The calculation of $N(A)$ becomes easy using the Echelon form of A .

Row operations as Matrix multiplication

- Each row operation is equivalent to multiplying the given matrix on the right with another matrix.

Row operations as Matrix multiplication

- Each row operation is equivalent to multiplying the given matrix on the right with another matrix.
- For example, row exchange is equivalent to the following matrix multiplication,

$$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} c & d \\ a & b \end{pmatrix}$$

- The operation $R_2 = R_2 + (g)R_1$ can be written as,

- The operation $R_2 = R_2 + (g)R_1$ can be written as,

$$\begin{pmatrix} 1 & 0 \\ g & 1 \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} a & b \\ g a + c & g b + d \end{pmatrix}$$

- Suppose that one wants to find $N(A)$, i.e. all vectors x such that,

$$Ax = 0.$$

- Suppose that one wants to find $N(A)$, i.e. all vectors x such that,

$$Ax = 0.$$

- Multiplication by a row operation B on both the sides leads to

$$BAx = 0.$$

- Suppose that one wants to find $N(A)$, i.e. all vectors x such that,

$$Ax = 0.$$

- Multiplication by a row operation B on both the sides leads to

$$BAx = 0.$$

- Finally when the matrix is brought to the Echelon form,

$$A_{echelon}x = 0$$

- If A is taken to be matrix of our previous Echelon example,

$$A = \begin{pmatrix} 1 & 3 & 3 & 2 \\ 2 & 6 & 9 & 7 \\ 0 & 0 & 6 & 6 \end{pmatrix}$$

- The equation defining the Null space vectors becomes,

- The equation defining the Null space vectors becomes,

$$\begin{pmatrix} 1 & 3 & 3 & 2 \\ 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

- The pivot entries multiply the variables x_1 and x_3 , x_1 and x_3 are called pivot variables.

- The pivot entries multiply the variables x_1 and x_3 , x_1 and x_3 are called pivot variables.
- x_2 and x_4 are called the free variables.

- The pivot entries multiply the variables x_1 and x_3 , x_1 and x_3 are called pivot variables.
- x_2 and x_4 are called the free variables.
- The pivot variables can be solved for in terms of the free variables.

- The pivot entries multiply the variables x_1 and x_3 , x_1 and x_3 are called pivot variables.
- x_2 and x_4 are called the free variables.
- The pivot variables can be solved for in terms of the free variables.
- The free variables can be assigned any value in R .

- Writing out the two equations in

$$\begin{pmatrix} 1 & 3 & 3 & 2 \\ 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

and solving for the pivot variables gives,

- $x_3 = -x_4$ and $x_1 = -3x_2 + x_4$

- A general null vector can thus be rewritten as,

$$x \uparrow = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} -3x_2 + x_4 \\ x_2 \\ -x_4 \\ x_4 \end{pmatrix} = x_2 \begin{pmatrix} -3 \\ 1 \\ 0 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} 1 \\ 0 \\ -1 \\ 1 \end{pmatrix}$$

- The previous equation describes the general form of vectors in the Null space of A . The dimension of the Null space is 2.

- The previous equation describes the general form of vectors in the Null space of A . The dimension of the Null space is 2.
- The dimension of $N(A) = \text{Number of columns in } A - \text{Rank of } A$.