Justification for row operations

Mitaxi Mehta: Lecture 6

 While diagonalising a matrix, row operations play an important role.

- While diagonalising a matrix, row operations play an important role.
- In this lecture, I justify use of row operations in the context of finding intersection point of two lines.

- While diagonalising a matrix, row operations play an important role.
- In this lecture, I justify use of row operations in the context of finding intersection point of two lines.
- Consider the row space of a matrix A

- While diagonalising a matrix, row operations play an important role.
- In this lecture, I justify use of row operations in the context of finding intersection point of two lines.
- Consider the row space of a matrix A
- What are the row vectors? Shubham: The coefficients of the x and y in each equation forms row vectors.

 Given the components of a row vector, one can define the slope of the corresponding line.

- Given the components of a row vector, one can define the slope of the corresponding line.
- What does $v_1 \uparrow$ give ?

- Given the components of a row vector, one can define the slope of the corresponding line.
- What does v₁ ↑ give ?
- Shubham: Elements of v₁ ↑ define the slope of the first line.

- Given the components of a row vector, one can define the slope of the corresponding line.
- What does v₁ ↑ give ?
- Shubham: Elements of v₁ ↑ define the slope of the first line.
- The RHS along with the coefficients of x and y define intercepts.

- Given the components of a row vector, one can define the slope of the corresponding line.
- What does v₁ ↑ give ?
- Shubham: Elements of v₁ ↑ define the slope of the first line.
- The RHS along with the coefficients of x and y define intercepts.
- What is the geometric meaning of $c_1v_1 \uparrow + c_2v_2 \uparrow$?

- Given the components of a row vector, one can define the slope of the corresponding line.
- What does v₁ ↑ give ?
- Shubham: Elements of v₁ ↑ define the slope of the first line.
- The RHS along with the coefficients of x and y define intercepts.
- What is the geometric meaning of $c_1v_1 \uparrow + c_2v_2 \uparrow$? Ans: Coefficients of a new line.

Row space

• If the RHS of the line equations are also added like $c_1b_1 + c_2b_2$ Then the new line equation formed is such that it passes through the same intersection point as the previous two lines.

Row space

- If the RHS of the line equations are also added like $c_1b_1 + c_2b_2$ Then the new line equation formed is such that it passes through the same intersection point as the previous two lines.
- Example: Consider two line equations x + y = 2, x y = 4. The intersection point is (3,-1).
- The line equation using suprposition and $c_1 = 2$, $c_2 = 1$ is 3x + y = 8. Notice that the new line also passes through the same intersection point.

• In general consider two lines $a_1x + b_1y - c_1 = 0$ and $a_2x + b_2y - c_2 = 0$.

- In general consider two lines $a_1x + b_1y c_1 = 0$ and $a_2x + b_2y c_2 = 0$.
- If the intersection point is (x_0, y_0) then the point lies on both the lines. $a_1x_0 + b_1y_0 c_1 = 0$ and $a_2x_0 + b_2y_0 c_2 = 0$.

- In general consider two lines $a_1x + b_1y c_1 = 0$ and $a_2x + b_2y c_2 = 0$.
- If the intersection point is (x_0, y_0) then the point lies on both the lines. $a_1x_0 + b_1y_0 c_1 = 0$ and $a_2x_0 + b_2y_0 c_2 = 0$.
- Considering the superposition, LHS = $d_1(a_1x + b_1y c_1) + d_2(a_2x + b_2y c_2) = 0$ is satisfied by the same intersection point.