# The Pendulum Equation

Mitaxi Mehta Lecture 3



What is a pendulum?

- What is a pendulum?
- Can you imagine pendulum like parts occuring in a machine? Where?

- What is a pendulum?
- Can you imagine pendulum like parts occuring in a machine? Where?
- What about human dynamics?

- What is a pendulum?
- Can you imagine pendulum like parts occuring in a machine? Where?
- What about human dynamics?
- Why should chemical engineers study a pendulum?

- What is a pendulum?
- Can you imagine pendulum like parts occuring in a machine? Where?
- What about human dynamics?
- Why should chemical engineers study a pendulum?
- Do an experiment at your leisure, Take a pendulum and make it oscillate. Does the oscillation time period depend on the amplitude?

 Please note that for a usual pendulum (a bob of mass m, hung by a rod of length I, having negligible mass and pivoted at a fixed point) the time period of the pendulum depends on the amplitude.

- Please note that for a usual pendulum (a bob of mass m, hung by a rod of length I, having negligible mass and pivoted at a fixed point) the time period of the pendulum depends on the amplitude.
- Only in the limit of small oscillations, the time period is independent of the amplitude.

• What is the force under which a pendulum moves?

- What is the force under which a pendulum moves?
- What is wrong with the above question?

- What is the force under which a pendulum moves?
- What is wrong with the above question? You don't require a force to move a body.

- What is the force under which a pendulum moves?
- What is wrong with the above question? You don't require a force to move a body.
- Let us re-frame the question, what is the force under which the bob accelerates?

• If you were a bob of mass m what are the forces that you will feel?

- If you were a bob of mass m what are the forces that you will feel?
- Which is the force that actually accelerates the bob at any given point of time?

• The equation of the pendulum is,

$$\vec{F} = m\vec{a}$$
  $\rightarrow$   $ml\ddot{\theta} = -mg\sin\theta$ 

The equation of the pendulum is,

$$\vec{F} = m\vec{a}$$
  $\rightarrow$   $ml\ddot{\theta} = -mg\sin\theta$ 

• Can you explain why did I write the acceleration as above?

The equation of the pendulum is,

$$\vec{F} = m\vec{a} \qquad o \qquad ml\ddot{ heta} = -mg\sin\theta$$

• Can you explain why did I write the acceleration as above? Hint: Arc length =  $I\theta$ 

• In other words,

$$\ddot{\theta} = -\frac{g}{I}\sin\theta$$

In other words,

$$\ddot{\theta} = -\frac{g}{I}\sin\theta$$

 This differential equation is not easy to solve (for a person working with second order ODE for the first time), the solution can be written in terms of Jacobi Elliptic function. • Important terms to know, dependent and independent variables.

- Important terms to know, dependent and independent variables.
- What is the dependent variable and what is the independent variable in the pendulum equation?

- Important terms to know, dependent and independent variables.
- What is the dependent variable and what is the independent variable in the pendulum equation?
- How do we decide? Give some examples.

- Important terms to know, dependent and independent variables.
- What is the dependent variable and what is the independent variable in the pendulum equation?
- How do we decide? Give some examples. What can you say about position, velocity and acceleration of a point mass are they dependent or independent?

- Important terms to know, dependent and independent variables.
- What is the dependent variable and what is the independent variable in the pendulum equation?
- How do we decide? Give some examples. What can you say about position, velocity and acceleration of a point mass are they dependent or independent?
- Example: Mass attached to a spring. Draw a coordinate system define variables and write the differential equation.

• Exercise: Force due to air damping on a body is given by  $\vec{F} = -\gamma \vec{v}$ .

 Exercise: Force due to air damping on a body is given by \vec{F} = -γ \vec{v}\$. Why is there a minus sign? Use this force to write the differential equation of (1) A free falling body under gravity (2) A ball thrown up with velocity \vec{v}\$ from the ground. (Define and draw appropriate coordinate system)

- Exercise: Force due to air damping on a body is given by \vec{F} = -γ \vec{v}\$. Why is there a minus sign? Use this force to write the differential equation of (1) A free falling body under gravity (2) A ball thrown up with velocity \vec{v}\$ from the ground. (Define and draw appropriate coordinate system)
- Exercise: Consider 2 simple dynamical systems in your surrounding, describe them, draw appropriate coordinate system and write down the corresponding differential equations. (It is alright to search the web if you can't start, but please give the webpage URL you have used and highlight if you have made any changes in that information).

 The pendulum equation is a difficult one to solve when one is just starting with differential equations. Let us start with a simpler equation.  The pendulum equation is a difficult one to solve when one is just starting with differential equations. Let us start with a simpler equation.

$$\frac{dy}{dx} + 2y = 0.$$

You already know how to solve this.

 The pendulum equation is a difficult one to solve when one is just starting with differential equations. Let us start with a simpler equation.

$$\frac{dy}{dx} + 2y = 0.$$

- You already know how to solve this.
- However there is a new way of thinking about this equation.

• Consider  $\frac{d}{dx}$ ,

• Consider  $\frac{d}{dx}$ , Is this a number ?

• Consider  $\frac{d}{dx}$ , Is this a number ? Is this a function ?

- Consider  $\frac{d}{dx}$ , Is this a number ? Is this a function ?
- $\frac{d}{dx}$  is a mathematical operation, which when it acts on a function, gives another function. Such an expression is called an operator. We shall denote  $D = \frac{d}{dx}$

 Note that D is not commutative with respect to the function it operates on.

 Note that D is not commutative with respect to the function it operates on.

D3 is not 3D. Also note that f(x)D is a differential operator, whereas Df(x) is a function.

 Note that D is not commutative with respect to the function it operates on.

D3 is not 3D. Also note that f(x)D is a differential operator, whereas Df(x) is a function.

D3 = 0, 
$$3D = 3\frac{d}{dx}$$
  
 $Df(x) = \frac{df(x)}{dx}$ 

 However, note that the following kind of commutation works.

$$(D-3)(D-2)f(x) = (D-2)(D-3)f(x)$$

 However, note that the following kind of commutation works.

$$(D-3)(D-2)f(x) = (D-2)(D-3)f(x)$$

Calculate and show.

 However, note that the following kind of commutation works.

$$(D-3)(D-2)f(x) = (D-2)(D-3)f(x)$$

- Calculate and show.
- What is the value of (D-3) operating on zero?