

Approximation of functions

Mitaxi Mehta
Lecture 6

- In the pendulum equation, why is $\sin \theta$ replaced by θ (and not θ^2 or 2θ) ? Give as many explanations as you can.

- In the pendulum equation, why is $\sin \theta$ replaced by θ (and not θ^2 or 2θ) ? Give as many explanations as you can.
- Suppose you were asked to approximate $\sin \theta$ in the neighborhood of $\pi/4$ how would you go about doing it?

- What information you have about $\sin \theta$ at $\theta = \pi/4$?

- What information you have about $\sin \theta$ at $\theta = \pi/4$?
- Did you notice that a large number of statements so far have been questions?

- What information you have about $\sin \theta$ at $\theta = \pi/4$?
- Did you notice that a large number of statements so far have been questions?
- Probing knowledge through interrogation is known as Socratic (sometimes also Platonic) dialogue, I picked this up from my late teacher Prof. C S R Murthy of PRL, who taught me statistics.

- How much information do you have of $\sin \theta$ about $\theta = \pi/4$?

- How much information do you have of $\sin \theta$ about $\theta = \pi/4$?
- Can that information be used to gain better approximation of $\sin \theta$?

Approximations.

- If Δx is small,

$$f(x_0 + \Delta x) \approx f(x_0) + f'(x_0)\Delta x.$$

Approximations.

- If Δx is small,

$$f(x_0 + \Delta x) \approx f(x_0) + f'(x_0)\Delta x.$$

(Note: Δx is different from dx and ∂x .)

Approximations.

- If Δx is small,

$$f(x_0 + \Delta x) \approx f(x_0) + f'(x_0)\Delta x.$$

(Note: Δx is different from dx and ∂x .)

- The above formula is useful in finding approximate value of a function, if $f(x_0)$ and $f'(x_0)$ are known.

Approximations.

- If Δx is small,

$$f(x_0 + \Delta x) \approx f(x_0) + f'(x_0)\Delta x.$$

(Note: Δx is different from dx and ∂x .)

- The above formula is useful in finding approximate value of a function, if $f(x_0)$ and $f'(x_0)$ are known.
- Select Δx to be as small as possible.

- Find the approximate value of $(7.8)^{1/3}$, using $f(x_0 + \Delta x) \approx f(x_0) + f'(x_0)\Delta x$.

- Find the approximate value of $(7.8)^{1/3}$, using $f(x_0 + \Delta x) \approx f(x_0) + f'(x_0)\Delta x$.
- $f(x) =$

- Find the approximate value of $(7.8)^{1/3}$, using $f(x_0 + \Delta x) \approx f(x_0) + f'(x_0)\Delta x$.
- $f(x) = x^{1/3}$,

- Find the approximate value of $(7.8)^{1/3}$, using $f(x_0 + \Delta x) \approx f(x_0) + f'(x_0)\Delta x$.
- $f(x) = x^{1/3}$, $x_0 =$

- Find the approximate value of $(7.8)^{1/3}$, using $f(x_0 + \Delta x) \approx f(x_0) + f'(x_0)\Delta x$.
- $f(x) = x^{1/3}$, $x_0 = 8$,

- Find the approximate value of $(7.8)^{1/3}$, using $f(x_0 + \Delta x) \approx f(x_0) + f'(x_0)\Delta x$.
- $f(x) = x^{1/3}$, $x_0 = 8$, $f(x_0) = (8)^{1/3} = 2$

- Find the approximate value of $(7.8)^{1/3}$, using $f(x_0 + \Delta x) \approx f(x_0) + f'(x_0)\Delta x$.
- $f(x) = x^{1/3}$, $x_0 = 8$, $f(x_0) = (8)^{1/3} = 2$
 $f'(x_0) =$

- Find the approximate value of $(7.8)^{1/3}$, using $f(x_0 + \Delta x) \approx f(x_0) + f'(x_0)\Delta x$.
- $f(x) = x^{1/3}$, $x_0 = 8$, $f(x_0) = (8)^{1/3} = 2$
 $f'(x_0) = \frac{1}{3}8^{-2/3} = \frac{1}{12}$,

- Find the approximate value of $(7.8)^{1/3}$, using $f(x_0 + \Delta x) \approx f(x_0) + f'(x_0)\Delta x$.
- $f(x) = x^{1/3}$, $x_0 = 8$, $f(x_0) = (8)^{1/3} = 2$
 $f'(x_0) = \frac{1}{3}8^{-2/3} = \frac{1}{12}$, $\Delta x =$

- Find the approximate value of $(7.8)^{1/3}$, using $f(x_0 + \Delta x) \approx f(x_0) + f'(x_0)\Delta x$.
- $f(x) = x^{1/3}$, $x_0 = 8$, $f(x_0) = (8)^{1/3} = 2$
 $f'(x_0) = \frac{1}{3}8^{-2/3} = \frac{1}{12}$, $\Delta x = -.2$

- Find the approximate value of $(7.8)^{1/3}$, using $f(x_0 + \Delta x) \approx f(x_0) + f'(x_0)\Delta x$.

- $f(x) = x^{1/3}$, $x_0 = 8$, $f(x_0) = (8)^{1/3} = 2$
 $f'(x_0) = \frac{1}{3}8^{-2/3} = \frac{1}{12}$, $\Delta x = -.2$

$$(7.8)^{1/3} = (8 - .2)^{1/3}$$

- Find the approximate value of $(7.8)^{1/3}$, using $f(x_0 + \Delta x) \approx f(x_0) + f'(x_0)\Delta x$.

- $f(x) = x^{1/3}$, $x_0 = 8$, $f(x_0) = (8)^{1/3} = 2$
 $f'(x_0) = \frac{1}{3}8^{-2/3} = \frac{1}{12}$, $\Delta x = -.2$

$$(7.8)^{1/3} = (8 - .2)^{1/3} \approx 2 + \frac{1}{12}(-.2) = 2 - .0167 = 1.9833$$

- Find the approximate value of $(7.8)^{1/3}$, using $f(x_0 + \Delta x) \approx f(x_0) + f'(x_0)\Delta x$.

- $f(x) = x^{1/3}$, $x_0 = 8$, $f(x_0) = (8)^{1/3} = 2$
 $f'(x_0) = \frac{1}{3}8^{-2/3} = \frac{1}{12}$, $\Delta x = -.2$

$$(7.8)^{1/3} = (8 - .2)^{1/3} \approx 2 + \frac{1}{12}(-.2) = 2 - .0167 = 1.9833$$

- The actual answer is close to 1.9831.

- The steps are (1) Identify $f(x)$, x_0 and Δx .

- The steps are (1) Identify $f(x)$, x_0 and Δx .
Note: x is a variable, x_0 and Δx are values.

- The steps are (1) Identify $f(x)$, x_0 and Δx .
Note: x is a variable, x_0 and Δx are values.
- Substitute in the formula.

- Find the approximate value of $\sin(1^\circ)$.

- Find the approximate value of $\sin(1^\circ)$. $f(x) =$

- Find the approximate value of $\sin(1^\circ)$. $f(x) = \sin(x)$,

- Find the approximate value of $\sin(1^\circ)$. $f(x) = \sin(x)$,
 $x_0 =$

- Find the approximate value of $\sin(1^\circ)$. $f(x) = \sin(x)$,
 $x_0 = 0^\circ$,

- Find the approximate value of $\sin(1^\circ)$. $f(x) = \sin(x)$,
 $x_0 = 0^\circ$, $\Delta x =$

- Find the approximate value of $\sin(1^\circ)$. $f(x) = \sin(x)$,
 $x_0 = 0^\circ$, $\Delta x = 1^\circ$.

- Find the approximate value of $\sin(1^\circ)$. $f(x) = \sin(x)$, $x_0 = 0^\circ$, $\Delta x = 1^\circ$.

$$\sin(1^\circ) \approx \sin(0^\circ + 1^\circ)$$

- Find the approximate value of $\sin(1^\circ)$. $f(x) = \sin(x)$, $x_0 = 0^\circ$, $\Delta x = 1^\circ$.

$$\sin(1^\circ) \approx \sin(0^\circ + 1^\circ)$$

$$\text{using } f(x_0 + \Delta x) \approx f(x_0) + f'(x_0)\Delta x$$

- Find the approximate value of $\sin(1^\circ)$. $f(x) = \sin(x)$, $x_0 = 0^\circ$, $\Delta x = 1^\circ$.

$$\sin(1^\circ) \approx \sin(0^\circ + 1^\circ)$$

$$\text{using } f(x_0 + \Delta x) \approx f(x_0) + f'(x_0)\Delta x$$

$$\Rightarrow \sin(1^\circ) \approx \sin 0 + 1 \cos 0 = 0 + 1 \times 1 = 1.$$

- Find the approximate value of $\sin(1^\circ)$. $f(x) = \sin(x)$, $x_0 = 0^\circ$, $\Delta x = 1^\circ$.

$$\sin(1^\circ) \approx \sin(0^\circ + 1^\circ)$$

$$\text{using } f(x_0 + \Delta x) \approx f(x_0) + f'(x_0)\Delta x$$

$$\Rightarrow \sin(1^\circ) \approx \sin 0 + 1 \cos 0 = 0 + 1 \times 1 = 1.$$

What is wrong ?

Approximation of trigonometric functions.

- While approximating trigonometric functions, the variable should be expressed in radians and not degrees.

Approximation of trigonometric functions.

- While approximating trigonometric functions, the variable should be expressed in radians and not degrees. The correct calculation is,

$$\sin\left(\frac{\pi}{180}\right) \approx \sin\left(0 + \frac{\pi}{180}\right)$$

Approximation of trigonometric functions.

- While approximating trigonometric functions, the variable should be expressed in radians and not degrees. The correct calculation is,

$$\sin\left(\frac{\pi}{180}\right) \approx \sin\left(0 + \frac{\pi}{180}\right)$$

$$f(x) = \sin x, \quad x_0 = 0, \quad \Delta x = \frac{\pi}{180} = .0174$$

Approximation of trigonometric functions.

- While approximating trigonometric functions, the variable should be expressed in radians and not degrees. The correct calculation is,

$$\sin\left(\frac{\pi}{180}\right) \approx \sin\left(0 + \frac{\pi}{180}\right)$$

$$f(x) = \sin x, \quad x_0 = 0, \quad \Delta x = \frac{\pi}{180} = .0174$$

$$\Rightarrow \sin(1^\circ) \approx 0 + .0174 \times 1 = .0174.$$

- What are the dimensions of an angle ?

- What are the dimensions of an angle ?
- How many different interpretations (Geometric or otherwise) can you find for the \sin and \cos functions ?

Some interpretations of sin and cos.

- Ratio of sides to hypotenuse in a right angle triangle.

Some interpretations of \sin and \cos .

- Ratio of sides to hypotenuse in a right angle triangle.
- Projection on the x and y axis, of a point on the unit circle.

Some interpretations of sin and cos.

- Ratio of sides to hypotenuse in a right angle triangle.
- Projection on the x and y axis, of a point on the unit circle.
- Summation of infinite power series.

Some interpretations of sin and cos.

- Ratio of sides to hypotenuse in a right angle triangle.
- Projection on the x and y axis, of a point on the unit circle.
- Summation of infinite power series.
- Solutions of the differential equation, $\frac{d^2 u}{dx^2} = -u$.

Some interpretations of sin and cos.

- Ratio of sides to hypotenuse in a right angle triangle.
- Projection on the x and y axis, of a point on the unit circle.
- Summation of infinite power series.
- Solutions of the differential equation, $\frac{d^2 u}{dx^2} = -u$.
- Often the best way of interpreting a new transcendental function is as a solution of a particular differential equation.

Use of differentials in Error Calculations.

- While doing experiments, a standard source of errors is the least count of an instrument.

Use of differentials in Error Calculations.

- While doing experiments, a standard source of errors is the least count of an instrument.
- One would like to find out the uncertainty in the final result due to the uncertainty in measurements.

Use of differentials in Error Calculations.

- While doing experiments, a standard source of errors is the least count of an instrument.
- One would like to find out the uncertainty in the final result due to the uncertainty in measurements.
- If the error in the measurement of the radius of a cylinder of height 10 cm.s and radius 5 is .1 cm.s (i.e. the radius is measured to be $(5 \pm .1)$ cm.s) What is the error in the calculation of its volume ?

Use of differentials in Error Calculations.

- While doing experiments, a standard source of errors is the least count of an instrument.
- One would like to find out the uncertainty in the final result due to the uncertainty in measurements.
- If the error in the measurement of the radius of a cylinder of height 10 cm.s and radius 5 is .1 cm.s (i.e. the radius is measured to be $(5 \pm .1)$ cm.s) What is the error in the calculation of its volume ?

$$V = \pi r^2 h \quad \Rightarrow \quad \Delta v \approx 2\pi r h \Delta r = 10\pi \text{ cm}^3$$

- In the previous example find the relative error in the volume.

- In the previous example find the relative error in the volume.
- Find the percentage error in the volume.

- The same analysis can also be used to find the change in an outcome due to a small change in a parameter value.

- The same analysis can also be used to find the change in an outcome due to a small change in a parameter value.
- Example: Blood flow through the arteries. A small change in the radius due to cholesterol deposition leads to a big change in the rate of blood flow.

- The same analysis can also be used to find the change in an outcome due to a small change in a parameter value.
- Example: Blood flow through the arteries. A small change in the radius due to cholesterol deposition leads to a big change in the rate of blood flow.

If $Flow \propto r^4$,
10% change in the radius \Rightarrow 40%.change in the flow.