

Pendulum equation and solution for small oscillations.

Mitaxi Mehta
Lecture 5

Homework for week-1

- Homework has been posted. To be submitted by students whose surnames start with "S" in Friday's class.

Homework for week-1

- Homework has been posted. To be submitted by students whose surnames start with "S" in Friday's class.
- An idle question "why don't we have 0, negative and complex roll numbers ?

Homework for week-1

- Homework has been posted. To be submitted by students whose surnames start with "S" in Friday's class.
- An idle question "why don't we have 0, negative and complex roll numbers ? If we want to devise them what are your suggestions?"

Homework for week-1

- Homework has been posted. To be submitted by students whose surnames start with "S" in Friday's class.
- An idle question "why don't we have 0, negative and complex roll numbers ? If we want to devise them what are your suggestions?"
- Note that numbers are just labels for the real life information we want to deal with.

- Consider the small angle approximation of the pendulum equation,

$$\ddot{\theta} = -\frac{g}{l} \sin \theta$$

to convert it into a harmonic oscillator.

- Consider the small angle approximation of the pendulum equation,

$$\ddot{\theta} = -\frac{g}{l} \sin \theta$$

to convert it into a harmonic oscillator.

- Write the general solution of the equation in terms of cosine and sine and write the arbitrary coefficients in terms of initial position $\theta(0)$ and initial velocity $\frac{d\theta}{dt}|_{t=0}$.

- Write the solution in terms of the the exponential functions.

- Write the solution in terms of the the exponential functions.
- Use $e^{\pm i\theta} = \cos\theta \pm i\sin\theta$ to rewrite the solution in terms of cos and sin functions.

- Write the solution in terms of the the exponential functions.
- Use $e^{\pm i\theta} = \cos\theta \pm i\sin\theta$ to rewrite the solution in terms of cos and sin functions.
- Use the initial conditions to evaluate the arbitrary constants in the solution.

- Write the solution in terms of the the exponential functions.
- Use $e^{\pm i\theta} = \cos\theta \pm i\sin\theta$ to rewrite the solution in terms of cos and sin functions.
- Use the initial conditions to evaluate the arbitrary constants in the solution.
- Note that g and l are system parameters defined in the equation, the constants of motion on the other hand are defined in terms of initial conditions.

- Write the expression of Amplitude in terms of system parameters and initial values.

- Write the expression of Amplitude in terms of system parameters and initial values.
- Note that this can be done by writing the solution as $A \sin(\omega t + \phi)$.

- Write the expression of Amplitude in terms of system parameters and initial values.
- Note that this can be done by writing the solution as $A \sin(\omega t + \phi)$.
- Find the time period T by insisting that as $t \rightarrow t + T$, the solution does not change.