

# The Harmonic Oscillator

Mitaxi Mehta  
Lecture 4

# Working with operators

- However, note that the following kind of commutation works.

$$(D - 3)(D - 2)f(x) = (D - 2)(D - 3)f(x)$$

# Working with operators

- However, note that the following kind of commutation works.

$$(D - 3)(D - 2)f(x) = (D - 2)(D - 3)f(x)$$

- Calculate and show.

# Working with operators

- However, note that the following kind of commutation works.

$$(D - 3)(D - 2)f(x) = (D - 2)(D - 3)f(x)$$

- Calculate and show.
- What is the value of  $(D - 3)$  operating on zero?

- We already know the following,

$$\frac{dy}{dx} - 2y = (D - 2)y = 0$$

has the solution  $y(x) = c_1 e^{2x}$

- We already know the following,

$$\frac{dy}{dx} - 2y = (D - 2)y = 0$$

has the solution  $y(x) = c_1 e^{2x}$

- Notice that the same function will also solve,

$$\frac{d^2y}{dx^2} - 5\frac{dy}{dx} + 6y = (D - 3)(D - 2)y = 0$$

- Similarly  $c_2 e^{3x}$  solves

$$\frac{dy}{dx} - 3y = (D - 3)y = 0$$

- Similarly  $c_2 e^{3x}$  solves

$$\frac{dy}{dx} - 3y = (D - 3)y = 0$$

- so it also is a solution of

$$\frac{d^2y}{dx^2} - 5\frac{dy}{dx} + 6y = (D - 3)(D - 2)y = 0$$



- Does that mean that there are two different solutions  $c_1 e^{2x}$  and  $c_2 e^{3x}$ , for the differential equation

$$\frac{d^2 y}{dx^2} - 5 \frac{dy}{dx} + 6y = 0?$$

what do you think?

- Does that mean that there are two different solutions  $c_1 e^{2x}$  and  $c_2 e^{3x}$ , for the differential equation

$$\frac{d^2 y}{dx^2} - 5 \frac{dy}{dx} + 6y = 0?$$

what do you think?

- The equation is a second order differential equation, so solving it requires two integration and hence two constants of integration.

- The expression  $y(x) = c_1 e^{2x} + c_2 e^{3x}$  is known as the general solution of the given differential equation.

- The expression  $y(x) = c_1 e^{2x} + c_2 e^{3x}$  is known as the general solution of the given differential equation.
- We shall discuss similar such equations later.

- Consider Oscillatory motion of a point mass inside a potential well  $V(x)$ .

- Consider Oscillatory motion of a point mass inside a potential well  $V(x)$ .
- The force on the mass is given by  $\vec{F} = -\frac{dV}{dx}$ .

- Consider Oscillatory motion of a point mass inside a potential well  $V(x)$ .
- The force on the mass is given by  $\vec{F} = -\frac{dV}{dx}$ .
- Notice that the force is zero at the maxima and minima of the potential. Those are equilibrium of the particle

- Also notice that near the minima the force is restoring, it is directed towards the equilibrium position.



- Also notice that near the minima the force is restoring, it is directed towards the equilibrium position.
- Hence near the minimum of a potential, the particle with a low energy exhibits oscillatory motion.

- Also notice that near the minima the force is restoring, it is directed towards the equilibrium position.
- Hence near the minimum of a potential, the particle with a low energy exhibits oscillatory motion.
- Why does not the particle stop as it arrives at the equilibrium?

# Harmonic oscillations

- The simplest expression of force near equilibrium is linear  $F = -kx$ , which is the force on a mass attached to a spring. (What happened to the equilibrium length?)

# Harmonic oscillations

- The simplest expression of force near equilibrium is linear  $F = -kx$ , which is the force on a mass attached to a spring. (What happened to the equilibrium length?)
- Applying Newton's second law, the equation of motion is,

$$m \frac{d^2 x}{dt^2} = -kx$$

- In operator notation the equation can be written as

$$\frac{d^2x}{dt^2} + \frac{k}{m}x = 0 \quad \rightarrow \quad (D^2 + \omega^2)x = 0$$

- In operator notation the equation can be written as

$$\frac{d^2x}{dt^2} + \frac{k}{m}x = 0 \quad \rightarrow \quad (D^2 + \omega^2)x = 0$$

- As seen in the previous example the general solution can be written as,

$$x(t) = c_1 e^{i\omega t} + c_2 e^{-i\omega t}$$

- Solve the Harmonic oscillator equation for initial conditions  $x(t = 0) = 2$  and  $\frac{dx}{dt}(t = 0) = 0$ .

- Solve the Harmonic oscillator equation for initial conditions  $x(t = 0) = 2$  and  $\frac{dx}{dt}(t = 0) = 0$ .
- Rewrite the expression of  $x(t)$  in terms of sin and cos functions.



- Solve the Harmonic oscillator equation for initial conditions  $x(t = 0) = 2$  and  $\frac{dx}{dt}(t = 0) = 0$ .
- Rewrite the expression of  $x(t)$  in terms of sin and cos functions.
- Rewrite expression as  $x(t) = C\sin(\omega t + \phi)$ . What is the physical meaning of  $\phi$  ?

- Note that we have looked at three different ways of writing the solution of the differential equation of a Simple Harmonic Oscillator (SHO), they are,

- Note that we have looked at three different ways of writing the solution of the differential equation of a Simple Harmonic Oscillator (SHO), they are,  
(1)  $x(t) = c_1 e^{i\omega t} + c_2 e^{-i\omega t}$

- Note that we have looked at three different ways of writing the solution of the differential equation of a Simple Harmonic Oscillator (SHO), they are,  
(1)  $x(t) = c_1 e^{i\omega t} + c_2 e^{-i\omega t}$   
(2)  $x(t) = A \cos(\omega t) + B \sin(\omega t)$

- Note that we have looked at three different ways of writing the solution of the differential equation of a Simple Harmonic Oscillator (SHO), they are,

$$(1) x(t) = c_1 e^{i\omega t} + c_2 e^{-i\omega t}$$

$$(2) x(t) = A \cos(\omega t) + B \sin(\omega t)$$

$$(3) x(t) = C \sin(\omega t + \phi)$$

- Note that all the three expressions are mathematically equivalent after rewriting the coefficients appropriately.

- Note that all the three expressions are mathematically equivalent after rewriting the coefficients appropriately.
- Once the general solution is known one can write the solution of SHO for any set of initial values.

- Exercise

(1) Make the small angle approximation of the pendulum equation  $\sin \theta \approx \theta$  and write down the solution.



- Exercise

(1) Make the small angle approximation of the pendulum equation  $\sin \theta \approx \theta$  and write down the solution.

(2) A mass attached to a spring hangs vertically down with the other end of the spring held fixed. Write the corresponding differential equation and solve.

- Write the position vector of a point in uniform circular motion as a function of time.

- Write the position vector of a point in uniform circular motion as a function of time.  
Write the position vector of a point on the rim of a bicycle wheel rolling in straight line.