

Nuclear Engineering 150 – Discussion Section
Team Exercises #3

Problem 1

A reactor is operating for a long time at some known power density P_0 . Then, it instantaneously changes power to some power density P_1 . One fission product of interest is ^{135}Xe , though it has a negligible yield from the initial fission reaction. ^{135}Xe precursors ^{135}Te and ^{135}I are produced with a combined yield of approximately 6%, before decaying via β^- decay to ^{135}I and ^{135}Xe respectively. Find the number density of ^{135}Xe as a function of time after the power change. (Your solution may be left as variables)

Nucleus	Half-life	Thermal σ_a
^{135}Te	19.0 s	~ 0
^{135}I	6.6 hr	~ 0
^{135}Xe	9.2 hr	2.6×10^6 barns

Problem 2

- a) Find the excitation energy in ^{236}U when a neutron with zero kinetic energy is absorbed by ^{235}U .
- b) Find the excitation energy in ^{239}U when a neutron with zero kinetic energy is absorbed by ^{238}U .
- c) The activation energy for ^{236}U is 6.2 MeV and the activation energy for ^{239}U is 6.6 MeV. Will fission occur in each of these cases? Identify ^{235}U and ^{238}U as fissile or fissionable and explain.
- d) A ^{238}U nuclei absorbs a 2 MeV neutron and fissions into ^{132}Sn , ^{106}Mo , and a neutron. If the neutron is produced with 2.5% of the total energy released in the reaction, does it have enough energy to fission another ^{238}U atom?

Nucleus	Mass
n	amu
^{106}Mo	amu
^{132}Sn	amu
^{235}U	amu
^{236}U	amu
^{238}U	amu
^{239}U	amu