

Paleo-oscillomics: inferring aspects of Neanderthal language abilities from gene regulation of neural oscillations

Elliot Murphy^{1,2} & Antonio Benítez-Burraco³

1) Division of Psychology and Language Sciences, University College London, London, UK
e-mail: elliott.murphy.13@ucl.ac.uk

2) Department of Psychology, University of Westminster, London, UK

3) Department of Spanish, Linguistics, and Theory of Literature. Faculty of Philology, University of Seville, Seville, Spain
e-mail: abenitez8@us.es

Summary - Language seemingly evolved from changes in brain anatomy and wiring. We argue that language evolution can be better understood if particular changes in phasal and cross-frequency coupling properties of neural oscillations, resulting in core features of language, are considered. Because we cannot track the oscillatory activity of the brain from extinct hominins, we used our current understanding of the language oscillogenome (that is, the set of genes responsible for basic aspects of the oscillatory activity relevant for language) to infer some properties of the Neanderthal oscillome. We have found that several candidates for the language oscillogenome show differences in their methylation patterns between Neanderthals and humans. We argue that differences in their expression levels could be informative of differences in cognitive functions important for language.

Keywords - Language evolution, Language disorders, Neanderthals, Candidate-gene approach, Neural oscillations, Methylation patterns.

Introduction

Language is the most distinctive human ability. Since it does not fossilize, language scientists instead need to rely on indirect evidence to gain some insight about how it evolved in our clade. Language is ultimately the product of neural activity and so skulls and brain endocasts are objects of particular interest. Because of the intimate cross-talk between the skull and the brain, we expect that the analysis of the hominin skull shape and growth pattern can inform how the brains of extinct hominin species were organized and how they developed during growth; factors which impact cognition (Zollikofer & Ponce de León, 2013). Language abilities in Neanderthals are a controversial issue (Johansson, 2015). The skull of anatomically-modern humans (AMH) differs both in size and shape from the skull of

Neanderthals (Bruner, 2004). However, it is still not clear if calvarian differences correlate with endocast differences, and hence with brain function (Mounier *et al.*, 2016), or if there is an AMH-specific brain developmental path (see Ponce de León *et al.*, 2016; Williams & Cofran, 2016). Comparisons with extant primates are helping to refine our understanding of these differences (Pearce *et al.*, 2013). Overall, gross differences in brain structure and function between hominin species have been used to infer differences in cognitive abilities important for language processing (Boeckx & Benítez-Burraco, 2014a).

In neighboring domains, neuronal oscillatory activity has been acknowledged as the basis of multiple cognitive and behavioural processes (Buzsáki, 2006). In our previous work we have argued that computational operations of language can be decomposed into generic processes

that can be implemented via neural oscillations (Murphy, 2015a,b) and that certain multiplexing algorithms responsible for the construction of hierarchical phrase structures appear to be species-specific (Murphy, 2016a). We have also argued that neural oscillations provide a more reliable explanatory level for investigating linguistic computation than the (correlational) examination of nerve tracks and brain regions involved in language processing; not just in neurotypical language processing, but in cognitive conditions that are human-specific and entail language dysfunction (Benítez-Burraco & Murphy, 2016; Murphy & Benítez-Burraco, 2016). Meyer (2017) reviews a range of experimental evidence suggesting that oscillations play a causal role in language comprehension, from the chunking (δ) and storage (α) of phrasal units, to the prediction of upcoming syntactic material (β) and the unification of this material into a coherent semantic structure (γ). Beaudet (2017) points out that ‘[s]peech capacity cannot be appropriately inferred only from the cerebral condition, therefore hypotheses aiming at reconstructing the timing and mode of emergence of language in the hominin lineage should seek to combine various lines of evidence’ – with such lines of evidence including, we hope, links between genetics and oscillatory brain activity.

While we cannot track the oscillatory activity of extinct hominins, we believe that the next best solution to moving beyond this shortcoming consists in examining the coding regions and the expression patterns of the genes responsible for the brain’s oscillatory activity putatively responsible for human language. We have recently identified and functionally characterized a set of 48 genes that comprises the core of our *language oscillogenome*; that is, the set of genes responsible for basic aspects of the oscillatory activity relevant for language (Murphy & Benítez-Burraco, 2017). In this current contribution, we have looked for differences between Neanderthals and us in the coding regions of these genes, and in their methylation patterns, as well as for signals of positive selection in our species. Our ultimate aim is to inferring, from these genomic differences, differences in the brain activity important for language processing.

Evolutionary changes in the language oscillogenome

In order to achieve our objective, we first gathered via systematic literature review and database searches the available information concerning fixed changes in AMH proteins compared to Neanderthal proteins, genomic regions positively selected in AMHs after our split from Neanderthals, and differentially methylated regions (DMRs) in AMHs compared to Neanderthals. We have extensively built on Green *et al.*’s (2010) seminal characterization of the Neanderthal genome, Prüfer *et al.*’s (2013) description of the Altai Neanderthal genome, and Gokhman *et al.*’s (2014, 2017) DNA methylation maps of modern humans and extinct hominins. We then looked for our candidates in these papers and databases. As noted in Murphy and Benítez-Burraco (2017), our candidates for the language oscillogenome fulfil three criteria: i) they are associated with language disorders (developmental dyslexia and/or specific language impairment) and/or language dysfunction in cognitive disorders entailing language deficits (schizophrenia and/or autism spectrum disorder), ii) they play a role in brain rhythmicity and/or are candidates for conditions entailing brain dysrhythmias, like epilepsy; and iii) gene-oscillations-language links can be confidently established for them. Our findings are summarized in Table 1.

Among the 48 genes we highlighted in our previous work as part of the shared signature of abnormal brain oscillations associated with language deficits and core components of the language oscillogenome, 11 of these exhibit some difference between AMHs and Neanderthals. Two proteins bear fixed changes in one position in AMHs, although these changes are not expected to impact significantly on their structure and function: CNTNAP2 exhibits a Ile345Val change (reported as benign according to PolyPhen, and as tolerated (0.79) according to SIFT), whereas FMR1 shows a nearly fixed Ser145Ala change (frequency of 0.98 in AMHs) (classified as benign according to PolyPhen, and as tolerated (1) according to SIFT). More

Tab. 1 - AMH-specific changes in genes involved in brain oscillations important for languages processing. The information about fixed aminoacid changes in the protein (column 1) derives from several sources, as noted in the main text. Evidence of positive selection (column 2) is from Green *et al.* (2010). The data about DMRs in AMH bone (column 3) are from Gokhman *et al.* 2017 (extended data Table 2). The data about transcription factors enriched in AMH DMRs (column 4) is from Gokhman *et al.* 2014 (Table S5). The information about the oscillomic/oscillopathic features of the gene is from Murphy & Benítez-Burraco, 2018.

GENE	FIXED AA CHANGE IN AMHS	POSITIVELY SELECTED IN AMHS	DIFFERENTIALLY METHYLATED IN AMH SKELETAL SAMPLES	ENRICHED IN AMH DMRS	OSCILLOMIC/OSCILLOPATHIC FEATURES
<i>AUTS2</i>		✓	↑(body gene)		Epilepsy
<i>CACNA1C</i>			↑ (body gene)		β, γ
<i>CNTNAP2</i>	✓				α
<i>COL4A2</i>			↑(body gene)		Epilepsy
<i>COMT</i>			↑(downstream the gene)		α
<i>DYRK1A</i>		✓			Inhibition of neural activity
<i>EGR1</i>				✓	Epilepsy
<i>ELP4</i>		✓			High amplitude centrotemporal sharp waves
<i>FMR1</i>	✓				θ, γ
<i>FOXP1</i>		✓	↑(body gene)		Epileptiform discharges
<i>ROBO1</i>		✓			Epilepsy

interestingly, 5 of our candidates are found within the regions that exhibit the strongest signals of positive selection in AMHs compared to Neanderthals, according to Green *et al.* (2010): *AUTS2*, *DYRK1A*, *ELP4*, *FOXP1*, and *ROBO1*. These positively selected regions contain numerous binding sites for transcription factors (Supplementary file, S1), which might have been affected by AMH-specific changes. Nonetheless, the most direct evidence of putative differences in the expression pattern of genes important for language under an oscillomic lens derive from the methylation maps of the AMH and the Neanderthal genomes obtained by Gokhman and colleagues (2014, 2017). We have found that several regions within the body gene of 4 of our candidates are hypermethylated in skeletal samples of AMHs compared to Neanderthals (*AUTS2*, *CACNA1C*, *COL4A2*,

and *FOXP1*). Likewise, one hypermethylated region is found downstream one of our candidates (*COMT*) (Table 1 and supplementary file, S2). All these regions contain binding sites for several transcription factors of interest, like FOXA1, FOXA2, EP300, CBPB, MEF2A, FOXP2, FOXA2, CEBPB (Supplementary file, S2), which we have highlighted as important for language evolution, particularly, for changes related to the externalization of our linguistic thoughts (speech) (Boeckx & Benítez-Burraco, 2014b, Benítez-Burraco & Boeckx 2015). Finally, we have found that *EGR1* is one of the enriched transcription factors in AMH DMRs.

Although methylation is commonly associated with promoter regulation, it is also a regulatory mechanism of gene expression when affecting splice sites, coding regions, binding sites for transcription factors, and distal regulatory

elements, like enhancers, silencers or insulators (Jones *et al.*, 2012; Miller & Grant, 2013; Lev Maor *et al.*, 2015). Accordingly, from the above differences in their methylation patterns, we might expect differences between Neanderthals and AMHs in the expression levels/patterns of several candidates for the language oscillome, namely, *AUTS2*, *CACNA1C*, *COL4A2*, *COMT*, *EGR1*, and *FOXP1*. Obviously, it is still to be determined whether the identified DMRs in skeletal methylation maps exist between the brains of AMHs and Neanderthals too. However, this might be the case (see Gokhman *et al.*, 2016 for discussion).

Interestingly, duplications and deletions of all these genes, purportedly resulting in abnormal high or low levels of the protein, respectively, result in cognitive and language deficits (Appendix, Supplementary file, S4).

AUTS2 is involved in cytoskeletal regulation to control neuronal migration and neurite extension, but nuclear *AUTS2* also controls transcription of several genes (Oksenberg *et al.*, 2014; Hori & Hoshino, 2017). The gene has been associated with differential processing speeds (Luciano *et al.*, 2011) and is a candidate for epilepsy (Mefford *et al.*, 2010). *CACNA1C* encodes the alpha 1C subunit of the Cav1.2 voltage-dependent L-type calcium channel, involved in β and γ generation (Kumar *et al.*, 2015). Pathogenic variants of *CACNA1C* have been associated with low performance scores in semantic verbal fluency tasks in schizophrenics (Krug *et al.*, 2010), as well as with childhood-onset epilepsy, executive dysfunction, intellectual disability, and/or autism spectrum disorder (Damaj *et al.*, 2015). *COL4A2* is a candidate for dyslexia, but it has also been associated with epilepsy and severe developmental delay (Giorgio *et al.*, 2015; Smigiel *et al.*, 2016). *COMT*, which encodes a catechol-O-methyltransferase that catalyzes the O-methylation of neurotransmitters like dopamine, epinephrine, and norepinephrine, has been regularly associated with schizophrenia, and particularly, with language dysfunction in this condition (Egan *et al.*, 2001; Malhotra *et al.*, 2002). The gene has been also associated with

reading abilities in the neurotypical population (Landi *et al.*, 2013), as well as with language performance and processing, and language acquisition, particularly to verbal fluency (Krug *et al.*, 2009; Soeiro-De-Souza *et al.*, 2013; Sugiura *et al.*, 2017). Low activity levels in *COMT* has been associated with low voltage α (Enoch *et al.*, 2003). Intriguingly, AMHs exhibit a hypomethylated region inside the body of one gene important for *COMT* function, namely *MSRA*, compared to Neanderthals (Gokhman *et al.*, 2014, Table S2). *EGR1* encodes a transcription factor that contributes to regulate neural plasticity and memory consolidation (Veyrac *et al.*, 2014). In human epileptic foci, *EGR1* expression correlates with the frequency, amplitude and area of the interictal spikes, a hallmark of epileptic neocortex (Rakhade *et al.*, 2007). Interestingly, *EGR1* is a target of *FOXP2*, the renowned “language gene” (Konopka *et al.*, 2009), whereas *EGR1* downregulates *PLAUR* (Matsunoshita *et al.*, 2011), another of *FOXP2*’s targets (Roll *et al.*, 2010). *FMRP1* is the main candidate for Fragile X syndrome (Kaufmann *et al.*, 2004; Smith *et al.*, 2012). *Fmr1*-knockout mice exhibit abnormal patterns of coupling between θ and γ oscillations in the CA1 area of the hippocampus, enhanced mGluR5 signalling resulting in altered neocortical rhythmic activity, and abnormally weak changes during tasks involving cognitive challenge (Hays *et al.*, 2011; Radwan *et al.*, 2016). *FOXP1* interacts with *FOXP2* to form functional heterodimers and *FOXP1* haplo-insufficiency has been associated with epileptiform discharges, speech delay, and motor delay (Carr *et al.*, 2010).

Although in this paper we are interested in changes that are specific to AMHs and that allegedly contributed to the emergence of our species-specific pattern of brain oscillations, we wish to note that several of our candidates are differentially methylated in Neanderthals compared to other hominins. These differences might help account as well for their distinctive cognitive abilities, as discussed below. Accordingly, the body gene of *COMT* and *KANSL1* are differentially hypomethylated in bone samples of Neanderthals compared to AMHs. Likewise,

one region downstream of *ERG1* is found hypomethylated in Neanderthals (Supplementary file, S3). We have already discussed both *COMT* and *ERG1*. Regarding *KANSL1*, it encodes a component of the NSL1 complex, which functions as a regulator of gene transcription and chromatin organization, and is a candidate for Koolen-de Vries syndrome, which is associated with epilepsy, developmental delay, and moderate intellectual disability, that mostly affects expressive language abilities (Koolen *et al.*, 2016).

The language network in Neanderthals: paleoneurology and oscillomics

Although very indirect due to the lack of actual evidence in the case of the former, comparative studies of Neanderthal and AMH cognition are vital if we are to hone our understanding of cognitive features unique to our species. As Mounier *et al.* (2016, p. 22) note, ‘the use of palaeoneurology to infer phylogenies of our genus is rare’. An emerging consensus indicates that while the skull size and shape of AMHs and Neanderthals are comparable, their internal neural organisation was in a number of respects distinct (Pearce *et al.*, 2013). For instance, Neanderthals lived at much higher latitudes than AMHs and would have likely had larger eyes and, consequently, larger visual cortices (Pearce & Dunbar, 2012; Balzeau *et al.*, 2012). Over the past couple of decades, digital anatomy and computed morphometrics have led to a seismic shift in functional craniology, exposing the generally plesiomorph structure of the Neanderthal braincase (Weber & Bookstein, 2011; Bruner, 2014; Williams & Cofran, 2016). Figure 1 summarizes the major findings of this research. Naturally, not every neuroanatomical difference between Neanderthals and AMHs will be relevant to evolutionary linguistics, but we will conservatively review those differences which strike us as most significant. Notably, Neanderthals display upper parietal bulging which approximates the AMH shape but crucially lacks what Bruner (2014, p.

123) describes as ‘the overall bulging of the parietal surface characterizing the modern human brain globularity, in which the parietal proportions also enlarge in the longitudinal and vertical directions’. As we have already noted (Murphy & Benítez-Burraco, 2016), the globularity of the AMH braincase appears to have had direct consequences for the efficiency of oscillatory couplings across the cortex and subcortex which we have, in turn, used to derive the cross-modular nature of linguistic representations. Bruner (2014, p. 123) concludes that the type of parietal changes exhibited by the AMH brain likely resulted in modifications to visuo-spatial integration and ‘mental experiment capabilities’, which we interpret as recursively generated symbolic manipulations or what Tattersall (2017, p. 65) regards as ‘imagination’. Likewise, due to the expanded occipital lobe and visual cortices in Neanderthals, and due to the role of this region in α -inhibition vital to the coordination of cross-cortical feature binding (Murphy, 2016a), it is a possibility that Neanderthals were capable of executing a degree of control over representation integration not seen in other primates close to AMHs, but nevertheless still reduced in cross-cortical scope relative to AMHs. Further support for this hypothesis comes from Appendix: *COMT* exhibits hypermethylation in AMHs compared to Neanderthals, and since low-voltage α has been associated with low activity levels in the gene (Enoch *et al.*, 2003), with *COMT* additionally being associated with language performance and verbal fluency (Krug *et al.*, 2009; Soeiro-De-Souza *et al.*, 2013), it is reasonable to assume that this hypermethylation contributes to the type of α -inhibition required to control inter-modular featural integration.

It was argued in Murphy (2016b) that a range of monkeys (putty-nosed monkeys, Campbell’s monkeys, King Colobus monkeys, and New World monkeys) seem capable only of single-instance serial order concatenation and not hierarchical phrasal construction. Drawing on much neuroethological data, the likely oscillatory basis of this capacity was explored. It was proposed that phase-amplitude coupling




Fig. 1 - Paleoneurological characters of Neanderthals from a replica of the skull and endocasts of Saccopastore 1. White arrows indicate derived traits while black arrows indicate plesiomorph traits, which impose constraints as a result of large brain size (reproduced from Bruner, 2014, p. 123). The colour version of this figure is available at the JASs website.

between parahippocampal θ and cross-cortical γ was responsible for constructing individual call representations, and that primarily basal ganglia β increases signalled the maintenance of such calls in memory (Murphy, 2016b).

At this point, we need to distinguish between the possibility that Neanderthals were capable of speech from the possibility that they were capable of internally generating hierarchically organised sets of mental representations, as in natural language (as assumed in much work in generative linguistics; Chomsky *et al.* 2018). Neanderthals may have been capable of speech similar to AMHs due to the hard portion of the hyoid apparatus typically being preserved and their aural ability to be sensitive to frequencies (up to 5Hz) relevant for speech (Martinez *et al.*, 2004). But what about the generation of hierarchically organised representations, the core of natural language (Chomsky *et al.* 2018)? We believe that since current work is revealing an oscillatory code for language, and since the

molecular basis of this code is in turn becoming amenable to reconstruction (Murphy & Benítez-Burraco, 2017), the most reliable existing method of examining whether Neanderthals had anything resembling a proto-language is to draw inferences from emerging genetic databases. For instance, while Neanderthals may have been sensitive to language-relevant frequencies, the question we are interested in is whether neural assemblies internally oscillating at the same frequencies were used in the service of constructing hierarchically organized representations. Relatedly, Mithen (2014) reviews the evidence for cognitive similarities and differences in AMHs and Neanderthals and concludes, for example, that while pigment use likely indicates a complex social and emotional life, it does not provide evidence for symbolic thought.

It is likely that the human-specific oscillatory profile documented in Murphy (2016a), differing from the Neanderthal oscillome proposed here, is responsible for Mithen's (1996, 2014)

seminal distinction between the domain-specific Neanderthal mind (in which distinct core knowledge systems are largely isolated from each other) and the ‘cognitive fluidity’ of the AMH mind. Without this fluidity, integrating representations from distinct domains, Neanderthals ‘could never develop such [complex social] relationships and initiate a process of domestication’ (Mithen, 2014, p. 12; concerning domestication, see also Benítez-Burraco *et al.*, 2016). More specifically, we would like to suggest that the distinct θ - γ code we predicted for the Neanderthal brain might also explain its presumably limited working memory (Wynn & Coolidge, 2012), given the crucial role that coupling between these rhythms appears to have in working memory operations across a number of modalities (Murphy, 2016a; Schomburg *et al.*, 2014; Vosskuhl *et al.*, 2015). A widening in the anterior fossa in Neanderthals (Fig. 1) is associated with an expansion in Broca’s area, argued in Murphy (2016a) to be implicated in language-relevant memory buffers (see also Rogalsky *et al.*, 2008 for a discussion of earlier proposals concerning Broca’s area in expanding working memory capacities). While linguistic representations may well have been assigned a special memory buffer (phonological loop) in Brodmann area 44, a part of Broca’s area (henceforth, BA 44), the computations associated with Merge, the core combinatorial operation in natural language that combines elementary linguistic units (Chomsky, 1995), are certainly well distributed. In fact, even this perspective might not be wholly accurate: Along with BA 44 being a memory buffer (see Badre & Wagner 2007 for seminal discussion), it may also serve to select between competing syntactic interpretations, both storing and comparing representations (Novick *et al.*, 2010). We confess that this perspective differs from the one articulated prominently by Friederici *et al.* (2017), who view parts of Broca’s area as playing a much more central role in phrase structure generation. Taken alongside the oscillogenic evidence presented above – that Neanderthals may have exhibited reduced cross-frequency coupling between θ and γ due to differences in *CACNA1C* and *ELP4* expression

– we should also conclude that the Neanderthal brain was likely incapable of exploiting this expanded Broca’s area, at least in the service of maintaining sets of symbolic representations. As with the partly globular parietal modifications discussed above, this appears to be another major feature of the Neanderthal brain which begins to approximate neuroanatomical characteristics that we hypothesize are required for core aspects of AMH syntax, but which ultimately displays other features which are likely incompatible with it. While the currently proposed oscillatory profile of Neanderthals does not necessarily exclude elements deemed necessary for phrasal construction (involving δ phase-amplitude coupling with θ and β , for instance; see Murphy, 2016b), the severe limitations which seem to have been imposed on it (most notably, working memory constraints and non-optimal cross-frequency couplings) suggests that even if Neanderthals did have core features of the neural code for AMH syntax there would have been a number of obstacles to implementing it.

Returning to Table 1, *CACNA1C*, which is differentially methylated in AMHs, contributes to β and γ generation, and has been correlated with semantic verbal fluency, may set AMHs apart from Neanderthals by contributing to speech-related fluency through β (implicated in storing ongoing speech representations in memory) and γ (involved in the initial construction of discourse representations, as discussed above). Relatedly, the abnormal dendritic spine morphology associated with *FMRI*, along with the decreased β power (Appendix), strengthens this notion.

With respect to the evolutionary implications, we conclude that the documented differences in the oscillatory profiles of humans and Neanderthals not only explain why we have language and Neanderthals likely did not, but they also shed some light on language-related cognitive differences, in particular concerning working memory limitations. We remain silent on the issue of the type of word-like representations Neanderthals may have possessed, and have focused purely on computational capacity,

although it is worth noting that the evidence for Neanderthal symbolic thought at least suggests that it is their processing capabilities which distinguishes them from us. Our view is therefore somewhat similar to the mainstream generative position on language evolution (insofar as this particular topic goes, that is), although our views on neural reorganization in our lineage differ substantially from, for instance, Berwick & Chomsky's (2016, 2017) or Friederici *et al.*'s (2017) positions (however the position articulated by these authors is also currently compatible with the data we present, given their focus on the computational novelty of Merge). Be that as it may, it should also be stressed that the current contribution constitutes highly speculative work, with the differences between Neanderthals and humans being a hotly debated and controversial topic. We hope at least to have opened up new avenues for exploring these differences, even if our present conclusions are soon revealed to be inaccurate and premature.

Conclusion

Overall, it is plausible that the oscillatory differences we have documented between AMHs and Neanderthals are connected to the major cultural and technological novelties which occurred towards the end of the Middle Pleistocene and which distinguish AMHs from our closest relatives. Expanding our molecular understanding of the Neanderthal brain can contribute to expanding the line of inquiry into what we have here called paleo-oscillomics. While it is hard to imagine anything in the fossil record shedding light on the *representational* basis of the Neanderthal brain (Berwick *et al.*, 2013), reconstructing the Neanderthal oscillome can at least reveal aspects of its *computational* capacity.

Acknowledgments

This work was supported by an Economic and Social Research Council scholarship (1474910) to

EM, and by funds from the Spanish Ministry of Economy and Competitiveness (grant number FFI-2013-43823-P) to ABB. The authors wish express their gratitude to David Gokhman for sharing his data on methylation maps of Neanderthals and humans, and for kindly reading an earlier draft of the paper.

Author contribution

ABB wrote Section 1-2. EM wrote Section 3-4.

References

- Badre D. & Wagner A.D. 2007. Left ventrolateral prefrontal cortex and the cognitive control of memory. *Neuropsychologia* 45: 2883-2901.
- Balzeau A., Holloway R. L. & Grimaud-Hervé D. 2012. Variations and asymmetries in regional brain surface in the genus *Homo*. *J. Hum. Evol.*, 62: 696-706.
- Benítez-Burraco A. & Boeckx C. 2015. Possible functional links among brain- and skull-related genes selected in modern humans. *Front. Psychol.*, 6: 794.
- Benítez-Burraco A. & Murphy E. 2016. The oscillopathic nature of language deficits in autism: from genes to language evolution. *Front. Hum. Neurosci.*, 10: 120.
- Benítez-Burraco A., Lattanzi W. & Murphy E. 2016. Language impairments in ASD resulting from a failed domestication of the human brain. *Front. Neurosci.*, 10: 373.
- Berwick R. C. & Chomsky N. *Why Only Us: Language and Evolution*. MIT Press Cambridge, MA.
- Berwick R. C. & Chomsky N. 2017. Why only us: recent questions and answers. *J. Neurolinguistics*, 43: 166-177.
- Berwick R. C., Hauser M. D. & Tattersall I. 2013. Neanderthal language? Just-so stories take center stage. *Front. Psychol.*, 4: 671.
- Beaudet A. 2017. The emergence of language in the hominin lineage: perspectives from fossil endocasts. *Front. Hum. Neurosci.*, 11: 427.

- Boeckx C. & Benítez-Burraco A. 2014a. The shape of the human language-ready brain. *Front. Psychol.*, 5: 282
- Boeckx C. & Benítez-Burraco A. 2014b. Globularity and language-readiness: Generating new predictions by expanding the set of genes of interest. *Front. Psychol.*, 5: 1324.
- Bruner E. 2004. Geometric morphometrics and paleoneurology: brain shape evolution in the genus *Homo*. *J. Hum. Evol.*, 47: 279-303.
- Bruner E. 2014. Functional craniology, human evolution, and anatomical constraints in the Neanderthal braincase. In Akazawa T., Ogiwara N., Tanabe H. C. & Terashima H. (eds): *Dynamics of Learning in Neanderthals and Modern Humans*, vol. II. *Cognitive and Physical Perspectives. Replacement of Neanderthals by Modern Humans Series*, pp. 121-129. Springer Japan, Tokyo.
- Buzsáki G. 2006. *Rhythms of the Brain*. Oxford University Press, New York.
- Carr C. W., Moreno-De-Luca D., Parker C., Zimmerman H. H., Ledbetter N., Martin C. L. et al. 2010. Chiari I malformation, delayed gross motor skills, severe speech delay, and epileptiform discharges in a child with FOXP1 haploinsufficiency. *Eur. J. Hum. Genet.*, 18: 1216-20.
- Chomsky N. 1995. *The Minimalist Program*. MIT Press, Cambridge.
- Chomsky N., Gallego Á. J. & Ott D. 2018. Generative grammar and the faculty of language: insights, questions, and challenges. In Gallego Á. J. & Ott D. (eds): *Generative Syntax: Questions, Crossroads, and Challenges*. Special issue of *Catalan Journal of Linguistics*.
- Damaj L., Lupien-Meilleur A., Lortie A., Riou É., Ospina L. H., Gagnon L. et al. 2015. CACNA1A haploinsufficiency causes cognitive impairment, autism and epileptic encephalopathy with mild cerebellar symptoms. *Eur. J. Hum. Genet.*, 23: 1505-12.
- Egan M. F., Goldberg T. E., Kolachana B. S., Callicott J. H., Mazzanti C. M., Straub R. E. et al. 2001. Effect of COMT Val108/158 Met genotype on frontal lobe function and risk for schizophrenia. *Proc. Natl. Acad. Sci. USA*, 98: 6917-6922.
- Enoch M. A., Xu K., Ferro E., Harris C. R. & Goldman D. 2003. Genetic origins of anxiety in women: a role for a functional catechol-O-methyltransferase polymorphism. *Psychiatr. Genet.*, 13: 33-41.
- Friederici A. D., Chomsky N., Berwick R. C., Moro A. & Bolhuis J. J. 2017. Language, mind and brain. *Nat. Hum. Behav.*, 1: 713-722.
- Giorgio E., Vaula G., Bosco G., Giacone S., Mancini C., Calcia A. et al. 2015. Two families with novel missense mutations in COL4A1: When diagnosis can be missed. *J. Neurol. Sci.*, 352: 99-104
- Gokhman D., Lavi E., Prüfer K., Fraga M. F., Riancho J. A., Kelso J. et al. 2014. Reconstructing the DNA methylation maps of the Neandertal and the Denisovan. *Science*, 344: 523-527.
- Gokhman D., Meshorer E. & Carmel L. 2016. Epigenetics: it's getting old. Past meets future in paleoepigenetics. *Trends Ecol. Evol.*, 31: 290-300.
- Gokhman D., Agranat-Tamir L., Housman G., Nissim-Rafinia M., Nieves-Colón M., Gu H. et al. 2017. Recent regulatory changes shaped human facial and vocal anatomy. *bioRxiv*, 106955; doi: <https://doi.org/10.1101/106955>
- Green R. E., Krause J., Briggs A. W., Maricic T., Stenzel U., Kircher M. et al. 2010. A draft sequence of the Neandertal genome. *Science*, 328: 710-722.
- Hays S. A., Huber K. M. & Gibson J. R. 2011. Altered neocortical rhythmic activity states in Fmr1 KO mice are due to enhanced mGluR5 signaling and involve changes in excitatory circuitry. *J. Neurosci.*, 31: 14223-14234.
- Hori K. & Hoshino M. 2017. Neuronal migration and AUTS2 syndrome. *Brain Sci.*, doi: 10.3390/brainsci7050054.
- Johansson S. 2015. Language abilities in Neanderthals. *Ann. Rev. Ling.*, 1: 311-332.
- Jones P. A. 2012. Functions of DNA methylation: islands, start sites, gene bodies and beyond. *Nat. Rev. Genet.*, 13: 484-492.
- Kaufmann W. E., Cortell R., Kau A. S., Bukelis I., Tierney E., Gray R. M. et al. 2004. Autism spectrum disorder in fragile X syndrome:

- Communication, social interaction, and specific behaviors. *Am. J. Med. Genet. A*, 129: 225–234
- Konopka G., Bomar J. M., Winden K., Coppola G., Jonsson Z. O., Gao F. *et al.* 2009. Human-specific transcriptional regulation of CNS development genes by FOXP2. *Nature*, 462: 213–217.
- Koolen D. A., Pfundt R., Linda K., Beunders G., Veenstra-Knol H. E., Conta J. H. *et al.* 2016. The Koolen-de Vries syndrome: a phenotypic comparison of patients with a 17q21.31 microdeletion versus a KANSL1 sequence variant. *Eur. J. Hum. Genet.*, 24: 652–659
- Krug A., Markov V., Sheldrick A., Krach S., Jansen A., Zerres K. *et al.* 2009. The effect of the COMT val(158)met polymorphism on neural correlates of semantic verbal fluency. *Eur. Arch. Psychiatry Clin. Neurosci.*, 259: 459–65.
- Krug A., Nieratschker V., Markov V., Krach S., Jansen A., Zerres K., *et al.* 2010. Effect of CACNA1C rs1006737 on neural correlates of verbal fluency in healthy individuals. *Neuroimage*, 49: 1831–1836.
- Kumar A., Sundaram S. K., Sivaswamy L., Behen M. E., Makki M. I., Ager J. *et al.* 2010. Alterations in frontal lobe tracts and corpus callosum in young children with autism spectrum disorder. *Cereb. Cortex*, 20: 2103–2113.
- Kumar D., Dedic N., Flachsamm C., Voulé S., Deussing J. M., & Kimura M. 2015. Cacna1c (Cav1.2) modulates electroencephalographic rhythm and rapid eye movement sleep recovery. *Sleep*, 38: 1371–1380.
- Landi N., Frost S. J., Mendel W. E., Preston J. L., Jacobsen L. K. *et al.* 2013. The COMT Val/Met polymorphism is associated with reading-related skills and consistent patterns of functional neural activation. *Dev. Sci.*, 16: 13–23.
- Lev Maor G., Yearim A. & Ast G. 2015. The alternative role of DNA methylation in splicing regulation. *Trends Genet.* 31: 274–280.
- Luciano M., Hansell N. K., Lahti J., Davies G., Medland S. E., Räikkönen K. *et al.* 2011. Whole genome association scan for genetic polymorphisms influencing information processing speed. *Biol. Psychol.*, 86: 193–202
- Malhotra A. K., Kestler L. J., Mazzanti C., Bates J. A., Goldberg T. & Goldman D. 2002. A functional polymorphism in the COMT gene and performance on a test of prefrontal cognition. *Am. J. Psychiatry*, 159: 652–654
- Martinez I., Rosa M., Arsuaga J.-L., Jarabo P., Quam R., Lorenzo C. *et al.* 2004. Auditory capacities in Middle Pleistocene humans from the Sierra de Atapuerca in Spain. *Proc. Natl. Acad. Sci. USA*, 101: 9976–9981.
- Matsunoshita Y., Ijiri K., Ishidou Y., Nagano S., Yamamoto T., Nagao H. *et al.* 2011. Suppression of osteosarcoma cell invasion by chemotherapy is mediated by urokinase plasminogen activator activity via up-regulation of EGR1. *PLoS One*, 6: e16234.
- Mefford H. C., Muhle H., Ostertag P., von Spiczak S., Buysse K., Baker C. *et al.* 2010. Genome-wide copy number variation in epilepsy: novel susceptibility loci in idiopathic generalized and focal epilepsies. *PLoS Genet.*, 6, e1000962.
- Meyer L. 2017. The neural oscillations of speech processing and language comprehension: state of the art and emerging mechanisms. *Eur. J. Neurosci.*, doi:10.1111/ejn.13748.
- Miller J. L. & Grant P. A. 2013. The role of DNA methylation and histone modifications in transcriptional regulation in humans. *Subcell Biochem.*, 61:289–317.
- Mithen S. 1996. *The Prehistory of the Mind: A Search for the Origins of Art, Science and Religion*. Thames & Hudson, London.
- Mithen S. 2014. The cognition of *Homo Neanderthalensis* and *H. sapiens*: does the use of pigment necessarily imply symbolic thought? In Akazawa T., Ogihara N., Tanabe H. C. & Terashima H. (eds): *Dynamics of Learning in Neanderthals and Modern Humans, vol. II: Cognitive and Physical Perspectives*, pp. 7–16. Replacement of Neanderthals by Modern Humans Series. Springer Japan, Tokyo.
- Mounier A., Balzeau A., Caparros M. & Grimaud-Hervé D. 2016. Brain, calvarium, cladistics: A new approach to an old question, who are modern humans and Neandertals? *J. Hum. Evol.*, 92: 22–36.
- Murphy E. 2015a. Labels, cognomes and cyclic computation: an ethological perspective. *Front. Psychol.*, 6: 715.

- Murphy E. 2015b. The brain dynamics of linguistic computation. *Front. Psychol.*, 6: 1515.
- Murphy E. 2016a. A theta-gamma neural code for feature set composition with phase-entrained delta nestings. *UCL Working Papers in Linguistics*, 28: 1-23.
- Murphy E. 2016b. Evolutionary monkey oscillomics: generating linking hypotheses from preserved brain rhythms. *Theor. Ling.*, 42: 117-137.
- Murphy E. & Benítez-Burraco A. 2016. Bridging the gap between genes and language deficits in schizophrenia: an oscillopathic approach. *Front. Hum. Neurosci.*, 10: 422.
- Murphy E. & Benítez-Burraco A. 2018. Toward the language oscillogenome. *Front. Psychol.*, 9: 1999.
- Novick J. M., Trueswell J.C. & Thompson-Schill S. L. 2010. Broca's area and language processing: evidence for the cognitive control connection. *Lang. Linguist. Compass*, 4: 906-924.
- Oksenberg N., Haliburton G. D., Eckalbar W. L., Oren I., Nishizaki S., Murphy K. *et al.* 2014. Genome-wide distribution of *Auts2* binding localizes with active neurodevelopmental genes. *Transl. Psychiatry*, 4: e431.
- Pearce E. & Dunbar R. I. M. 2012. Latitudinal variation in light levels drives human visual system size. *Biol. Lett.*, 8: 90-93.
- Pearce E., Stringer C. & Dunbar R. I. M. 2013. New insights into differences in brain organization between Neanderthals and anatomically modern humans. *Proc. Biol. Sci.*, 280: 20130168.
- Ponce de León M. S., Bienvenu T., Akazawa T. & Zollikofer C. P. 2016. Brain development is similar in Neanderthals and modern humans. *Curr. Biol.*, 26: R665-666.
- Prüfer K., Racimo F., Patterson N., Jay F., Sankararaman S., Sawyer S. *et al.* 2014. The complete genome sequence of a Neanderthal from the Altai Mountains. *Nature*, 505: 43-49.
- Radwan B., Dvorak D. & Fenton A.A. 2016. Impaired cognitive discrimination and discoordination of coupled theta-gamma oscillations in *Fmr1* knockout mice. *Neurobiol. Dis.*, 88: 125-138.
- Rakhade S. N., Shah A. K., Agarwal R., Yao B., Asano E. & Loeb J. A. 2007. Activity-dependent gene expression correlates with interictal spiking in human neocortical epilepsy. *Epilepsia*, 48: 86-95.
- Rogalsky C., Matchin W. & Hickok G. 2008. Broca's area, sentence comprehension, and working memory: an fMRI study. *Front. Hum. Neurosci.*, 2: 14.
- Roll P., Vernes S. C., Bruneau N., Cillario J., Ponsole-Lenfant M., Massacrier A. *et al.* 2010. Molecular networks implicated in speech-related disorders: FOXP2 regulates the SRPX2/uPAR complex. *Hum. Mol. Genet.*, 19: 4848-4860.
- Schomburg E. W., Fernandez-Ruiz A., Mizuseki K., Berenyi A., Anastassiou C. A., Koch C. *et al.* 2014. Theta phase segregation of input-specific gamma patterns in entorhinal-hippocampal networks. *Neuron*, 84: 470-485.
- Smigiel R., Cabala M., Jakubiak A., Koder H., Sasiadek M. J., Matsumoto N. *et al.*, 2016. Novel COL4A1 mutation in an infant with severe dysmorphic syndrome with schizencephaly, periventricular calcifications, and cataract resembling congenital infection. *Birth Defects Res. Part A Clin. Mol. Teratol.*, 106: 304-307. doi:10.1002/bdra.23488.
- Smith L. E., Barker E. T., Seltzer M. M., Abbeduto L. & Greenberg J. S. 2012. Behavioral phenotype of fragile X syndrome in adolescence and adulthood. *Am. J. Intellect. Dev. Disabil.*, 117: 1-17.
- Soeiro-De-Souza M. G., Bio D. S., David D. P., Missio G., Lima B., Fernandes F. *et al.* 2013. Gender effects of the COMT Val 158 Met genotype on verbal fluency in healthy adults. *Mol. Med. Rep.*, 8: 837-844.
- Sugiura L., Toyota T., Matsuba-Kurita H., Iwayama Y., Mazuka R., Yoshikawa T. *et al.* 2017. Age-dependent effects of Catechol-O-Methyltransferase (COMT) gene Val158Met polymorphism on language function in developing children. *Cereb. Cortex*, 27: 104-116.
- Tattersall I. 2017. How can we detect when language emerged? *Psychon. Bull. Rev.*, 24: 64-67.

- Veyrac A., Besnard A., Caboche J., Davis S. & Laroche S. 2014. The transcription factor Zif268/Egr1, brain plasticity, and memory. *Prog. Mol. Biol. Transl. Sci.*, 122: 89-129.
- Vosskuhl J., Huster R. J. & Herrmann C. S. 2015. Increase in short-term memory capacity induced by down-regulating individual theta frequency via transcranial alternating current stimulation. *Front. Hum. Neurosci.*, 9: 257.
- Weber G. W. & Bookstein F. L. 2011. *Virtual Anthropology: A Guide to a New Interdisciplinary Field*. Springer, Wien.
- Williams F. L. & Cofran Z. 2016. Postnatal craniofacial ontogeny in neandertals and modern humans. *Am. J. Phys. Anthropol.*, 159: 394-409.
- Wynn T. & Coolidge F. L. 2012. *How to Think Like a Neandertal*. Oxford University Press, Oxford.
- Zollikofer C. P. & Ponce de León M. S. 2013. Pandora's growing box: Inferring the evolution and development of hominin brains from endocasts. *Evol. Anthropol.*, 22: 20-33.

Editor, Giovanni Destro Bisol



This work is distributed under the terms of a Creative Commons Attribution-NonCommercial 4.0 Unported License <http://creativecommons.org/licenses/by-nc/4.0/>

Appendix - Abnormal brain, cognitive, and/or behavioural features associated with mutations in genes involved in brain oscillations important for language processing. The abnormal features associated to the genes were retrieved from the Phenotypes tool of Ensembl (<http://www.ensembl.org/index.html>). Abnormal features observed in humans (column 1) derive from several databases of diseases and phenotypes directly associated with a gene, including the Online Mendelian Inheritance in Man (OMIM) compendium, Orphanet, and DDG2P, as well as from several sources of diseases and traits associated with variants of interest, including COSMIC, ClinVar, dbGaP, EGA, GIAN, HGMD-Public, MAGIC, NHGRI-EBI GWAS catalog, and UniProt. The abnormal phenotypes linked to mutations of the orthologous genes in other species come from several databases, including the International Mouse Phenotyping Consortium and Europhenome (for mouse), ZFIN (for zebrafish), Rat Genome Database (for rat), Animal_QTLdb and Online Mendelian Inheritance in Animals. Because of biological differences between species, some animal phenotypes will not be easy to translate to our species. Column 3 contains the available information about abnormal brain, cognitive, and/or behavioural features associated in humans with deletions of the full gene or the promoter region of the gene, as provided by DECIPHER (<https://decipher.sanger.ac.uk/>). Column 4 displays the abnormal brain, cognitive, and/or behavioural traits associated to duplications of the full sequence of the genes of interest, as also provided by DECIPHER.

GENE	ABNORMAL BRAIN/COGNITIVE/BEHAVIOURAL FEATURES ASSOCIATED TO THE MUTATION OF THE GENE IN HUMANS	ABNORMAL BRAIN/COGNITIVE/BEHAVIOURAL FEATURES ASSOCIATED TO THE MUTATION OF ORTHOLOGUES OF THE GENE	ABNORMAL BRAIN/COGNITIVE/BEHAVIOURAL FEATURES ASSOCIATED TO DELETIONS OF THE FULL GENE OR THE PROMOTER REGION	ABNORMAL BRAIN/COGNITIVE/BEHAVIOURAL FEATURES ASSOCIATED TO DUPLICATION OF THE FULL GENE
<i>AUTS2</i>	Autism Mental retardation Intellectual disability Global developmental delay Abnormal response to serotonin reuptake inhibitors in non-psychotic unipolar depression	Alterations of forebrain neuron development Increased cell proliferation in the midbrain and the hindbrain Decreased size of the anterior region of the head Decreased number of cerebellar neurons Generalized epilepsy with febrile seizures	Global brain atrophy Moderate intellectual disability (DECIPHER patient 279043)	N/A
<i>CACNA1C</i>	Timothy syndrome Autism Schizophrenia Bipolar disorder Intellectual deficiency	Abnormal central nervous system synaptic transmission Abnormal nervous system physiology Abnormal learning/memory/conditioning	Autistic behavior (DECIPHER patient 290015)	N/A
<i>COL4A2</i>	Porencephaly Brain small vessel disease with hemorrhage	Cerebral hemorrhage Porencephaly Brain small vessel disease with hemorrhage	Behavioral abnormality (DECIPHER patient 278109)	Intellectual dissability (DECIPHER patient 273803)
<i>COMT</i>	22q11.2 Deletion Syndrome Susceptibility to schizophrenia	Abnormal increased prepulse inhibition Increased anxiety-related response Enhanced contextual conditioning behavior Impaired cued conditioning behavior Increased dopamine level Hyperactivity Abnormal spatial working memory	N/A	N/A

Appendix - Continued.

GENE	ABNORMAL BRAIN/ COGNITIVE/ BEHAVIOURAL FEATURES ASSOCIATED TO THE MUTATION OF THE GENE IN HUMANS	ABNORMAL BRAIN/COGNITIVE/ BEHAVIOURAL FEATURES ASSOCIATED TO THE MUTATION OF ORTHOLOGUES OF THE GENE	ABNORMAL BRAIN/ COGNITIVE/ BEHAVIOURAL FEATURES ASSOCIATED TO DELETIONS OF THE FULL GENE OR THE PROMOTER REGION	ABNORMAL BRAIN/ COGNITIVE/ BEHAVIOURAL FEATURES ASSOCIATED TO DUPLICATION OF THE FULL GENE
<i>EGR1</i>	Schizophrenia	Abnormal forebrain, midbrain, and hindbrain development Decreased dendrite morphogenesis Generalized epilepsy with febrile seizures	N/A	Abnormal emotion/ affect behavior Global developmental delay (DECIPHER patient 304609)
<i>FMR1</i>	Fragile X syndrome Intellectual disability Mental Competency	Abnormal cranial nerve position Disrupted sensory neuron axon guidance Abnormal axons of the lateral longitudinal fasciculus Disorganized hindbrain axons Increased synapse assembly of cholinergic neurons Abnormal dendritic spine morphology Decreased long-term synaptic potentiation Abnormal long term depression Decreased β wave amplitude Impaired learning and memory Impaired turning behavior	Global developmental delay (DECIPHER patient 284912)	N/A
<i>FOXP1</i>	Intellectual disability Severe speech delay Mental retardation with language impairment and with or without autistic features Attention deficit hyperactivity disorder	Cerebral visual impairment Intellectual disability Malformations of cortical development Autistic features	Abnormality of the face Abnormality of the nervous system Aggressive behavior Autism Hyperactivity Intellectual disability Short attention span (DECIPHER patient 264639)	N/A