

# Mathematical Analysis II

Felix Chen

## Contents

0.1 Absolute continuous functions . . . . .	4
1 Multi-dimensional Calculus . . . . .	8
1.1 Directional derivatives . . . . .	8

### Step 2

#### Proposition 0.1

The jump function  $J(x)$  is differentiable almost everywhere, and  $J'(x) = 0, a.e..$

*Proof.* The Dini derivatives of  $J(x)$  exist and are non-negative (since  $J$  is increasing).

$$\overline{D}(J)(x) = \max\{D^+(J)(x), D^-(J)(x)\}.$$

Let  $E_\varepsilon = \{\overline{D}(J)(x) > \varepsilon > 0\}$ . We'll prove  $E_\varepsilon$  is null for all  $\varepsilon$ . If  $x \in E_\varepsilon$ ,  $\exists h$  s.t.

$$\frac{J(x+h) - J(x)}{h} > \varepsilon \implies J(x+h) - J(x-h) > \varepsilon h.$$

Let  $N \in \mathbb{N}$  s.t.  $\sum_{n>N} \alpha_n < \frac{\varepsilon \delta}{10}$ . Define  $J_N(x) = \sum_{n>N} j_n(x)$ .

$$E_{\varepsilon,N} = \{\overline{D}(J_N)(x) > \varepsilon\}, \quad E_\varepsilon \subset E_{\varepsilon,N} \cup \{x_1, \dots, x_N\},$$

Since for  $x \neq x_i$ ,

$$\begin{aligned} \overline{D}(J)(x) &= \limsup_{h \rightarrow 0} \frac{J(x+h) - J(x)}{h} \\ &= \limsup_{h \rightarrow 0} \left( \frac{J_N(x+h) - J_N(x)}{h} + \frac{1}{h} \sum_{n=1}^N (j_n(x+h) - j_n(x)) \right) = \overline{D}(J_N)(x). \end{aligned}$$

Next we need to control the measure of  $E_{\varepsilon,N}$ .

For all  $y \in E_{\varepsilon,N}$ , there exists sufficiently small  $h$  s.t.  $J_N(y+h) - J_N(y) > h\varepsilon$ . So the intervals  $(y-h, y+h)$  is a covering of  $E_{\varepsilon,N}$ , and it can be controlled using the value of  $J_N$ . Therefore we hope to find some *disjoint* intervals which cover certain ratio of  $E_{\varepsilon,N}$ .

#### Lemma 0.2

Let  $\mathcal{B}$  be a collection of balls with bounded radius in  $\mathbb{R}^d$ . There exists countably many disjoint balls  $B_i$  s.t.

$$\bigcup_{B \in \mathcal{B}} B \subset \bigcup_{i=1}^{\infty} 5B_i.$$

*Proof.* Let  $r(B)$  denote the radius of  $B$ . Take  $B_1$  s.t.  $r(B_1) > \frac{1}{2} \sup_{b \in B} r(B)$ .  
The rest is the same as before.  $\square$

By lemma, there exists countably many disjoint intervals  $(x_i + h_i, x_i - h_i)$  s.t.

$$\begin{aligned} m^*(E_{\varepsilon, N}) &\leq 5 \sum_{i=1}^{\infty} 2h_i \\ &\leq 10 \sum_{i=1}^{\infty} \varepsilon^{-1} (J_N(x_i + h_i) - J_N(x_i - h_i)) \\ &\leq 10 \varepsilon^{-1} (J_N(b) - J_N(a)) < \delta. \end{aligned}$$

Hence  $m^*(E_{\varepsilon}) \leq m^*(E_{\varepsilon, N}) < \delta \implies m^*(E_{\varepsilon}) = 0$ , which gives  $\overline{D}(J) = 0, a.e..$   $\square$

**Step 3** First we prove  $D^+(F) < \infty, a.e..$

Let  $E_{\gamma} = \{x : D^+(F)(x) > \gamma\}$ .

When  $h \in [\frac{1}{n+1}, \frac{1}{n}]$  :

$$\begin{aligned} \frac{F(x+h) - F(x)}{h} &\leq \frac{n+1}{n} \frac{F(x + \frac{1}{n}) - F(x)}{\frac{1}{n}}, \\ &\geq \frac{n}{n+1} \frac{F(x + \frac{1}{n+1}) - F(x)}{\frac{1}{n+1}}. \end{aligned}$$

Thus

$$D^+(F)(x) = \limsup_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \limsup_{n \rightarrow \infty} \frac{F(x + \frac{1}{n}) - F(x)}{\frac{1}{n}}$$

is measurable.

**Lemma 0.3** (Riesz sunrise lemma)

Let  $G(x)$  be a continuous function on  $\mathbb{R}$ . Define

$$E = \{x : \exists h > 0, \text{ s.t. } G(x+h) > G(x)\}.$$

Then  $E$  is open and  $E = \bigcup_{i=1}^{\infty} (a_i, b_i)$ , where  $(a_i, b_i)$  are disjoint finite intervals s.t.  $G(a_i) = G(b_i)$ .

When  $G$  is defined on finite interval  $[a, b]$ , we also have  $G(a) \leq G(b_1)$ .

*Proof.* Note that  $E$  is open since  $G$  is continuous.

Take a maximum open interval  $(a, b) \subset E$ , i.e.  $a, b \notin E$ , so  $G(a) \geq G(b)$ .

Since  $b \notin E, G(x) \leq G(b), \forall x > b$ . If  $G(a) > G(b)$ , Let  $G(a + \varepsilon) > G(b)$ , as  $a + \varepsilon \in E$ , exists  $h > 0$  s.t.  $G(a + \varepsilon + h) > G(a + \varepsilon)$ .

But  $G$  has a maximum on  $[a + \varepsilon, b]$ , say  $G(c)$ , we must have  $c \neq a + \varepsilon, b$ . This leads to a contradiction.  $\square$

**Remark 0.4** — This lemma provides a better estimation than previous covering lemmas, since it directly claims that  $E$  can be broken into disjoint intervals.

For  $x \in E_\gamma$ ,  $\exists h > 0$  s.t.  $F(x+h) - F(x) > \gamma h$ , by [Lemma 0.3](#) on  $F(x) - \gamma x$ ,

$$m(E_\gamma) \leq \sum_{k=1}^{\infty} (b_k - a_k) \leq \gamma^{-1} \sum_{k=1}^{\infty} (F(b_k) - F(a_k)) \leq \gamma^{-1} (F(b) - F(a)).$$

Therefore when  $\gamma \rightarrow \infty$ ,  $m(E_\gamma) \rightarrow 0$ .

The last part is  $D^+(F) \leq D_-(F)$ , a.e..

Similarly it's sufficient to prove the following set is null for all rational numbers  $r < R$ :

$$E_{r,R} = \{D^+(F)(x) > R, D_-(F)(x) < r\}.$$

Since  $D^+(F)$  is measurable,  $E_{r,R}$  is measurable. If  $m(E_{r,R}) > 0$ , we can restrict it to a smaller interval  $[c, d] \subset [a, b]$  such that  $d - c < \frac{R}{r} m(E_{r,R})$ .

Let  $G(x) = F(-x) + rx$ , by [Lemma 0.3](#) on  $[-d, -c]$ ,

$$\{s : \exists h > 0, G(x+h) > G(x)\} = \bigcup_k (-b_k, -a_k).$$

Note that  $-E_{r,R}$  is contained in the above set, and  $G(-b_k) \leq G(-a_k) \iff F(b_k) - F(a_k) \leq r(b_k - a_k)$ ,

We use [Lemma 0.3](#) again on each  $(a_k, b_k)$  and  $F(x) - Rx$ ,

$$E_{r,R} \cap (a_k, b_k) \subset \{x : \exists h > 0, F(x+h) - F(x) \geq Rh\} = \bigcup_{l=1}^{\infty} (a_{k,l}, b_{k,l}).$$

Hence

$$\begin{aligned} m(E_{r,R}) &\leq \sum_{k,l=1}^{\infty} (b_{k,l} - a_{k,l}) \\ &\leq R^{-1} \sum_{k,l=1}^{\infty} (F(b_{k,l}) - F(a_{k,l})) \leq R^{-1} \sum_{k=1}^{\infty} (F(b_k) - F(a_k)) \\ &\leq R^{-1} r \sum_{k=1}^{\infty} (b_k - a_k) \leq R^{-1} r (d - c), \end{aligned}$$

which gives a contradiction! So  $m(E_{r,R}) = 0$  for all rationals  $r < R$ . Therefore we're done by

$$m(\{D^+(F) > D_-(F)\}) \leq \sum_{r,R} m(E_{r,R}) = 0$$

Now we can complete the proof of ???. Here we state the theorem again:

Let  $F$  be an increasing function on  $[a, b]$ , then  $F$  is differentiable almost everywhere, and

$$\int_a^b F'(x) dx \leq F(b) - F(a).$$

Let  $F_n(x) = n(F(x + \frac{1}{n}) - F(x))$ , where  $F(x) = F(b)$  for  $x > b$ . Since  $F_n \geq 0$ , by Fatou's

Lemma, (we've already proved  $F$  is differentiable almost everywhere and  $F' \geq 0$ )

$$\begin{aligned}
 \int_a^b \liminf_{n \rightarrow \infty} F_n \, dx &\leq \liminf_{n \rightarrow \infty} \int_a^b F_n \, dx \\
 \implies \int_a^b F'(x) \, dx &\leq \liminf_{n \rightarrow \infty} \int_a^b n \left( F\left(x + \frac{1}{n}\right) - F(x) \right) \, dx \\
 &= \liminf_{n \rightarrow \infty} n \left( \int_{a+\frac{1}{n}}^{b+\frac{1}{n}} F(x) \, dx - \int_a^b F(x) \, dx \right) \\
 &= \liminf_{n \rightarrow \infty} \left( F(b) - n \int_a^{a+\frac{1}{n}} F(x) \, dx \right) \\
 &\leq F(b) - F(a)
 \end{aligned}$$

### §0.1 Absolute continuous functions

**Definition 0.5** (Absolute continuity). We say a function  $F(x)$  is **absolutely continuous** on interval  $[a, b]$ , if  $\forall \varepsilon > 0, \exists \delta > 0$ , such that for all disjoint intervals  $(a_k, b_k), k = 1, \dots, N$  with

$$\sum_{k=1}^N (b_k - a_k) < \delta,$$

we must have

$$\sum_{k=1}^N |F(b_k) - F(a_k)| < \varepsilon.$$

The space consisting of all the absolutely continuous functions on  $[a, b]$  is denoted by  $Ac([a, b])$ .

#### Example 0.6

A  $C^1$  function with bounded derivative or a Lipschitz function is absolutely continuous.

Some obvious properties of absolutely continuous function  $F$ :

- $F$  is continuous;
- $F$  has bounded variation, i.e.  $F \in BV$ .
- $F$  is differentiable almost everywhere, since  $F = F_1 - F_2$ , where  $F_1, F_2$  are increasing.

In fact we have

$$T_F([a, b]) = \int_a^b |F'(x)| \, dx.$$

- If  $N$  is a null set, then  $F(N)$  is also null. In particular  $F$  maps measurable sets to measurable sets.

*Proof of the last property.* Take intervals  $(a_k, b_k)$  s.t.  $N \subset \bigcup_{k=1}^{\infty} (a_k, b_k)$ . Since  $F(N) \subset F(\bigcup_{k=1}^{\infty} (a_k, b_k))$ ,

$$|F(N)| \leq \sum_{k=1}^{\infty} |F([a_k, b_k])| \leq \sum_{k=1}^{\infty} |F(\tilde{b}_k) - F(\tilde{a}_k)| < \varepsilon.$$

□

**Proposition 0.7**

The space  $Ac([a, b]) \subset BV([a, b])$ , moreover it's an algebra, and it's a separable Banach space under the norm induced from  $BV$ .

Finally we come to the full generalization of Newton-Lebniz formula:

**Theorem 0.8** (Fundamental theorem of Calculus)

A function  $F \in Ac([a, b]) \implies F$  is differentiable almost everywhere,  $F'$  is integrable, and

$$F(x) - F(a) = \int_a^x F'(\tilde{x}) d\tilde{x}, \quad \forall x \in [a, b].$$

*Proof.* Let  $\tilde{F}(x) = F(a) + \int_a^x F'(y) dy \in Ac([a, b])$  (by the absolute continuity of integrals).

We have  $F - \tilde{F} \in Ac([a, b])$  and  $(F - \tilde{F})' = 0, a.e..$

Thus it suffices to prove the following theorem:

**Theorem 0.9**

Let  $F \in Ac([a, b])$ , and  $F' = 0, a.e.$ , then  $F(a) = F(b)$ , i.e.  $F$  is constant on  $[a, b]$ .

To prove this, we'll need Vitali covering theorem:

**Definition 0.10** (Vitali covering). Let  $\mathcal{B} = \{B_\alpha\}$ , where  $B_\alpha$  are closed balls in  $\mathbb{R}^d$ . We say  $\mathcal{B}$  is a **Vitali covering** of a set  $E$ , if  $\forall x \in E, \forall \eta > 0$ , exists  $B_\alpha \in \mathcal{B}$  s.t.  $m(B_\alpha) < \eta, x \in B_\alpha$ .

**Theorem 0.11** (Vitali)

Let  $E \subset \mathbb{R}^d$  with  $m^*(E) < \infty$ , for any Vitali covering  $\mathcal{B}$  of  $E$  and  $\delta > 0$ , exists disjoint balls  $B_1, \dots, B_n \in \mathcal{B}$ , such that

$$m^*\left(E \setminus \bigcup_{i=1}^n B_i\right) < \delta.$$

*Proof.* For all  $\varepsilon > 0$ , exists an open set  $A$  s.t.  $E \subset A$  and  $m(A) < m^*(E) + \varepsilon < +\infty$ .

Remove all the balls in  $\mathcal{B}$  with radius greater than 1. Each time we take a ball  $B_i$  with radius greater than  $\frac{1}{2} \sup_{B \in \mathcal{B}'} r(B)$ , where  $\mathcal{B}'$  are the remaining balls, and remove all the balls which intersect with  $B_i$ .

If we end up with finitely many balls  $B_1, \dots, B_n$ , we must have  $E \subset \bigcup_{i=1}^n B_i$ , otherwise  $x \notin B_i \implies \exists B \ni x, B \cap B_i = \emptyset$ , contradiction!

If we take out countably many balls  $B_1, B_2, \dots \subset A$ , since  $\sum_{i=1}^{\infty} m(B_i) \leq m(A) < \infty$ , there exists  $N$  s.t.  $\sum_{i=N+1}^{\infty} m(B_i) < 5^{-d}\delta$ .

Now we only need to prove

$$E \setminus \bigcup_{i=1}^N B_i \subset \bigcup_{i>N} 5B_i.$$

□

Let  $E = \{x : F'(x) = 0\}$ ,  $\forall x \in E$ ,  $\exists \delta(x) > 0$ , s.t.

$$|F(y) - F(x)| < \varepsilon|y - x|, \forall |y - x| < \delta(x).$$

Hence  $[x - h, x + h]$ ,  $0 < h < \delta(x)$  is a Vitali covering of  $E$ . By [Theorem 0.11](#), there exists finitely many disjoint intervals  $[x_k - h_k, x_k + h_k] = I_k$  s.t.

$$m^*\left(E \setminus \bigcup_{k=1}^N I_k\right) < \varepsilon.$$

Assume  $a \leq a_1 < b_1 < \dots < a_N < b_N \leq b$ , by absolute continuity and  $|F(b_k) - F(a_k)| < \varepsilon(b_k - a_k)$ ,

$$F(b) - F(a) \leq \sum_{k=1}^N |F(b_k) - F(a_k)| + \sum_{k=0}^N |F(a_{k+1}) - F(b_k)| \leq \varepsilon(b - a) + \delta.$$

Here we complete the proof of the generalized Fundamental theorem of Calculus.  $\square$

There's another version of this theorem which looks like Newton-Lebniz formula more:

**Theorem 0.12**

Let  $F$  be a differentiable function on  $[a, b]$ , if  $F'$  is Lebesgue integrable, then

$$F(b) - F(a) = \int_a^b F'(x) dx.$$

We need to prove a lemma first.

**Theorem 0.13**

Let  $F$  be real function on  $[a, b]$ , if  $F$  is differentiable on  $E$ , and  $|F'| \leq M$  in  $E$ , then

$$m^*(F(E)) \leq Mm^*(E).$$

*Proof.* For all  $\varepsilon > 0$ ,  $x \in E$ ,  $\exists \delta > 0$ ,

$$\left| \frac{F(x+h) - F(x)}{h} - M \right| < \varepsilon, \quad \forall |h| < \delta.$$

So  $[x - h, x + h]$  is a Vitali covering of  $E$ . By Vitali's theorem ([0.11](#)), exists disjoint intervals  $I_i = [x_i - h_i, x_i + h_i]$  s.t.

$$m^*\left(E \setminus \bigcup_{i=1}^{\infty} I_i\right) = 0, \quad \sum_{i=1}^{\infty} 2h_i \leq m^*(E) + \varepsilon.$$

But for  $y \in I_i$ ,  $|F(y) - F(x_i)| \leq (M + \varepsilon)h_i$ , thus  $m^*(F(I_i)) \leq 2(M + \varepsilon)h_i = (M + \varepsilon)|I_i|$ .

$$\begin{aligned} m^*(F(E)) &\leq m^*\left(F\left(E \cap \bigcup_{i=1}^{\infty} I_i\right)\right) + m^*\left(F\left(E \setminus \bigcup_{i=1}^{\infty} I_i\right)\right) \\ &\leq \sum_{i=1}^{\infty} m^*(F(I_i)) + m^*\left(F\left(E \setminus \bigcup_{i=1}^{\infty} I_i\right)\right) \\ &\leq (M + \varepsilon)(m^*(E) + \varepsilon) + m^*\left(F\left(E \setminus \bigcup_{i=1}^{\infty} I_i\right)\right) \end{aligned}$$

So it suffices to prove the case when  $E$  is null. Define

$$E_n = \left\{ x \in E : |F(y) - F(x)| \leq (M + \varepsilon)|y - x|, \forall |y - x| < \frac{1}{n} \right\}.$$

Observe that  $E_n \nearrow E$  and  $F(E_n) \nearrow F(E)$ . There exists disjoint intervals  $J_{n,k}$  s.t.

$$E_n \subset \bigcup_{k=1}^{\infty} J_{n,k}, \quad \sum_{k=1}^{\infty} |J_{n,k}| \leq \min \left\{ \frac{1}{n}, \varepsilon \right\}.$$

Thus

$$m^*(F(E_n)) \leq \sum_{k=1}^{\infty} m^*(F(E_n \cap J_{n,k})) \leq \sum_{k=1}^{\infty} (M + \varepsilon)|J_{n,k}| \leq \varepsilon(M + \varepsilon).$$

Taking  $\varepsilon \rightarrow 0$  we get  $F(E_n)$  is null. So  $F(E) = \lim_{n \rightarrow \infty} F(E_n)$  is null, which completes the proof.  $\square$

Returning to the proof of the theorem, in fact we only need to prove

$$|F(b) - F(a)| \leq \int_a^b |F'(x)| dx,$$

since this implies  $F$  is absolutely continuous. For all  $\varepsilon > 0$ , let

$$E_n = \{x \in [a, b] : n\varepsilon \leq |F'(x)| < (n+1)\varepsilon\}.$$

By our lemma,  $m^*(F(E_n)) \leq (n+1)\varepsilon m(E_n) \leq \varepsilon m(E_n) + \int_{E_n} |F'(x)| dx$ .

Hence

$$\begin{aligned} |F(b) - F(a)| &\leq m(F([a, b])) \leq \sum_{n=0}^{\infty} m^*(F(E_n)) \\ &\leq \varepsilon(b-a) + \int_a^b |F'(x)| dx. \end{aligned}$$

#### Theorem 0.14

A rectifiable curve  $\gamma(t) = (x(t), y(t))$  with  $x, y$  absolutely continuous has length

$$L(\gamma) = \int_a^b |\gamma'(t)| dt.$$

*Proof.* Since  $|\gamma(t_i) - \gamma(t_{i-1})| = \left| \int_{t_{i-1}}^{t_i} \gamma'(t) dt \right| \leq \int_{t_{i-1}}^{t_i} |\gamma'(t)| dt$ , thus  $L(\gamma) \leq \int_a^b |\gamma'(t)| dt$ .

$\forall \varepsilon > 0$ , we can take a step function (with vector values)  $g$  s.t.  $\gamma' = g + h$ , and  $\int_a^b |h| dx < \varepsilon$ .

Define

$$G(x) = G(a) + \int_a^x g(t) dt, \quad H(x) = H(a) + \int_a^x h(t) dt.$$

We have  $\gamma(t) = G(t) + h(t)$ , and  $T_\gamma([a, b]) \geq T_G([a, b]) - T_H([a, b])$ .

$$\begin{aligned} L(\gamma) &= T_\gamma([a, b]) \geq \int_a^b |g| dt - \int_a^b |h| dt \\ &\geq \int_a^b |\gamma'(t)| dt - 2 \int_a^b |h| dt \\ &\geq \int_a^b |\gamma'(t)| dt - 2\varepsilon. \end{aligned}$$

which gives the opposite inequality.  $\square$

**Proposition 0.15** (substitution formula)

Let  $\phi : [a, b] \rightarrow [c, d]$  be strictly increasing AC function. For a function  $f$  on  $[c, d]$ , we have

$$\int_c^d f(y) dy = \int_a^b f(\phi(x))\phi'(x) dx.$$

*Proof.* It's equivalent to  $m(\phi(E)) = \int_E \phi' dx$ .  $\square$

## §1 Multi-dimensional Calculus

In this section we'll generalize the differentiation and integration theory to higher dimensions, and finally reach the generalized Fundamental Theorem of Calculus (Stokes' formula). Recall that Lebesgue integral is already defined on higher dimensions, so here we mainly study the differentiation of multi-dimensional functions.

### §1.1 Directional derivatives

Let  $\Omega$  be a simply connected open set in  $\mathbb{R}^d$ .  $f$  is a multi-variable function on  $\Omega$ . Let  $(x_1, \dots, x_n)$  be a coordinate system on  $\Omega$ , we can write  $f = f(x_1, \dots, x_n)$ .

**Definition 1.1** (Directional derivatives). Let  $v \in \mathbb{R}^d$  be a nonzero vector. If

$$\lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

exists, then we say the **directional derivative** of  $f$  in direction  $v$  exists at  $x_0$ , denoted by

$$\frac{\partial f}{\partial v}(x_0) = (\nabla_v f)(x_0) = \lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}.$$

**Definition 1.2** (Partial derivatives). Let  $(x_1, \dots, x_n)$  be a coordinate system, let  $e_i = (0, \dots, 1, \dots, 0)$  be the  $i$ -th vector of the standard basis. The directional derivative in  $e_i$

$$(\nabla_{e_i} f)(x_0) = \frac{\partial f}{\partial x_i}(x_0) = \lim_{h \rightarrow 0} \frac{f(x_1, \dots, x_i + h, \dots, x_n) - f(x_1, \dots, x_n)}{h}.$$

is called the  $i$ -th **partial derivative** of  $f$ . Here  $\frac{\partial}{\partial x_i}$  is also called a “vector field”.

**Remark 1.3** — The partial derivatives rely on the coordinate, but the directional derivatives is independent of the coordinate (i.e. geometry quantities).

**Example 1.4**

Let  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ , and  $f(x, y) = g(x)$  for some  $g$ .

$$\frac{\partial f}{\partial y} = \lim_{h \rightarrow 0} \frac{f(x, y + h) - f(x, y)}{h} = 0.$$



**Example 1.5**

Consider  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ ,

$$f(x, y) = \begin{cases} \frac{xy}{x^2 + y^2}, & x^2 + y^2 \neq 0; \\ 0, & x = y = 0. \end{cases}$$

The partial derivative

$$\frac{\partial f}{\partial x}(0, 0) = \lim_{h \rightarrow 0} \frac{f(h, 0) - f(0, 0)}{h} = 0 = \frac{\partial f}{\partial y}(0, 0).$$

But the directional derivative in  $v = (v_1, v_2)$  is

$$(\nabla_v f)(0, 0) = \lim_{h \rightarrow 0} \frac{f(hv_1, hv_2) - f(0, 0)}{h} = \lim_{h \rightarrow 0} \frac{v_1 v_2}{h(v_1^2 + v_2^2)},$$

which doesn't exist for  $v_1 v_2 \neq 0$ .

The main idea of differentiation in 1 dimensional is to estimate a function locally using a straight line. Likely, in higher dimensions, the differentiation is also estimating a function locally using a *linear map*.

**Definition 1.6** (Differentiation). Let  $f : \Omega \rightarrow \mathbb{R}$ ,  $x_0 \in \Omega$ . If there exists a linear map  $A : \mathbb{R}^d \rightarrow \mathbb{R}$  s.t.

$$f(x_0 + v) = f(x_0) + A(v) + o(|v|) \iff \lim_{|v| \rightarrow 0} \frac{|f(x_0 + v) - f(x_0) - A(v)|}{|v|} = 0,$$

then we say  $f$  is **differentiable** at  $x_0$ , and the linear map  $A$  is called the **differentiation** of  $f$  at  $x_0$ , denoted by

$$df|_{x_0} = df(x_0) = A : \mathbb{R}^d \rightarrow \mathbb{R}.$$

If  $f$  is differentiable everywhere, we say  $f$  is a differentiable function.

**Remark 1.7** — In fact this definition can be generalized to any Banach space. Keep in mind that  $df(x_0)$  is a *linear map* instead of a number, the reason why the one dimensional differentiation is a number is that a linear map in one dimension is identical to a scalar.

**Theorem 1.8**

Let  $f$  be a function differentiable at  $x_0$ , then its directional derivatives exist at  $x_0$ ,  $\forall v \in \mathbb{R}^d$ ,

$$(\nabla_v f)(x_0) = (df(x_0))(v) = \sum_{i=1}^d \frac{\partial f}{\partial x_i} \cdot v_i = \nabla f \cdot v.$$

where  $\nabla f = (\frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_n})$  is the **gradient vector** of  $f$ .

*Proof.* Note that

$$\frac{f(x_0 + hv) - f(x_0)}{h} = \frac{df(x_0)(hv) + o(h|v|)}{h} \rightarrow df(x_0)(v).$$

$$df(x_0)(v) = df(x_0) \left( \sum_{i=1}^d v_i e_i \right) = \sum_{i=1}^d v_i df(x_0)(e_i) = \sum_{i=1}^d v_i \frac{\partial f}{\partial x_i}.$$

□

**Example 1.9**

Let  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ ,

$$f(x, y) = \begin{cases} \frac{y^2}{x}, & x \neq 0; \\ 0, & x = 0. \end{cases}$$

Note that the directional derivatives of  $f$  exists at  $(0, 0)$ , but  $f$  is not continuous at  $x_0$ , so not differentiable.

**Theorem 1.10**

Let  $\Omega \subset \mathbb{R}^d$ . If the partial derivatives of  $f$  exists and are continuous at  $x_0$ , then  $f$  is differentiable at  $x_0$ .

*Proof.* Let  $u_j = (v_1, \dots, v_j, 0, \dots, 0)$ .

$$\begin{aligned} f(x_0 + v) - f(x_0) - (\nabla f)(x_0) \cdot v &= \sum_{j=1}^d f(x_0 + u_j) - f(x_0 + u_{j-1}) - \frac{\partial f}{\partial x_j}(x_0) v_j \\ &= \sum_{j=1}^d \frac{\partial f}{\partial x_j}(x_0 + u_{j-1} + \xi_j e_j) v_j - \frac{\partial f}{\partial x_j}(x_0) v_j \end{aligned}$$

where the last step used Lagrange's theorem. Since  $v_j < |v|$  and the partial derivatives are continuous at  $x_0$ , so when  $|v| \rightarrow 0$ , the above also approach to 0. □

**Corollary 1.11**

If  $f$  is differentiable on  $\Omega$ , and  $df = 0$ , then  $f$  is constant on  $\Omega$ .

**Proposition 1.12**

Let  $f : \Omega \rightarrow \mathbb{R}$  be a function differentiable at  $x_0$ , and  $f$  achieves its local extremum at  $x_0$ , then  $df(x_0) = 0$ .

*Proof.* Trivial. □

**Definition 1.13.** Let  $\Omega \subset \mathbb{R}^d$ ,  $\Omega' \subset \mathbb{R}^{d'}$ ,  $f : \Omega \rightarrow \Omega'$ . If there exists a linear map

$$df|_{x_0} : \mathbb{R}^d \rightarrow \mathbb{R}^{d'},$$

s.t.

$$f(x_0 + v) = f(x_0) + df(x_0)(v) + o(|v|),$$

then we say  $f$  is differentiable at  $x_0$ , the linear map  $df(x_0)$  is called the differentiation of  $f$  at  $x_0$ .

**Proposition 1.14**

Let  $f = (f_1, \dots, f_{d'})$ .  $f$  is differentiable at  $x_0$  is equivalent to  $f_i$  is differentiable at  $x_0$ , and  $df(x_0) : \mathbb{R}^d \rightarrow \mathbb{R}^{d'}$  can be represent as the matrix

$$\left( \frac{\partial f_i}{\partial x_j}(x_0) \right)_{i,j}$$

this is called the **Jacobi matrix** of  $f$  at  $x_0$ , denoted by  $J(f)(x_0)$ .