

Mathematical Analysis II

Felix Chen

Contents

0.1	Stokes' formula	1
0.2	Differential forms	4

§0.1 Stokes' formula

Intuitively, Stokes' formula states that: Let D be a region, $d\omega$ be a differential form, then

$$\int_D d\omega = \int_{\partial D} \omega.$$

Here ∂D means the “boundary” of D .

Of course we need some “regularity” requirements of D and ω , and it's the generalization of Newton-Lebniz formula into higher dimensions.

Definition 0.1.1 (Bounded regions with boundary). Let $\Omega \subset \mathbb{R}^n$ be a compact set, we say it's a **bounded region with boundary** if $\forall x \in \partial\Omega$, there exists open sets $U, V \subset \mathbb{R}^n$, $x \in U$ and a continuous homeomorphism $\Phi : U \rightarrow V$, such that

$$\Phi(U \cap \Omega) = V \cap \{x_n \geq 0\}, \quad \Phi(U \cap \partial\Omega) = V \cap \{x_n = 0\}.$$

If Φ is also C^1 , we say $x \in \partial\Omega$ is a **regular point**, otherwise a **singular point**.

Lemma 0.1.2

Let Ω be a bounded region with boundary, for all regular $p \in \partial\Omega$, there exists a unique unit vector $\nu(p) \in \mathbb{R}^n$, and $\varepsilon > 0$, s.t.

$$\nu(p) \perp T_p \partial\Omega, |\nu(p)| = 1, p - t\nu(p) \in \Omega, \forall 0 < t < \varepsilon.$$

We call $\nu(p)$ the **outward unit normal vector** of p .

Proof. By the definition of regular points, we may assume that:

$$\Omega \cap V = \{x \in V \mid f(x) \geq 0\}, \quad \partial\Omega \cap V = \{x \in V \mid f(x) = 0\}.$$

Where f is a C^1 function.

Since ∇f is nonzero, the tangent space $T_p \partial\Omega = \{v \mid v \cdot \nabla f = 0\}$.

Let $\nu(p) = -\frac{\nabla f}{|\nabla f|}$, then it's obvious $\nu(p)$ points outside of Ω . □

Now for a cuboid I and a C^1 function ϕ ,

$$\begin{aligned} \int_I \frac{d\phi}{dx_n} dx &= \int_{I_{n-1}} \phi(x_1, \dots, x_{n-1}, b_n) dx_1 \cdots dx_{n-1} - \int_{I_{n-1}} \phi(x_1, \dots, x_{n-1}, a_n) dx_1 \cdots dx_{n-1} \\ &= \int_{\partial I} \phi \cdot \nu_n d\sigma. \end{aligned}$$

Where σ is the measure on the boundary, ν is the outward unit normal vector.

Lemma 0.1.3

Let K be a compact set in \mathbb{R}^n , $U \supset K$ is open, there exists a smooth function f such that $\text{supp } f \subset U$, and $f|_K > 0$.

Proof. Let $\rho(x)$ be a smooth function s.t. $\rho(x) = 1$ for $|x| \leq 1$ and $\rho(x) = 0$ for $|x| \geq 2$. Let

$$g(x) = \int_{|y| \leq 2} f(x - \delta y) \rho(y) dy.$$

Then g is a smooth non-negative function. □

Theorem 0.1.4 (Unit decomposition on compact sets)

Let K be a compact set, $\{U_1, \dots, U_k\}$ is an open covering of K . There exists smooth functions f_1, \dots, f_k s.t.

$$1 = f_1(x) + f_2(x) + \cdots + f_k(x), \quad \text{supp } f_i(x) \subset U_i.$$

Proof. For $1 \leq i \leq k$, $\delta > 0$, define

$$K_i^\delta = \{x \in U_i \mid d(x, U_i^c) > \delta\}.$$

Note that $\{\bigcup_{i=1}^k K_i^{\frac{1}{m}}\}_{m=1}^\infty$ is also an open covering of K , thus there exists N s.t.

$$K \subset \bigcup_{i=1}^k K_i^{\frac{1}{N}}.$$

Hence by lemma we have g_i s.t. $\text{supp } g_i \subset U_i$ and $g_i > 0$ on the closure of $K_i^{\frac{1}{N}}$. Similarly we have a smooth function g s.t. $g(x) = 0$ on K , and $g > 0$ outside of the closure of $\bigcup_{i=1}^k K_i^{\frac{1}{N}}$.

Let $G(x) = g_1(x) + \cdots + g_k(x) + g(x) > 0$ on $\bigcup_{i=1}^k U_i$, then we can define $f_i(x) = \frac{g_i(x)}{G(x)}$ which satisfy the condition. □

Theorem 0.1.5

Let Φ be a C^1 homeomorphism from a cuboid I to Ω , then Ω satisfies Stokes' formula:

$\forall \phi \in C^1(\overset{\circ}{D}) \cap C(\overline{D})$, we have

$$\int_D \nabla \phi dx = \int_{\partial D} \phi \nu d\sigma.$$

Proof. Since $\Omega = \Phi(I)$, let y be the coordinates on I , $x = \Phi(y)$,

$$\int_{\Phi(I)} \nabla \varphi \, dx = \int_I \nabla \varphi(\Phi(y)) (\mathrm{d}\Phi)^{-1} J_\Phi \, dy.$$

Let $A = \mathrm{d}\Phi$, WLOG $J_\Phi > 0$. Using the index notation and Einstein summation,

$$A_{kj} A^{ji} = A^{kj} A_{ji} = \delta_{ki}.$$

Thus

$$\partial_{y_j} \varphi A^{ji} |A| = \partial_{y_j} (\varphi A^{ji} |A|) - \varphi \partial_{y_j} (A^{ji} |A|)$$

Since $|A| = A_{kl} A^{kl} |A|$, $A_{kl} = \frac{\partial \Phi_k}{\partial y_l}$.

$$\begin{aligned} \partial_{y_j} (A^{ji} |A|) &= |A| \partial_{y_j} A^{ji} + A^{ji} \partial_{y_j} |A| \\ &= |A| \partial_{y_j} A^{ji} + A^{ji} |A| \partial_{y_j} A_{kl} A^{kl} \\ &= |A| (\partial_{y_j} A^{ji} + \partial_{y_l} A_{kj} A^{kl}) \\ &= |A| (\partial_{y_j} A^{ji} - \partial_{y_j} A^{ji}) = 0. \end{aligned}$$

Hence by our previous work,

$$\int_I \partial_{y_j} (\varphi A^{ji} |A|) \, dy = \int_{\partial I} \varphi A^{ji} |A| \nu_j \, d\sigma.$$

Putting this together for all i 's, note that $\tilde{\nu} = \frac{\nabla \Phi_n^{-1}}{|\nabla \Phi_n^{-1}|}$,

TODO

□

Let (ϕ_1, \dots, ϕ_n) be an element in the tangent bundle TM , it can represent a vector field

$$X = (\phi_1, \dots, \phi_n) = \sum_{i=1}^n \phi_i \frac{\partial}{\partial x_i}.$$

Here $X \in TM$, $X(p) \in T_p M$.

We define the **divergence** of X to be

$$\mathrm{div}(X) = \sum_{i=1}^n \frac{\partial \phi_i}{\partial x_i}.$$

The Stokke's formula can be presented as divergence theorem:

Theorem 0.1.6 (Divergence theorem)

Let X be a vector field,

$$\int_D \mathrm{div}(X) \, dx = \int_{\partial D} X \cdot \nu \, d\sigma.$$

Another commonly-used operator is the **Laplace operator**:

$$\Delta = \mathrm{div} \cdot \nabla, \quad \Delta \phi = \mathrm{div}(\nabla \phi) = \mathrm{tr}(H_\phi) = \sum_{i=1}^n \frac{\partial^2 \phi}{\partial x_i^2}.$$

When $n = 2$, we have $X = P \frac{\partial}{\partial x} + Q \frac{\partial}{\partial y}$, $\operatorname{div}(X) = \frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y}$,

$$\int_D \left(\frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} \right) dx dy = \int_{\partial D} X \cdot \nu d\sigma.$$

Since ∂D is a curve $\gamma(t)$, so $d\sigma = |\gamma'(t)| dt$. Let $\gamma(t) = (x(t), y(t))$, then $\nu(t) = \frac{(y'(t), -x'(t))}{|\gamma'(t)|}$. Here we must take $\gamma(t)$ to be *counterclockwise* to ensure ν points outside of D .

Thus we get

$$\int_{\partial D} X \cdot \nu d\sigma = \int_{\gamma} \frac{Py'(t) - Qx'(t)}{|\gamma'(t)|} |\gamma'(t)| dt = \int_D (P dy - Q dx).$$

This result is known as *Green's formula*.

This leads to the curve integrals of the second type: let $\gamma(t) \in \mathbb{R}^d$, X a vector field, we call the integral

$$\int_{\gamma} \sum_{i=1}^d X^i dx_i = \int_{\gamma} X \cdot d\gamma(t).$$

the **curve integral of the second type**.

When $n = 3$, the result is called *Gauss's formula*, we have $X = P \frac{\partial}{\partial x} + Q \frac{\partial}{\partial y} + R \frac{\partial}{\partial z}$,

$$\int_D \operatorname{div}(X) dx dy dz = \int_{\partial D} X \cdot \nu d\sigma.$$

Let $\gamma(u, v) = (x, y, z)$ be a parametrization of ∂D . We have two tangent vector γ_u, γ_v , so the normal vector is defined as $\nu = \frac{\gamma_u \times \gamma_v}{|\gamma_u \times \gamma_v|}$. Also $d\sigma = |\gamma_u \times \gamma_v| du dv$. After some computation we can get

$$\begin{aligned} \nu d\sigma &= (dy dz, dz dx, dx dy). \\ \int_D \operatorname{div}(X) dx dy dz &= \int_{\partial D} (P dy dz + Q dz dx + R dx dy). \end{aligned}$$

§0.2 Differential forms

Let T_p^*M denote the *dual space* of T_pM , and dx_i is the dual basis of $\frac{\partial}{\partial x_i}$. The linear combination of dx_i are called **differential forms**, and a differential form on a manifold can be written as $\sum_{i=1}^n a_i dx_i$, where a_i are functions on M .

We can construct differential forms of higher order, the order is $1 \leq k \leq n$, called **k-forms**, which is a linear combination of

$$dx_{i_1} dx_{i_2} \cdots dx_{i_k}, \quad i_1 < i_2 < \cdots < i_k.$$

Here the product is *wedge product*, i.e. $dx_i dx_j = -dx_j dx_i$. We denote the space of all k -forms by $\Lambda^k(\Omega)$.

We can define the multiplication of forms: let $\omega_1 \in \Lambda^{k_1}, \omega_2 \in \Lambda^{k_2}$, then $\omega_1 \wedge \omega_2 \in \Lambda^{k_1+k_2}$ by multiplying the coefficients and dx_i 's respectively.

There's also an operator called **exterior differentiation** $d : \Lambda^k \rightarrow \Lambda^{k+1}$, where

$$d(a dx_{i_1} \cdots dx_{i_k}) = \sum_{j=1}^n \frac{\partial a}{\partial x_j} dx_j dx_{i_1} \cdots dx_{i_k}.$$

This operator behaves like the derivatives very much:

$$d(\omega_1 + \omega_2) = d\omega_1 + d\omega_2, \quad d(\omega_1 \omega_2) = d\omega_1 \wedge \omega_2 + (-1)^{k_1 k_2} \omega_1 \wedge d\omega_2.$$

Note that $\omega_1 \wedge \omega_2 = (-1)^{k_1 k_2} \omega_2 \wedge \omega_1$, so when $k_1 k_2$ is even, the wedge product may not be anti-symmetrical.

If we have a coordinate transformation $\Phi : (x_1, \dots, x_n) \rightarrow (y_1, \dots, y_n)$, we have

$$dy_i = \sum_{j=1}^n \frac{\partial y_i}{\partial x_j} dx_j,$$

thus $dy_1 \cdots dy_n = J_\Phi dx_1 \cdots dx_n$. Here J_Φ can be negative, so the differential forms already contains the information of orientation.

Theorem 0.2.1

Let ω be a differential form, $d(d\omega) = 0$.

Proof. Partial derivatives commute. □

Definition 0.2.2. Let ω be a differential form, if $d\omega = 0$, we say ω is a **closed form**, if there exists ω_1 s.t. $d\omega_1 = \omega$, then ω is a **exact form**.

The theorem above tells us that exact forms must be closed, but in general closed forms may not be exact, it depends on the topology structure of Ω .

Theorem 0.2.3 (Poincare)

The closed forms on \mathbb{R}^n must be exact.

Proof. Use induction, when ω is an n -form this can be proved by computation.

For a generic form $\omega = \omega_1 + dx_1 \wedge \omega_2$, where ω_1, ω_2 do not contain dx_1 . We want to find ω_3 s.t. $d\omega_3 = dx_1 \wedge \omega_2 + \omega_4$, where ω_3, ω_4 don't contain dx_1 as well. (The construction is direct)

Since $\omega - d\omega_3 = \omega_1 - \omega_4$, and

$$d(\omega - d\omega_3) = d\omega = 0 \implies d(\omega_1 - \omega_4) = 0.$$

Since $d(\omega_1 - \omega_4) = 0$ and it doesn't contain dx_1 , hence all its coefficients can't contain dx_1 . Thus we can view it as a differential form in \mathbb{R}^{n-1} . □

Remark 0.2.4 — When Ω is simply connected, then all the closed 1-forms are exact. Also this is equivalent to the integral on any closed curves are 0.

We can rewrite Stokes' formula using differential forms:

Theorem 0.2.5 (Stokes' formula)

Let D be a $k+1$ dimensional orientable manifold, $\omega \in \Lambda^k(\mathbb{R}^n)$, we have

$$\int_D d\omega = \int_{\partial D} \omega.$$