

MINIMAL RESIDUAL DISEASE (MRD) BY 8-COLOR FLOW
CYTOMETRY

Date: September 09, 2026

Patient Name:	Neha Gupta	MRN / Patient ID:	MED-HEM-2026-5510
Age / Gender:	35 Yrs / Female	Blood Group & Rh:	A- (A Rh Negative)
Clinical Diagnosis:	B-Cell Acute Lymphoblastic Leukemia (B-ALL Post-Induction)	Referring Consultant:	Dr. Ashok Vaid, MD, DM

Analyzed Cellular Fraction	Events Acquired	Residual Blasts (%)	Clinical Assessment
Bone Marrow Mononuclear Cells	1,250,000 Events	< 0.01% (Negative)	MRD Negative (< 10 ⁻⁴)
CD19+ / CD10+ Immature Precursors	840,000 Events	Undetected Aberrant	Normal Hematogones Only
Leukemia-Associated Immunophenotype (LAIP)	CD34+ CD19+ CD20- CD58+	Zero Clonal Events	No Relapse Pattern

CLINICAL INTERPRETATION & RECOMMENDATIONS:

- High-sensitivity 8-color flow cytometry acquired > 1 million cellular events (analytical sensitivity threshold of 0.01%).
- Patient has achieved deep immunophenotypic Minimal Residual Disease negative complete remission (MRD < 0.01%).
- Negative MRD status is a highly favorable prognostic indicator prior to consolidation and stem cell matching.

Prepared By:
Technical Specialist
Transplant Immunology Lab

Verified By:
Dr. S. Damodar, MD, DM
Head of Hemato-Oncology & BMT

Laboratory Director:
Dr. Meenakshi Sundaram, FRCPATH
Director of Cellular Therapy