Spotted Python

Week 8 Tutorial

UNSW SYDNEY

- BPT Diagram
- Fitting a Gaussian
- Spectroscopic redshift

BPT Diagram

- Simple Diagnostic Diagram (Kewley+2006)
  - *HII & Composites:* $\log\left(\frac{[OIII]}{[OII]}\right) < -1.701\log\left(\frac{[OI]}{H_\alpha}\right) - 2.163$
  - *LINERS:* $\log\left(\frac{[OIII]}{[OII]}\right) < 1.0\log\left(\frac{[OI]}{H_\alpha}\right) + 0.7$
  - *Seyferts:* $\log\left(\frac{[OIII]}{[OII]}\right) > 1.0\log\left(\frac{[OI]}{H_\alpha}\right) + 0.7$
- What are HII, LINERS, Seyferts?




Figure 5. The [O III]/[O II] versus [O I]/Hα diagnostic diagram for SDSS galaxies with S/N > 3. Galaxies have been classified using the standard BPT diagnostic diagrams (Fig. 4). Ambiguous galaxies are not included. Our new preferred classification scheme is shown in blue. The H80 LINER line (orange dashed) is also shown.

UNSW SYDNEY

SDSS Navigate Visual Tool

- https://skyserver.sdss.org/dr18/VisualTools/navi
- Select an emission line galaxy with a measured spectrum
- Using the flux (EW), place the objects on the $\log\left(\frac{[OIII]}{[OII]}\right)$ vs $\log\left(\frac{[OI]}{H\alpha}\right)$ plot
  - Wavelengths of the emission lines: [O I] λ6300, [O II] λλ3726, [O III] λ5007, [Hα] λ6564
  - λλ indicates that you must sum across the doublet near those wavelengths
- Determine the type of emitter the galaxy is

UNSW SYDNEY

Spectroscopic Redshift

- Recall that redshift can be determined using the following:

$$z = \frac{\lambda_{observed}}{\lambda_{rest}} - 1$$

- List of emission lines are available here: http://astronomy.nmsu.edu/drewski/tableofemissionlines.html#:~:text=The%20ultraviolet%20and%20optical%20spectra,nature%20of%20the%20ionizing%20source