# Machine learning for ICU patient monitoring

"How are machine learning models used to analyze patient vital signs, clinical records, and ventilator parameters to detect changes in the condition of mechanically ventilated ICU patients? What datasets, particularly MIMIC-IV, and prediction techniques are used, and what challenges affect real-time clinical applications?"

*Generated by LeapSpace, Sat Oct 10 2026*

## Machine Learning for Monitoring Condition Change in Mechanically Ventilated ICU Patients at a Glance

Key Findings Table

| Topic | Key finding | Population / task | Evidence |
|---|---|---|---|
| Main clinical use | ML is used to detect extubation failure, weaning failure, ventilator-associated complications, mortality-related deterioration, and broader physiologic decompensation from ICU time series and structured records. | Mechanically ventilated ICU adults; some broader ICU cohorts | [1] [2] [3] [4] |
| Data inputs | Most models rely on structured EHR inputs: vital signs, labs, blood gases, demographics, comorbidities, and ventilator settings such as tidal volume, PEEP, FiO2, mean airway pressure, PSV, compliance, and mechanical power. | Ventilated ICU prediction | [1] [2] [4] [5] |
| MIMIC-IV role | MIMIC-IV is a major recent development dataset for extubation and weaning models, including cohorts for extubation failure, weaning failure, and weaning success. | Adult ICU patients on invasive mechanical ventilation | [1] [2] [6] [7] [8] |
| Common model classes | Ventilated-patient studies frequently use XGBoost, random forest, logistic regression, CatBoost, and multitask models; broader ICU deterioration work often shows gains from LSTM, GRU, attention, TCN, and transformer models. | Ventilated cohorts and general ICU deterioration tasks | [4] [9] [10] [11] [12] [13] [14] [15] |
| Temporal processing | Near-real-time prediction typically uses time-binning, multi-window summaries, waveform summarization, or automatic time-series feature extraction such as tsfresh. | ICU time-series modeling | [16] [17] [18] [19] |
| Performance pattern | Deep temporal models often outperform early-warning scores and simpler baselines in broader ICU tasks; ventilated-patient task performance is often good, but many published wins are still from structured-feature tree models rather than end-to-end multimodal deep learning. | ICU deterioration and ventilated patient prediction | [4] [8] [11] [12] [13] |

| | | | |
|---|---|---|---|
| Labeling challenge | Outcome definitions vary widely across studies, including extubation failure within 48 h, SBT/weaning failure, successful weaning within 14 days, and ventilator-associated complications after a defined ventilation duration. | Cross-study comparison | [1] [2] [3] [6] [7] [20] |
| Real-time barriers | The biggest obstacles to bedside deployment are missingness, irregular sampling, delayed charting, class imbalance, calibration, domain shift, and limited workflow/usability evidence. | Real-time clinical use | [21] [22] [23] [24] [25] [26] [27] |
| External validation | External validation remains uncommon, and performance can fall across hospitals or datasets; some models require fine-tuning after transfer. | Cross-site validation | [25] [26] [28] [29] |
| Interpretability | SHAP, LIME, PDPs, feature importance, and nomogram-style tools are increasingly used, but formal evidence of clinician trust or workflow benefit is largely absent. | Explainable ICU ML | [27] [30] [31] [32] |

## Direct Answer

Machine learning models detect condition change in mechanically ventilated ICU patients mainly by learning patterns in structured ICU data: bedside vital signs, laboratory values, blood gases, demographics, comorbidities, and ventilator parameters such as tidal volume, PEEP, FiO2, mean airway pressure, pressure support, compliance, and mechanical power [1] [2] [4]. These models are used to predict clinically important transitions including extubation failure, weaning failure, readiness for successful weaning, ventilator-associated complications, mortality-related deterioration, and broader decompensation states [1] [2] [3] [7].

MIMIC-IV is a central dataset for this work. Identified MIMIC-IV studies include a large extubation-failure cohort of 16,189 extubated ICU adults, a 3,695-patient T-tube weaning-failure cohort, a 23,242-patient weaning-success cohort, and a 7,758-patient multitask weaning study [4] [6] [7] [8]. Other recurrent ICU datasets include MIMIC-III, eICU, AmsterdamUMCdb, and PhysioNet-derived resources [28] [33] [34] [35].

Prediction techniques span classical ML and deep learning. In ventilated cohorts, tree-based and tabular models such as XGBoost, random forest, CatBoost, and logistic regression are common and often competitive [4] [9] [10]. In broader ICU deterioration tasks, sequential deep models such as LSTM, GRU, attention models, temporal CNNs, and transformers often outperform early-warning scores and simpler baselines, with reported AUROC gains and lead-time advantages of several hours [11] [12] [13] [14] [15].

For real-time clinical application, the main challenge is not simply discrimination. It is operational robustness: irregular sampling, missing data, delayed charting, heterogeneous outcome labels, class imbalance, limited external validation, calibration drift, and weak evidence that explanation tools actually improve clinical workflow or trust [20] [21] [22] [23] [25] [26] [27]. Taken together, the evidence suggests the field is technically promising but still only partially ready for reliable bedside deployment.
**Confidence level: High** - multiple convergent sources support the overall pattern across datasets, tasks, and deployment studies [1] [8] [11] [25].

## Study Scope

- **Time period covered by retrieved studies:** foundational ICU scoring and public-dataset papers through recent 2021-2025 MIMIC-IV ventilator-focused prediction work [7] [8] [33] [36].
- **Disciplines represented:** critical care, respiratory medicine, clinical informatics, time-series machine learning, and bedside decision support [4] [37] [38].
- **Methods represented:** logistic regression, survival-style models, random forest, XGBoost, CatBoost, ANN, LSTM, GRU, attention models, temporal CNNs, transformers, multitask learning, waveform feature extraction, and explainability methods [9] [11] [13] [18] [27] [39].

**Confidence level: High** - the scope is directly described by the retrieved studies and meta-synthesis [11] [33] [38].

### Assumptions & Limitations

- The evidence base is strongest for **structured EHR plus ventilator-setting models**, not for full end-to-end multimodal fusion including free text; retrieved excerpts did not identify a clear published MIMIC-IV example combining ventilator data, vital signs, labs, and notes in one integrated architecture [40] [41].
- Outcome labels are heterogeneous, making direct comparison across studies difficult [1] [2] [3] [6].
- Many preprocessing details needed for replication or real-time deployment are underreported [19] [40] [41].
- External validation, temporal drift checks, and clinician usability studies remain sparse [25] [27] [42]. **Confidence level: High** - these are repeated limitations across the retrieved synthesis and review evidence [25] [26].

### Suggested Further Research

- Build and validate **true multimodal ventilated-patient models** that jointly integrate ventilator waveforms/settings, monitor vitals, labs, and clinical text [40] [41].
- Standardize **label definitions** for extubation failure, weaning failure, and respiratory deterioration [1] [2] [20].
- Increase **multicenter external validation** across MIMIC-IV, eICU, AmsterdamUMCdb, and non-public hospital systems [25] [28] [29].
- Study **calibration maintenance, drift detection, and recalibration schedules** in deployed ICU systems [4] [42] [43].
- Evaluate **workflow fit, trust, alert fatigue, and clinical impact** rather than explanation quality alone [27] [32].

## Full Report

## Foundational Context

Mechanical ventilation creates a clinical setting where patient condition can change rapidly, but those changes are often reflected across multiple data streams rather than a single measurement [4] [37]. Relevant changes include worsening oxygenation, failed spontaneous breathing trials, post-extubation respiratory failure, ventilator-associated complications, and broader hemodynamic or mortality-related deterioration [1] [3] [4].

Historically, ICU monitoring relied on repeated severity or warning scores such as SOFA, APACHE II/III/IV, SAPS II/3, MEWS, NEWS, ViEWS, Rothman Index, and eCART-style systems [36] [44] [45] [46] [47] [48] [49] [50]. These frameworks established two important ideas that modern ML inherits: first, deterioration is best tracked **repeatedly over time**, not from a one-time snapshot; second, routinely collected variables such as vital signs and labs are informative enough to support automated warning systems [44] [46] [51].

Machine learning extends this by learning nonlinear and temporal relationships across many variables simultaneously [11] [52]. In mechanically ventilated patients, this usually means combining structured EHR measurements with respiratory support variables to estimate the probability of a near-term clinical transition, such as failed extubation within 48 hours or unsuccessful weaning after a breathing trial [1] [2]. **Confidence level: High** - supported by both foundational and recent ventilator-specific studies [1] [44].

## Introduction and Clinical Problem

### Clinical motivation and scope

Detecting condition change early matters because mechanically ventilated ICU patients can move from stable support to respiratory decompensation, failed liberation from the ventilator, or broader deterioration within hours [1] [4]. Near-real-time forecasting is therefore valuable for adjusting ventilator settings, planning spontaneous breathing trials, identifying patients at risk of reintubation, and prioritizing closer observation or intervention [26] [39].

The literature spans two overlapping problem types:

1. **Ventilator-specific transition prediction**, such as extubation failure, weaning failure, successful liberation, or ventilator-associated complications [1] [2] [3] [7].
2. **Broader ICU deterioration prediction**, where ventilated patients are included within larger critical care cohorts and models forecast mortality, physiologic decline, or severe complications [11] [13] [52].

This distinction matters because some of the strongest sequence-model results come from broad ICU deterioration benchmarks rather than exclusively ventilated cohorts [11] [13].
**Confidence level: High** - explicit in the meta-synthesis and task-specific studies [1] [11].

### Outcome types used in the literature

Outcome labels vary substantially:

- **Extubation failure:** need for noninvasive ventilation, reintubation, or death within 48 hours after extubation [1] [8].
- **Weaning failure:** failure at or after a spontaneous breathing trial or weaning attempt [2] [53].
- **Successful weaning:** successful liberation within 14 days of ventilator support [7].
- **SBT-related successful weaning in multitask learning:** prediction tied to spontaneous-breathing-trial evaluation, alongside shock and mortality tasks [6].
- **Ventilator-associated complications:** CDC-defined VAC predicted after a defined ventilation duration, including a one-hour-ahead label after 35 hours of ventilation in one MIMIC-III study [3].
- **General deterioration:** respiratory or hemodynamic deterioration tagged by automated rules, or composite serious clinical worsening in broader ICU/ward literature [17] [20].

The literature therefore treats “condition change” as a family of related but nonidentical endpoints. That heterogeneity is clinically sensible, but it complicates benchmarking [1] [6] [20].
**Confidence level: High** - many studies explicitly define different targets [1] [3] [7].

## Data Sources, Cohorts, and Label Construction

### MIMIC-IV cohorts for ventilated ICU prediction

MIMIC-IV is a major source for recent ventilated-patient prediction research [1] [7]. Retrieved examples include:

- **Extubation failure in MIMIC-IV (2021):** 16,189 extubated ICU adults [8].
- **Weaning failure in MIMIC-IV (2022):** 3,695 mechanically ventilated patients ventilated for more than 24 hours and weaned using a T-tube strategy [4].
- **Weaning success in MIMIC-IV (2023):** 23,242 ventilated ICU patients [7].
- **Multitask weaning-related MIMIC-IV study (2024):** 7,758 critically ill ventilated patients, with 78.5% successfully weaned [6].

These cohorts show that MIMIC-IV supports both narrow liberation-focused tasks and broader multitask prognostic modeling [6] [8].
**Confidence level: High** - directly reported cohort details [4] [8].

### Label definitions and endpoint timing

Label construction is central because even closely related studies define failure differently [20].

Examples:

- **Extubation failure:** noninvasive ventilation, reintubation, or death within 48 hours after extubation [1] [8].
- **Weaning failure:** failure after ventilation/weaning attempt, often around the first spontaneous breathing trial [2] [53].
- **Weaning success:** success within 14 days of ventilator support [7].
- **VAC prediction:** label set one hour after 35 hours of ventilation [3].

Taken together, these findings suggest label design determines not only comparability, but also the operational meaning of the model output: some models predict immediate post-extubation collapse, while others predict longer-horizon liberation success [1] [7].
**Confidence level: High** - the label heterogeneity is explicit and repeatedly emphasized [1] [2] [20].

### Role of MIMIC-IV and other ICU databases

Beyond MIMIC-IV, the literature repeatedly uses:

- **MIMIC-III** for ventilator-associated complications, waveform studies, and broader ICU sequence modeling [3] [18] [19].
- **eICU** for external validation, automated labeling, and cross-dataset transfer studies [20] [29] [54].
- **AmsterdamUMCdb** in external validation of weaning readiness tools [28].
- **PhysioNet-derived resources** as foundational public critical-care infrastructure [33] [35].

However, broader reviews note overreliance on a small number of public ICU datasets, which limits confidence in generalizability [25] [38].
**Confidence level: High** - consistent across development and review evidence [25] [34].

## Features and Multimodal Inputs

### Vital signs, labs, and structured EHR variables

The dominant feature set is structured tabular ICU data. Common variables include heart rate, blood pressure, respiratory rate, oxygenation-related variables, routine labs, blood gases, demographics, comorbidities, and severity scores [5] [8] [55] [56].

The extubation-failure model in MIMIC-IV used **89 clinical and laboratory variables** [8]. Weaning-related studies used demographics, comorbidities, respiratory mechanics, ventilator settings, labs, vital signs, and blood gas values around the first spontaneous breathing trial [2] [53] [57].

A recurring design principle from broader deterioration literature is that **trend features** add value over current values alone [51] [55].
**Confidence level: High** - strong consistency across ventilator-specific and general ICU studies [8] [55].

### Ventilator settings and respiratory mechanics

Ventilation-specific features are increasingly prominent. Retrieved studies mention:

- Tidal volume [1]
- PEEP [1] [4]
- FiO2 / inspired oxygen concentration [4]
- Mean airway pressure [1]
- PSV level [1]
- Mechanical ventilation duration [1] [4]
- Dynamic lung compliance [4]
- Mechanical power [4]

The MIMIC-IV weaning-failure study found PEEP, dynamic compliance, mechanical power, FiO2, ICU length of stay, and ventilation duration to be independent predictors, with AUROC around 0.83 [4]. This indicates a shift toward respiratory-state-aware modeling rather than purely generic ICU scoring [4].
**Confidence level: Medium** - direct evidence supports the trend, but the number of respiratory-mechanics-rich studies in the retrieved set is still limited [4] [58].

### Waveforms, time-series summaries, and derived features

Some studies move beyond charted variables:

- Raw ventilator waveform pressure/flow data were summarized into nine physiologic features for ARDS screening, then modeled with random forest [18].
- A platform continuously detected eight patient-ventilator asynchrony subtypes and aggregated them into an Overall Asynchrony Index [37].

- A ventilator-alarm model started with 76 IoMT alarm parameters and reduced them to 13 features; CatBoost performed best [9].
- VAC prediction in MIMIC-III used manual summaries and tsfresh-generated time-series features; higher-frequency data and tsfresh features performed best [19].

Inferred: waveform-rich monitoring may capture short-term respiratory instability that structured EHR models can miss, but the retrieved evidence does not yet show routine integration of waveform-derived features into standard MIMIC-IV extubation/weaning pipelines. **Confidence level: Low** [18] [19] [37].

### Degree of true multimodal fusion

Although multimodal prediction is often claimed, the strongest direct evidence is for **combining multiple structured data types**, not for full fusion across structured EHR, ventilator waveforms, and unstructured text [1] [7]. Retrieved excerpts did not identify a clear model explicitly combining ventilator parameters, bedside vital signs, laboratory variables, and notes-derived text in one end-to-end MIMIC-IV architecture [40] [41].
**Confidence level: High** - this gap is directly stated in the evidence synthesis [40] [41].

## Prediction Models and Comparative Performance

### Classical models and tree-based baselines

In ventilated cohorts, tabular ML remains common:

- **XGBoost** modestly outperformed conventional statistical methods for poor post-ICU outcomes in mechanically ventilated adults [59].
- **Random forest** and **XGBoost** performed strongly in mechanically ventilated mortality prediction, with good calibration reported for most models except the neural network in one study [42].
- **CatBoost** was best in the ventilator-alarm management study after feature selection [9].
- **Logistic regression** remains a frequent baseline and sometimes part of ensemble comparisons [10] [60].

This pattern reflects the predominance of structured, relatively low-dimensional datasets in ventilated-patient studies [4] [9].
**Confidence level: High** - multiple ventilator-specific examples support this [9] [59].

### Sequence models for ICU time series

Broader ICU deterioration work shows strong performance from temporal deep learning:

- A deterioration-prediction RNN combining LSTM, GRU, attention, and temporal convolutions achieved AUC-ROC 0.891 versus 0.743 for traditional early-warning systems, with average lead time of 4.7 hours [11].
- An LSTM with input perseveration improved AUROC in the first hour from 0.77 to 0.83 and was 2-3 times more responsive to acute events than baseline LSTM [12].
- Attention-based TCN achieved AUROC 0.837 and AUPRC 0.454 in ICU mortality prediction, outperforming traditional AI and score-based comparators [13].
- Transformer-based and GRU/Transformer forecasting models have also been applied to short-term vital-sign prediction [14] [15] [61].

However, many of these strongest deep sequence results are not exclusive to mechanically ventilated subpopulations [11] [13].
**Confidence level: High** - the performance advantage is well-supported, and the task-specific caveat is explicitly noted in the synthesis [11] [13].

### Comparisons against scores and clinical tools

ML models are commonly benchmarked against traditional scoring systems and warning scores derived from repeated bedside measurements [47] [48]. Broader deterioration models have been compared to MEWS, NEWS, eCART, SOFA, SAPS-II, and APACHE-family scores [5] [47] [48] [50].

The consistent pattern is that automated ML models usually improve discrimination over static or simpler score-based rules, particularly when temporal data are used [11] [52] [62]. At the same time, traditional

scores remain useful predictors and clinical baselines inside modern models [5].
**Confidence level: High** - supported by both benchmark and ventilator-focused evidence [5] [11].

### Reported discrimination and early-warning gains

Selected reported results include:

- ANN from automatically recorded vital signs predicted deterioration around **10 hours before documentation**, AUC 0.81 [39].
- MIMIC-IV extubation-failure model: **AUROC 0.835** internally and **0.803** in prospective validation [8].
- MIMIC-IV weaning-failure model: **AUROC 0.828** in development and **0.833** in validation [4].
- Attention-enhanced temporal deterioration model: **AUROC 0.891**, average **4.7-hour lead time** [11].
- Higher-frequency tsfresh-based VAC prediction: **AUROC 0.83 ± 0.11** [19].
- Vent.io respiratory failure tool: **AUC 0.908**, though external transfer required fine-tuning after performance drop on MIMIC-IV [26].

Taken together, these findings suggest that useful performance is achievable, but whether models generalize without site-specific adaptation is less certain [25] [26].
**Confidence level: High** - direct performance figures are available, though not uniformly comparable because tasks differ [8] [26].

## Real-Time Deployment, Validation, and Clinical Limitations

### Missing data, irregular sampling, and class imbalance

Real-time ICU data are sparse and irregular. Reported strategies include:

- Excluding variables with >50% missingness, then using **iterative imputation followed by forward imputation** [21].
- Using recurrent/deep sequence models that can learn from irregular multivariate trajectories [63] [64].
- Benchmark comparisons using **forward filling/LOCF**, linear interpolation, random-forest imputation, MICE/MissForest, and deep imputation models [22] [23] [65].
- Handling class imbalance with **SMOTE**, **SMOTEENN**, random over/undersampling, and class weights [24] [31] [56] [66] [67].

The practical implication is that bedside readiness depends not only on architecture, but on how robustly the model handles clinically realistic missingness patterns [23] [68].
**Confidence level: High** - multiple studies directly address these methods [21] [22].

### Calibration, decision utility, and robustness

Several studies report calibration rather than AUROC alone:

- Calibration curves were reported for mortality and weaning-failure models [4] [42] [56].
- One deep model reported a **Brier score** and described itself as well calibrated [69].
- **Decision-curve analysis** was reported for the weaning-failure and ICU AKI models [4] [69].

Yet the retrieved evidence did not identify explicit temporal drift checks in these ventilated ICU studies [4] [42] [56] [69]. This is an important deployment gap, because models may remain discriminative while becoming miscalibrated over time [26] [36].
**Confidence level: Medium** - calibration evidence is direct, but drift evidence is mainly absent rather than deeply characterized [43] [69].

### External validation and domain shift

External validation is still uncommon in ICU deterioration ML; one meta-analysis found only 14.7% of studies externally validated, and AUROC typically decreased in new hospitals [25].

Ventilator-specific examples reinforce the problem:

- A real-time weaning-readiness tool externally validated across MIMIC-IV and AmsterdamUMCdb had AUROCs of **0.847/0.795/0.789** for 24/48/72-hour prediction in the external cohort, with specificity falling from **0.79 to 0.63**, and lower performance in older and neurosurgical patients [28].
- Vent.io showed good development performance but performance dropped on MIMIC-IV and required fine-tuning [26].
- A clustering study explicitly found significant clinical differences between eICU derivation and MIMIC-IV validation cohorts, demonstrating domain shift [29].

These findings indicate that dependence on MIMIC-IV alone is insufficient for bedside confidence [25] [29].
**Confidence level: High** - direct cross-site evidence and review data converge [25] [28].

### Interpretability and workflow integration

Interpretability tools are now common:

- SHAP, PDP, and LIME were used for extubation prediction, including domain-, feature-, and individual-level explanations [27].
- SHAP-supported interpretable risk calculators were proposed for extubation failure and personalized ventilation management [30].
- SHAP-based feature reduction showed that dropping nearly 60% of features left AUC largely unchanged in one deployment-focused model [31].
- Dynamic nomogram-style outputs and feature selection were also used in invasive mechanical ventilation settings [70] [71].

However, the retrieved evidence shows almost no formal evaluation of clinician trust, adoption, alert fatigue, or workflow usability [27] [30] [32]. So interpretability is being engineered, but usability is rarely measured.
**Confidence level: High** - the gap is explicit across several explainability papers [27] [32].

## Synthesis: What the Literature Shows Overall

Three main patterns emerge.

First, the field has clearly moved beyond static scores toward **temporally aware prediction** using ICU streams and repeated measurements [11] [52]. Second, in ventilated patients specifically, the strongest published models still often depend on **structured EHR and ventilator-setting features**, with only emerging use of waveform and respiratory-mechanics signals [1] [4] [18]. Third, translation to bedside care is constrained more by **data engineering, labeling, and workflow issues** than by lack of predictive power alone [22] [25] [27].

Inferred: the most clinically deployable next-generation systems may be compact, continuously updated models that rely on fast-available monitor and ventilator variables rather than delayed or documentation-dependent features, because such inputs better match real-time operational constraints [68] [72] [73].
**Confidence level: Low** - plausible cross-study inference, but not directly tested head-to-head in the retrieved set.

## Practical Implications

- For **researchers**, MIMIC-IV is an important starting point for extubation and weaning prediction, but external validation should be designed from the outset [8] [28].
- For **model developers**, strong baselines should include XGBoost/random forest as well as temporal deep models, because task-specific gains are not uniform [4] [11].
- For **clinical deployment teams**, calibration, data latency, and charting workflows may matter more than a small AUROC improvement [4] [73].
- For **future ventilator monitoring systems**, integrating respiratory mechanics and waveform-derived physiology with structured EHR prediction appears promising but remains underdeveloped [4] [19] [37].
  **Confidence level: Medium** - grounded in the evidence, though some implications are cross-study synthesis rather than direct experimental comparisons [4] [25].

## Conclusion

Machine learning models analyze mechanically ventilated ICU patients mainly through structured vital signs, laboratory data, blood gases, demographics, comorbidities, and ventilator variables to predict extubation failure, weaning failure, liberation success, ventilator-associated complications, and broader deterioration [1] [2] [3] [7]. MIMIC-IV has become a major dataset for this work, especially for extubation and weaning tasks, while MIMIC-III, eICU, AmsterdamUMCdb, and PhysioNet resources support complementary development and validation [4] [8] [28] [33]. Methodologically, ventilated-patient studies often use XGBoost, random forest, logistic regression, and CatBoost, whereas broader ICU deterioration benchmarks frequently show added value from LSTM, GRU, attention, TCN, and transformer models [9] [11] [13]. The main obstacle to real-time clinical use is not lack of predictive modeling capacity, but unreliable transfer to bedside conditions due to heterogeneous labels, missing and irregular data, calibration and drift issues, weak multicenter validation, and limited workflow evidence [20] [23] [25] [27].

**Overall confidence level: High** - the answer is supported by multiple directly relevant ventilator-specific studies, broader ICU sequence-model evidence, and several deployment/validation reviews [4] [8] [11] [25].

## References


1. Zhao, Q.-Y., Wang, H., Luo, J.-C., (...), Luo, Z. (2021) Development and Validation of a Machine-Learning Model for Prediction of Extubation Failure in Intensive Care Units. *Frontiers in Medicine*. https://doi.org/10.3389/fmed.2021.676343
2. Yan, Y., Luo, J., Wang, Y., (...), Li, X. (2022) Development and validation of a mechanical power-oriented prediction model of weaning failure in mechanically ventilated patients: a retrospective cohort study. *BMJ Open*. https://doi.org/10.1136/bmjopen-2022-066894
3. Tan, Y., Young, M., Girish, A., (...), Bergmann, J. (2023) Predicting respiratory decompensation in mechanically ventilated adult ICU patients. *Frontiers in Physiology*. https://doi.org/10.3389/fphys.2023.1125991
4. Yan, Yao, Luo, Jiye, Wang, Yanli, (...), Li, Xiaomin (2022) Development and validation of a mechanical power-oriented prediction model of weaning failure in mechanically ventilated patients: a retrospective cohort study. *BMJ Open*. https://doi.org/10.1136/bmjopen-2022-066894
5. Zhu, Y., Zhang, J., Wang, G., (...), Yu, Q. (2021) Machine Learning Prediction Models for Mechanically Ventilated Patients: Analyses of the MIMIC-III Database. *Frontiers in Medicine*. https://doi.org/10.3389/fmed.2021.662340
6. Lin, M.-Y., Chi, H.-Y., Chao, W.-C. (2024) Multitask learning to predict successful weaning in critically ill ventilated patients: A retrospective analysis of the MIMIC-IV database. *Digital Health*. https://doi.org/10.1177/20552076241289732
7. Kim, J., Kim, Y.K., Kim, H., (...), Kim, H.-J. (2023) Machine Learning Algorithms Predict Successful Weaning From Mechanical Ventilation Before Intubation: Retrospective Analysis From the Medical Information Mart for Intensive Care IV Database. *JMIR Formative Research*. https://doi.org/10.2196/44763
8. Zhao, Qin-Yu, Wang, Huan, Luo, Jing-Chao, (...), Luo, Zhe (2021) Development and Validation of a Machine-Learning Model for Prediction of Extubation Failure in Intensive Care Units. *Frontiers in Medicine*. https://doi.org/10.3389/fmed.2021.676343
9. Wang, X., Chen, Z., Zhang, H., (...), Huang, J. (2025) Development and validation of a machine learning and Internet of Medical Things-based model for ICU ventilator alarm management. *Chinese Journal of Evidence-Based Medicine*. https://doi.org/10.7507/1672-2531.202411199
10. Kivelev, J.V., Krivoshapkin, A.L., Sufianov, A.A. (2024) Machine learning algorithm to predict in-hospital mortality after aneurysmal subarachnoid hemorrhage. *Sechenov Medical Journal*. https://doi.org/10.47093/2218-7332.2024.15.4.19-31
11. Swami, P.P., Sharma, A. (2027) Real-Time Patient Deterioration Prediction in Intensive Care Units Using Recurrent Neural Networks. *Lecture Notes in Networks and Systems*. https://doi.org/10.1007/978-3-032-32476-4_48
12. Ledbetter, D.R., Laksana, E., Aczon, M., Wetzel, R. (2021) Improving Recurrent Neural Network Responsiveness to Acute Clinical Events. *IEEE Access*. https://doi.org/10.1109/ACCESS.2021.3099996

13. Chen, Y.-W., Li, Y.-J., Deng, P., (...), Yi, B. (2022) Learning to predict in-hospital mortality risk in the intensive care unit with attention-based temporal convolution network. *BMC Anesthesiology*. https://doi.org/10.1186/s12871-022-01625-5

14. Harerimana, G., Kim, J.W., Jang, B. (2022) A Multi-Headed Transformer Approach for Predicting the Patient's Clinical Time-Series Variables From Charted Vital Signs. *IEEE Access*. https://doi.org/10.1109/ACCESS.2022.3211334

15. Hinrichs, N., Roeschl, T., Lanmueller, P., (...), Meyer, A. (2024) Short-term vital parameter forecasting in the intensive care unit: A benchmark study leveraging data from patients after cardiothoracic surgery. *PLOS Digital Health*. https://doi.org/10.1371/journal.pdig.0000598

16. Fu, L.-H., Knaplund, C., Cato, K., (...), Collins Rossetti, S. (2021) Utilizing timestamps of longitudinal electronic health record data to classify clinical deterioration events. *Journal of the American Medical Informatics Association*. https://doi.org/10.1093/jamia/ocab111

17. Churpek, M.M., Yuen, T.C., Winslow, C., (...), Edelson, D.P. (2016) Multicenter Comparison of Machine Learning Methods and Conventional Regression for Predicting Clinical Deterioration on the Wards. *Critical Care Medicine*. https://doi.org/10.1097/CCM.0000000000001571

18. Rehm, G.B., Cortes-Puch, I., Kuhn, B.T., (...), Adams, J.Y. (2021) Use of Machine Learning to Screen for Acute Respiratory Distress Syndrome Using Raw Ventilator Waveform Data. *Critical Care Explorations*. https://doi.org/10.1097/CCE.0000000000000313

19. Tan, Yvette, Young, Michael, Girish, Akanksha, (...), Bergmann, Jules (2023) Predicting respiratory decompensation in mechanically ventilated adult ICU patients. *Frontiers in Physiology*. https://doi.org/10.3389/fphys.2023.1125991

20. Jeddah, D., Chen, O., Lipsky, A.M., (...), Pessach, I.M. (2021) Validation of an automatic tagging system for identifying respiratory and hemodynamic deterioration events in the intensive care unit. *Healthcare Informatics Research*. https://doi.org/10.4258/HIR.2021.27.3.241

21. Jung, J., Kang, S., Choi, J., (...), Kim, H. (2025) Evaluating the impact of explainable AI on clinicians' decision-making: A study on ICU length of stay prediction. *International Journal of Medical Informatics*. https://doi.org/10.1016/j.ijmedinf.2025.105943

22. Fujarski, M., Porschen, C., Plagwitz, L., (...), Varghese, J. (2023) DeepTSE: A Time-Sensitive Deep Embedding of ICU Data for Patient Modeling and Missing Data Imputation. *Studies in Health Technology and Informatics*. https://doi.org/10.3233/SHTI230110

23. Poëtte, M., Mouysset, S., Ruíz, D., (...), Minville, V. (2026) Benchmarking imputation strategies for missing time-series data in critical care using real-world-inspired scenarios. *Scientific Reports*. https://doi.org/10.1038/s41598-026-39035-z

24. Nguyen, Q.T., Tran, M.P., Prabhakaran, V., (...), Nguyen, G.H. (2024) Compact machine learning model for the accurate prediction of first 24-hour survival of mechanically ventilated patients. *Frontiers in Medicine*. https://doi.org/10.3389/fmed.2024.1398565

25. Rockenschaub, P., Akay, E.M., Carlisle, B.G., (...), Madai, V.I. (2025) External validation of AI-based scoring systems in the ICU: a systematic review and meta-analysis. *BMC Medical Informatics and Decision Making*. https://doi.org/10.1186/s12911-024-02830-7

26. Lam, J.Y., Lu, X., Shashikumar, S.P., (...), Nemati, S. (2024) Development, deployment, and continuous monitoring of a machine learning model to predict respiratory failure in critically ill patients. *JAMIA Open*. https://doi.org/10.1093/jamiaopen/ooae141

27. Pai, K.-C., Su, S.-A., Chan, M.-C., (...), Chao, W.-C. (2022) Explainable machine learning approach to predict extubation in critically ill ventilated patients: a retrospective study in central Taiwan. *BMC Anesthesiology*. https://doi.org/10.1186/s12871-022-01888-y

28. Zappalà, S., Scaravilli, V., Rovati, L., (...), Grasselli, G. (2025) Development and validation of a machine learning model for real-time prediction of invasive mechanical ventilation weaning readiness. *Journal of Critical Care*. https://doi.org/10.1016/j.jcrc.2025.155105

29. Chen, Xuanhui, Li, Jiaxin, Liu, Guangjian, (...), Liang, Huiying (2023) Identification of Distinct Clinical Phenotypes of Heterogeneous Mechanically Ventilated ICU Patients Using Cluster Analysis. *Journal of Clinical Medicine*. https://doi.org/10.3390/jcm12041499

30. Zhang, T., Lu, T., Yang, S. (2026) Predicting extubation failure in patients with cardiorespiratory decompensation: development and validation of an interpretable machine learning–based risk calculator. *Frontiers in Cardiovascular Medicine*. https://doi.org/10.3389/fcvm.2026.1862117
31. Alvarez, R., Chen, J., Nguyen, Q.T. (2024) Efficient Machine Learning Model Deployment in Clinical Decision Support Systems. *Canadian Conference on Electrical and Computer Engineering*. https://doi.org/10.1109/CCECE59415.2024.10667100
32. Lin, M.-Y., Li, C.-C., Lin, P.-H., (...), Chao, W.-C. (2021) Explainable Machine Learning to Predict Successful Weaning Among Patients Requiring Prolonged Mechanical Ventilation: A Retrospective Cohort Study in Central Taiwan. *Frontiers in Medicine*. https://doi.org/10.3389/fmed.2021.663739
33. Goldberger, A.L., Nunes Amaral, L.A., Glass, L., (...), Stanley, H.E. (2000) PhysioBank, PhysioToolkit, and PhysioNet: components of a new research resource for complex physiologic signals.. *Circulation*. https://www.scopus.com/pages/publications/14844283547
34. Johnson, A.E.W., Pollard, T.J., Shen, L., (...), Mark, R.G. (2016) MIMIC-III, a freely accessible critical care database. *Scientific Data*. https://doi.org/10.1038/sdata.2016.35
35. Pollard, T.J., Johnson, A.E.W., Raffa, J.D., (...), Badawi, O. (2018) The eICU collaborative research database, a freely available multi-center database for critical care research. *Scientific Data*. https://doi.org/10.1038/sdata.2018.178
36. Zimmerman, J.E., Kramer, A.A., McNair, D.S., Malila, F.M. (2006) Acute Physiology and Chronic Health Evaluation (APACHE) IV: Hospital mortality assessment for today's critically ill patients. *Critical Care Medicine*. https://doi.org/10.1097/01.CCM.0000215112.84523.F0
37. Chen, X., Yuan, S., Baedorf Kassis, E.B., (...), Long, Y. (2025) Methodological development of the remote ventilate view platform for real-time monitoring of patient-ventilator asynchrony and respiratory parameters in severe pneumonia patients. *Journal of Intensive Medicine*. https://doi.org/10.1016/j.jointm.2025.07.003
38. Ji, Q., Wu, Y., Yang, M., (...), Hu, W. (2026) Integration, challenges, and future of artificial intelligence in critical care medicine: comprehensive applications from predictive models to clinical integration. *Frontiers in Medicine*. https://doi.org/10.3389/fmed.2026.1874474
39. Thiele, D., Rodseth, R., Friedland, R., (...), Nee, J. (2025) Machine Learning Models for the Early Real-Time Prediction of Deterioration in Intensive Care Units—A Novel Approach to the Early Identification of High-Risk Patients. *Journal of Clinical Medicine*. https://doi.org/10.3390/jcm14020350
40. Martin, J., Afshar, M., Afshar, A.S., (...), Churpek, M.M. (2026) Explainable multimodal deep learning models for variable-length sequences in critically ill patients. *Journal of Biomedical Informatics*. https://doi.org/10.1016/j.jbi.2026.105001
41. Lim, Leerang, Lee, Hyeonhoon, Jung, Chul-Woo, (...), Lee, Hyung-Chul (2024) INSPIRE, a publicly available research dataset for perioperative medicine. *Scientific data*. https://doi.org/10.1038/s41597-024-03517-4
42. Zhu, Yibing, Zhang, Jin, Wang, Guowei, (...), Yu, Qian (2021) Machine Learning Prediction Models for Mechanically Ventilated Patients: Analyses of the MIMIC-III Database. *Frontiers in Medicine*. https://doi.org/10.3389/fmed.2021.662340
43. Bansal, Vikas, Smischney, Nathan J., Kashyap, Rahul, (...), Freeman, William David (2021) Reintubation Summation Calculation: A Predictive Score for Extubation Failure in Critically Ill Patients. *Frontiers in Medicine*. https://doi.org/10.3389/fmed.2021.789440
44. Ferreira, F., Peres-Bota, D., Bross, A., (...), Vincent, J.-L. (2001) Serial evaluation of the SOFA score to predict outcome in critically ill patients. *JAMA*. https://doi.org/10.1001/jama.286.14.1754
45. Knaus, W.A., Draper, E.A., Wagner, D.P., Zimmerman, J.E. (1985) APACHE II: A severity of disease classification system. *Critical Care Medicine*. https://doi.org/10.1097/00003246-198510000-00009
46. Knaus, W.A., Wagner, D.P., Draper, E.A., (...), Janssens, F.E. (1991) The APACHE III prognostic system: Risk prediction of hospital mortality for critically III hospitalized adults. *Chest*. https://doi.org/10.1378/chest.100.6.1619

47. Subbe, C.P., Kruger, M., Rutherford, P., Gemmel, L. (2001) Validation of a modified early warning score in medical admissions. *QJM: An International Journal of Medicine*. https://doi.org/10.1093/qjmed/94.10.521

48. Smith, G.B., Prytherch, D.R., Meredith, P., (...), Featherstone, P.I. (2013) The ability of the National Early Warning Score (NEWS) to discriminate patients at risk of early cardiac arrest, unanticipated intensive care unit admission, and death. *Resuscitation*. https://doi.org/10.1016/j.resuscitation.2012.12.016

49. Prytherch, D.R., Smith, G.B., Schmidt, P.E., Featherstone, P.I. (2010) ViEWS-Towards a national early warning score for detecting adult inpatient deterioration. *Resuscitation*. https://doi.org/10.1016/j.resuscitation.2010.04.014

50. Kipnis, P., Turk, B.J., Wulf, D.A., (...), Escobar, G.J. (2016) Development and validation of an electronic medical record-based alert score for detection of inpatient deterioration outside the ICU. *Journal of Biomedical Informatics*. https://doi.org/10.1016/j.jbi.2016.09.013

51. Rothman, M.J., Rothman, S.I., Beals, J. (2013) Development and validation of a continuous measure of patient condition using the Electronic Medical Record. *Journal of Biomedical Informatics*. https://doi.org/10.1016/j.jbi.2013.06.011

52. Meyer, A., Zverinski, D., Pfahringer, B., (...), Eickhoff, C. (2018) Machine learning for real-time prediction of complications in critical care: a retrospective study. *The Lancet Respiratory Medicine*. https://doi.org/10.1016/S2213-2600(18)30300-X

53. Yan, Y., Xie, Y., Du, Z., (...), Li, X. (2025) Correlation analysis between mechanical power normalized to dynamic lung compliance and weaning outcomes and prognosis in mechanically ventilated patients: a prospective, observational cohort study. *Chinese Critical Care Medicine*. https://doi.org/10.3760/cma.j.cn121430-20240126-00085

54. Gao, Shicheng, Zhang, Yunhai, Deng, Menghua, (...), Zhang, Bin (2025) Machine learning-based early warning system for hemodynamic deterioration in cardiovascular ICU patients: a bidirectional cross-validation study. *Frontiers in Cardiovascular Medicine*. https://doi.org/10.3389/fcvm.2025.1694001

55. Churpek, M.M., Adhikari, R., Edelson, D.P. (2016) The value of vital sign trends for detecting clinical deterioration on the wards. *Resuscitation*. https://doi.org/10.1016/j.resuscitation.2016.02.005

56. Ashrafi, N., Liu, Y., Xu, X., (...), Pishgar, M. (2024) Deep learning model utilization for mortality prediction in mechanically ventilated ICU patients. *Informatics in Medicine Unlocked*. https://doi.org/10.1016/j.imu.2024.101562

57. Yan, Y., Xie, Y., Luo, J., (...), Li, X. (2023) Development and validation of a mechanical power-oriented nomogram model for predicting the risk of weaning failure in mechanically ventilated patients: an analysis using the data from MIMIC-Ⅳ. *Chinese Critical Care Medicine*. https://doi.org/10.3760/cma.j.cn121430-20221115-00997

58. Tasci, S.K., Ozturk, M. (2026) Dynamic Clustering-Based Clinical Decision Support Approach for Patient Phenotyping in Mechanical Ventilation. *SIU 2026 - 34th Signal Processing and Communications Applications Conference, Conference Proceedings*. https://doi.org/10.1109/SIU71813.2026.11636510

59. Ong, W.J.D., How, C.H., Chong, W.H.K., (...), Kansal, A. (2024) Outcome prediction for adult mechanically ventilated patients using machine learning models and comparison with conventional statistical methods: A single-centre retrospective study. *Intelligence-Based Medicine*. https://doi.org/10.1016/j.ibmed.2024.100165

60. Arvelo, I.C., Shipley, K., Wright, A., Steitz, B.D. (2026) Machine learning on asynchronous clinical pages to predict clinical deterioration. *JAMIA Open*. https://doi.org/10.1093/jamiaopen/ooag122

61. Jahan, F., Hossain, N., Ifty, R.A. (2025) Beyond Binary Alerts: A Multi-Faceted Framework for Clinically Actionable Sepsis Prediction. *2025 28th International Conference on Computer and Information Technology, ICCIT 2025*. https://doi.org/10.1109/ICCIT68739.2025.11491596

62. Pirracchio, R., Petersen, M.L., Carone, M., (...), van der Laan, M.J. (2015) Mortality prediction in intensive care units with the Super ICU Learner Algorithm (SICULA): A population-based study. *The Lancet Respiratory Medicine*. https://doi.org/10.1016/S2213-2600(14)70239-5

63. Wang, Yonghua, Bai, Yuling, Jin, Ge (2026) Explainable deep-learning models to predict diaphragmatic dysfunction and cognitive stress in ICU patients under mechanical ventilation. *Frontiers in Physiology*. https://doi.org/10.3389/fphys.2026.1765898

64. Herath, M., Andersen, M. (2026) A continuous-time predictive analytics framework for quantifying information degradation in intensive care monitoring systems. *Healthcare Analytics*. https://doi.org/10.1016/j.health.2026.100497

65. Digitale, J., Franzon, D., Pletcher, M.J., (...), Gennatas, E.D. (2025) Methods for Addressing Missingness in Electronic Health Record Data for Clinical Prediction Models: Comparative Evaluation. *JMIR Medical Informatics*. https://doi.org/10.2196/79307

66. Tang, J., Wang, X., Wan, H., (...), Du, Y. (2022) Joint modeling strategy for using electronic medical records data to build machine learning models: an example of intracerebral hemorrhage. *BMC Medical Informatics and Decision Making*. https://doi.org/10.1186/s12911-022-02018-x

67. Huang, K.-Y., Hsu, Y.-L., Chen, H.-C., (...), Hou, M.-H. (2023) Developing a machine-learning model for real-time prediction of successful extubation in mechanically ventilated patients using time-series ventilator-derived parameters. *Frontiers in Medicine*. https://doi.org/10.3389/fmed.2023.1167445

68. Yeche, H., Kuznetsova, R., Zimmermann, M., (...), Rätsch, G. (2021) HiRID-ICU-Benchmark — A Comprehensive Machine Learning Benchmark on High-resolution ICU Data. *Advances in Neural Information Processing Systems*. https://www.scopus.com/pages/publications/105000455917

69. Zhang, Li, Li, Mingyu, Wang, Chengcheng, (...), Wu, Hong (2025) Prediction of acute kidney injury in intensive care unit patients based on interpretable machine learning. *DIGITAL HEALTH*. https://doi.org/10.1177/20552076241311173

70. Zhao, Y., Chen, S., Pang, C., (...), Lin, S. (2026) Development and external validation of a machine-learning model for classifying documented pressure injury status during ICU stays among patients receiving invasive mechanical ventilation: A MIMIC-IV and eICU study. *Journal of Tissue Viability*. https://doi.org/10.1016/j.jtv.2026.101058

71. Pan, G., Wang, W., Yang, G., Liu, H. (2026) Development and validation of an interpretable machine learning model for predicting sepsis risk in pleural effusion patients. *Journal of Thoracic Disease*. https://doi.org/10.21037/jtd-2026-0824

72. Akel, M.A., Carey, K.A., Winslow, C.J., (...), Edelson, D.P. (2021) Less is more: Detecting clinical deterioration in the hospital with machine learning using only age, heart rate, and respiratory rate. *Resuscitation*. https://doi.org/10.1016/j.resuscitation.2021.08.024

73. Churpek, M.M., Carey, K.A., Snyder, A., (...), Edelson, D.P. (2025) Multicenter Development and Prospective Validation of eCARTv5: A Gradient-Boosted Machine-Learning Early Warning Score. *Critical Care Explorations*. https://doi.org/10.1097/CCE.0000000000001232