Predicting respiratory deterioration in ICU patients

"How can machine learning and artificial intelligence predict early respiratory deterioration in mechanically ventilated ICU patients using vital signs and clinical data? What methods are used, how accurately can they predict deterioration, and what limitations remain?"

Generated by LeapSpace, Sat Oct 10 2026

Early Respiratory Deterioration in Mechanically Ventilated ICU Patients: Quick Insights

Key Findings Table

Key finding	Population / task	Evidence
Most published ICU ML models for ventilated adults do **not** predict imminent respiratory rescue events directly; they usually predict proxies such as prolonged mechanical ventilation, weaning/extubation outcomes, mortality, VAP, or future intubation need.	Adult ICU and mechanically ventilated cohorts	[1] [2] [3] [4] [5] [6] [7] [8] [9] [10] [11] [12]
Tree-based structured-EHR models are the practical workhorse, especially XGBoost and random forest; they often equal or exceed logistic regression on routine ICU data.	Ventilated adult ICU prediction studies using vitals, labs, severity scores, and sometimes ventilator settings	[6] [13] [14] [15] [16] [17] [18] [19]
Common high-value predictors include respiratory rate, oxygenation metrics, ABGs, PEEP, FiO2, plateau pressure, tidal volume, pH/PaCO2, plus broader severity markers such as SOFA/APACHE, vasopressor use, lactate, and renal variables.	Ventilated ICU adults across mortality, weaning, ARDS, and prolonged ventilation tasks	[1] [2] [3] [4] [20] [21] [22] [23] [24] [25]
Internal discrimination is often good, commonly around AUROC 0.80-0.90 for related ventilated-patient outcomes, but external validation is scarce and performance/calibration often worsen at new sites.	Ventilated adult ICU cohorts and broader ICU validation reviews	[1] [6] [17] [18] [26] [27] [28] [29] [30]
Temporal and multimodal models are emerging, including LSTM/GRU and EHR-plus-notes/waveform/radiomics systems, but consistent superiority over strong structured-data baselines is not established for this exact use case.	ICU time-series and multimodal prediction studies	[31] [32] [33] [34] [35] [36]
Implementation evidence lags far behind model development: few studies report real-time deployment, PPV at usable thresholds, alert burden, clinician response, or workflow impact.	Ventilated ICU AI and broader deterioration-alert literature	[5] [37] [38] [39] [40] [41] [42] [43] [44]

Direct Answer

Machine learning can predict early respiratory deterioration in mechanically ventilated ICU patients primarily by learning from **structured longitudinal ICU data**: vital signs, oxygenation measures, ventilator settings, blood gases, laboratory values, and severity scores, sometimes enhanced by notes, waveforms,

or imaging [13] [16] [17] [18] [35]. In current practice, the most consistently effective methods are **tree-based models** such as XGBoost, random forest, LightGBM, and related ensembles, with logistic regression remaining a strong baseline and LSTM/GRU-style models used when richer time-series modeling is attempted [6] [17] [18] [31] [32].

Reported accuracy is often promising for **related** ventilated-patient outcomes: many internal studies report AUROCs around **0.80-0.90**, including about **0.821** for XGBoost in ventilated-patient mortality prediction, **0.79** for 30-day mortality in another cohort, and roughly **0.805-0.872** for difficult-weaning prediction [6] [17] [18] [27] [28]. However, for the precise question of predicting **imminent respiratory deterioration during ongoing mechanical ventilation**—for example re-intubation, escalation of ventilatory support, ARDS worsening, or urgent rescue within 6-48 hours—the evidence base is thin, and few retrieved studies directly match that target [3] [10] [11] [12] [16].

The main limitations are not only technical but definitional and operational: studies use inconsistent outcome labels, variable prediction windows, mostly retrospective single-center data, limited external validation, imperfect calibration after transport, missing data, class imbalance, and sparse evidence on bedside usefulness or false-alarm burden [26] [30] [45] [46] [47] [48] [49] [50]. Taken together, these findings suggest that ML is already useful for **risk stratification of ventilated ICU trajectories**, but is **not yet mature** for reliable, standardized short-horizon respiratory rescue alerts across hospitals [17] [19] [30] [40].

Confidence level: High - The broad pattern is strongly repeated across the supplied evidence, although exact short-horizon respiratory-event studies are sparse [1] [17] [26] [50].

Study Scope

This report synthesizes evidence from adult ICU and mechanically ventilated patient studies focused on ML/AI prediction using routine clinical data, including structured EHR variables, time-series vitals, labs, blood gases, severity scores, and selected multimodal additions such as notes, waveforms, and imaging [17] [35] [36]. It covers prediction targets including prolonged ventilation, extubation/weaning outcomes, mortality, ARDS-related outcomes, invasive ventilation need, and broader deterioration alerts, along with validation, calibration, interpretability, and implementation issues [1] [4] [51] [52]. The evidence spans retrospective cohorts, multicenter validations, systematic/scoping reviews, and a small number of real-time deployment examples [26] [42] [50].

Assumptions & Limitations

The key limitation is that the published evidence only partially matches the user's exact use case: many studies involve ventilated ICU patients, but relatively few define deterioration specifically as **near-term respiratory worsening while already ventilated** [10] [12]. Outcome definitions vary substantially, especially for extubation or weaning failure, including whether noninvasive ventilation counts as failure and whether the window is 24, 48, 72 hours, or longer, which limits comparability [45] [46] [47] [53]. External validation, calibration reporting, and deployment metrics are incomplete in much of the literature [26] [42].

Suggested Further Research

- Standardize a consensus definition of **early respiratory deterioration** in already ventilated ICU adults, distinct from mortality or prolonged ventilation [45] [54].
- Build hourly rolling-window models for explicit 6-, 12-, 24-, and 48-hour respiratory events such as re-intubation, rescue escalation, or hypoxemic worsening [12] [55].
- Require external, temporal, and prospective validation with calibration repair and decision-curve analysis [26] [30] [56].
- Report operational metrics needed for implementation, including PPV, alert burden, clinician uptake, and workflow effects [42] [43].
- Compare strong structured-EHR baselines against temporal deep learning and multimodal fusion in the same ventilated cohorts [17] [31] [35].

Full Report

Foundational Context

What “early respiratory deterioration” means in this literature

In theory, early respiratory deterioration in a mechanically ventilated ICU patient should mean a **near-term worsening in respiratory status that requires action**, such as ventilator escalation, failure to maintain oxygenation/ventilation, re-intubation after extubation, or urgent rescue before collapse [57] [58]. In practice, the ML literature often substitutes downstream or proxy endpoints such as prolonged ventilation, extubation failure, difficult weaning, ICU mortality, VAP, or need for initial invasive ventilation [1] [2] [3] [4] [8] [9].

That distinction matters because a model predicting **30-day mortality** answers a different clinical question from a model predicting **respiratory decompensation in the next 12 hours** [57] [59]. The field therefore blends at least two use cases:

1. **Short-horizon alerting** for imminent deterioration and rescue [1] [16] [55].
2. **Trajectory prediction** for prolonged ventilation, weaning failure, or mortality planning [3] [11] [60].

Confidence level: High - This framing is directly supported by repeated contrasts between early-deterioration endpoints and longer-horizon prognostic outcomes [12] [57].

Prediction Targets and Outcome Definitions

What models are actually predicting

Among mechanically ventilated ICU adults, common ML targets include prolonged mechanical ventilation, extubation or weaning success/failure, ICU or hospital mortality, ventilator-associated complications, and related respiratory-risk states [1] [2] [3] [4] [5] [6] [7] [8] [9]. Reviews explicitly note that many studies presented as “respiratory deterioration” work are in fact modeling these proxy outcomes rather than true near-term rescue events [3] [10] [11] [12].

For the user’s exact target—already ventilated ICU adults with deterioration defined as re-intubation, escalation of ventilatory support, ARDS worsening, or urgent ICU rescue within 6-48 hours—no retrieved paper clearly met all criteria simultaneously [10] [12] [16] [61].

Why label choice changes interpretation

Extubation and weaning studies illustrate the problem well. “Failure” may mean re-intubation within 24 hours, 48 hours, 72 hours, or 7 days; some definitions also include noninvasive positive-pressure ventilation, while death is usually reported as a consequence rather than included in the formal label [46] [47] [53] [62] [63] [64]. A review concluded that these original definitions could not be standardized and recommended a consensus core outcome such as re-intubation within 48 hours with broader criteria handled separately [45].

This heterogeneity makes reported model accuracy hard to compare. A model that predicts “extubation success at 48 hours” is not directly comparable to one predicting “weaning failure” or “prolonged ventilation >14 days” [65] [66] [67].

Prediction windows and lead time

Prediction windows vary widely. Some models use admission or the first ICU day, others use 24 or 48 hours after diagnosis, and some evaluate 72-hour landmarks after ARDS diagnosis [3] [4] [8] [11]. Rolling hourly windows are used in broader ICU deterioration modeling, where predictions are updated each hour for outcomes such as death within 24 hours or ventilator onset within 4 hours [55]. But comparable hourly designs for explicit respiratory deterioration in already ventilated adults remain scarce [12].

Confidence level: High - Outcome and horizon heterogeneity is one of the clearest findings across the supplied evidence [1] [45] [46].

Methods and Model Architectures

Classical machine learning

The dominant deployed-style approach uses structured EHR features with conventional supervised models. Across ventilated adult ICU cohorts, studies compare logistic regression, random forest, XGBoost,

LightGBM, SVM, bagging, KNN, and multilayer perceptrons [6] [17] [18]. Tree-based methods repeatedly perform strongly:

- In a MIMIC-III mechanically ventilated cohort, XGBoost had the best AUC at **0.821**, narrowly ahead of random forest (**0.819**) and logistic regression (**0.818**) [17].
- In a 30-day mortality study, XGBoost reached **0.79**, slightly ahead of balanced random forest at **0.78** [6].
- For difficult weaning, random forest performed best with **AUC 0.805** and **accuracy 0.748** [18].

These results suggest that on routine ICU data, careful feature curation and split design may matter more than using highly complex architectures [17] [19].

Deep learning and temporal sequence models

Deep models are increasingly used for sequential ICU data, including LSTM, GRU, temporal CNN, and Transformer-style models [31] [32] [68]. In broader ICU tasks, LSTM/GRU often show strong discrimination and temporal CNNs can be competitive on AUPRC-focused tasks [31]. Transformer models also perform strongly in some ICU prediction settings and can outperform classical comparators in broader prognostic tasks [32] [68].

However, the supplied evidence does **not** show a clear head-to-head superiority of these deep sequence models for the exact task of respiratory worsening in already ventilated adults [31] [32] [68]. The strongest evidence still favors robust structured-data pipelines as the current practical baseline [17] [18].

Multimodal and fusion approaches

Multimodal work is promising. Examples include:

- EHR plus nursing notes for respiratory failure prediction, improving AUC by about **6.8%** and F1 by about **5.8%** versus single-modality/conventional baselines [33].
- EHR plus chest CT radiomics for ARDS mortality, improving AUC from **0.758** to **0.813** [34].
- EHR plus physiologic waveforms outperforming EHR-only and waveform-only models in postoperative ICU deterioration prediction [36].
- Structured EHR plus notes improving event prediction, including ventilator onset, especially at longer horizons [35].

Taken together, these findings suggest multimodal fusion can add value, but the evidence is still too sparse to call it established for ventilated respiratory deterioration specifically [33] [35].

Confidence level: Medium-High - Classical model evidence is strong; deep and multimodal superiority remains less certain in the ventilated-respiratory niche [17] [31] [34].

Feature Inputs and Feature Engineering

Core predictor domains

Across studies, the most recurrent predictor groups are:

- **Vitals:** respiratory rate, heart rate, SpO2, blood pressure [13] [16] [17].
- **Respiratory/ventilator variables:** PEEP, FiO2, plateau pressure, tidal volume, ventilator-derived indices [1] [2] [3] [4] [20] [22].
- **Blood gases and acid-base variables:** pH, PaCO2, PaO2/FiO2 [1] [2] [4] [20].
- **Severity and multisystem markers:** SOFA, APACHE, SAPS, vasopressor use, lactate, renal variables, age [1] [17] [23] [25].

This pattern implies that respiratory deterioration is usually embedded in broader physiologic instability rather than isolated respiratory mechanics alone [1] [25].

Engineered temporal features

Several studies used engineered summaries to capture trends:

- Hourly standard deviation plus 12-hour rolling mean for weaning prediction, generating large hourly feature sets before recursive feature elimination [69].

- Minimum, maximum, and mean summaries to capture early dynamic change [70].
- 24-hour aggregation windows in AHRF/ARDS-related prediction [22] [71].
- tsfresh-derived high-frequency features outperforming manual summaries in a ventilator-associated complication study [72].

These methods reflect a shift from static “first-value” prediction toward trajectory-aware modeling [72] [73].

Missing data and class balancing

ICU data pipelines often remove highly missing variables, backward-fill or forward-fill remaining streams, impute residual gaps with prior values or cohort means, and normalize features before modeling [74] [75] [76]. Class imbalance is commonly addressed with SMOTE or other balanced sampling approaches [29] [48] [77] [78].

These steps are practical necessities, but incomplete reporting means replication and bias assessment remain difficult [49] [50].

Predictive Performance and Validation

Internal validation

Internal performance is often good for related ventilated-patient outcomes:

- Random-split ventilated ICU cohorts reported AUCs from **0.806 to 0.821** across common models, with good calibration curves for most models except one neural network [28].
- A difficult-weaning model achieved **AUC 0.8722**, **sensitivity 0.7559**, and **specificity 0.875** [27].
- A deep ventilated-patient model achieved **AUROC 0.879** and **Brier score 0.0974** [29].
- Some broader respiratory or mortality tasks reached AUROCs into the **0.84-0.94** range [1] [17] [19] [20] [23] [79].

These numbers support real signal in routine ICU data, but they should be interpreted as mostly **derivation-cohort performance** [26].

External and temporal validation

External validation is uncommon. A review found only **14.7%** of ICU deterioration models were externally validated, with mean AUROC dropping by **0.037** in external data [26] [37]. In ventilated ICU mortality modeling, a MIMIC-IV logistic model had acceptable external discrimination in eICU but poor calibration, with a reported calibration slope of **0.17**, while exploratory XGBoost improved external AUROC/calibration and local recalibration was recommended [30].

Other multicenter ICU studies show a similar pattern: internal validation overestimates external performance, and recalibration often improves calibration more than discrimination [26] [56] [80] [81].

Calibration and clinical utility

Calibration reporting exists but is inconsistent. Examples include Hosmer-Lemeshow testing, Brier scores, calibration curves, and post-hoc Platt scaling or isotonic regression in selected ICU studies [29] [82] [83]. Still, the evidence that respiratory deterioration models remain well calibrated after deployment is limited [84] [85].

Decision-curve analysis appears in some external validations and sometimes supports net clinical benefit, but the provided evidence also notes that no model consistently beats simpler benchmarks across all settings [51] [86] [87].

Lead time

For the exact question of “how far in advance,” the literature is thinner than it appears. Useful lead-time framing exists for prolonged ventilation and general deterioration, but not many explicit respiratory rescue horizons in ventilated adults. Examples include prediction at intubation, within the first 24-48 hours, or using 72-hour reassessment for prolonged ventilation tasks [11] [60] [88]. Broader ICU deterioration models use hourly rolling labels, including 4-hour ventilator onset and 24-hour mortality windows [55]. But explicit 6-, 12-, 24-, or 48-hour prediction of re-intubation or ventilator escalation remains largely unfilled [12].

Confidence level: High - The evidence consistently supports promising internal discrimination but weaker transportability and sparse true lead-time evidence for the exact endpoint [26] [30].

Clinical Utility, Interpretability, and Workflow Integration

Interpretability

Interpretability tools such as SHAP, LIME, DALEX, and feature importance have identified clinically sensible predictors: respiratory dysfunction, age, sodium and creatinine trajectories, vasopressor adjustments, respiratory rate variability, tidal volume extremes, BNP, lactate, FiO2, and severity scores [1] [24] [25] [89] [90]. In prolonged ventilation models, day-2 PEEP and SOFA repeatedly appeared important [92].

These findings support biological plausibility and may help clinician trust, but they do not by themselves prove bedside usefulness [93] [94].

Deployment and alerting

Real-world deployment evidence is limited. A 2024 real-time system for mechanical power prediction reported workflow-integrated visual alerts intended to support earlier intervention [42]. Another extubation-oriented model was described as producing predictions every 3 minutes using only ventilator-derived inputs [95].

However, the provided studies do **not** report detailed alert burden, false-positive rates, clinician response, or acceptance metrics for these deployments [42] [95]. Broader early-warning literature shows that alerting systems can increase escalation activity without necessarily improving hard outcomes, and workload depends heavily on chosen thresholds [43] [44] [96].

Practical implication

A model with AUROC 0.85 may still be operationally poor if PPV is low and alerts are frequent [40] [43]. That is especially relevant in ICUs already saturated with alarms.

Confidence level: Medium-High - Interpretability evidence is good; operational deployment evidence remains sparse [1] [42] [44].

Limitations, Bias, and Generalizability

The literature repeatedly identifies systematic risks:

- **Retrospective bias and confounding** from local practice and documentation patterns [49] [97] [98].
- **Class imbalance**, which can suppress sensitivity for rare deterioration events [48] [77] [99].
- **Missing data**, often requiring exclusion or imputation [100] [101] [102].
- **Label leakage and overfitting**, especially with repeated measurements and overlapping windows [49] [103].
- **Dataset shift**, including differences across hospitals, ventilator brands, case mix, and time [104] [105] [106].

A systematic review of mechanical ventilation AI studies reported high risk of bias in **89%** of studies, with sparse reporting of calibration, ethnicity, and code/data sharing [50]. Proposed remedies include stricter temporal/geographic splits, multicenter prospective validation, recalibration, transfer learning, federated learning, and cross-database training [19] [107] [108] [109].

Confidence level: High - Methodological quality issues are repeatedly documented and appear structural rather than incidental [50] [110].

Conclusion

Machine learning can predict clinically important adverse trajectories in mechanically ventilated ICU patients using routine vital signs and clinical data, especially when models combine respiratory physiology with broader severity and organ-failure information [1] [17] [25]. The most robust current methods are structured-EHR models built with tree-based algorithms such as XGBoost and random forest, which often achieve internal AUROCs around **0.80-0.90** for related outcomes including mortality, difficult weaning, and prolonged ventilation [6] [17] [18] [27] [28].

But the evidence base does **not** yet strongly support a mature, standardized solution for predicting **imminent respiratory deterioration during ongoing mechanical ventilation** on a 6-48 hour horizon [10] [12]. The biggest barriers are inconsistent outcome definitions, sparse direct-event studies, limited external

and prospective validation, calibration drift under transport, and minimal data on alert burden or real workflow benefit [30] [42] [45] [50].

Taken together, these findings suggest a clear near-term strategy: use ML now for **trajectory-level risk stratification and decision support**, but require standardized labels, multicenter calibration-aware validation, and operational deployment studies before relying on these systems for high-stakes bedside respiratory rescue alerts [17] [19] [26] [107].

Overall conclusion confidence: High - The main synthesis is strongly supported across multiple evidence streams, while the principal uncertainty is the scarcity of exact endpoint-matched studies rather than contradictory results [1] [17] [26] [50].

References

1. Mendiratta, S., Mukkelli, V.G., Baidya Kayal, E.B., (...), Mehndiratta, A. (2025) ML-Driven Prediction of Invasive Mechanical Ventilation in Sepsis: Insights from a tertiary care hospital in India. *Proceedings of the 2025 International Conference on Artificial Intelligence and Emerging Technology, Global AI Summit 2025*. https://doi.org/10.1109/AISummit66170.2025.11411108
2. Vali, M., Paydar, S., Seif, M., (...), Ghaem Maralani, H. (2023) Prediction prolonged mechanical ventilation in trauma patients of the intensive care unit according to initial medical factors: a machine learning approach. *Scientific Reports*. https://doi.org/10.1038/s41598-023-33159-2
3. Villar, J., González-Martín, J.M., Férnandez, C., (...), Szakmany, T. (2025) Predicting Outcome and Duration of Mechanical Ventilation in Acute Hypoxemic Respiratory Failure: The PREMIER Study. *Journal of Clinical Medicine*. https://doi.org/10.3390/jcm14227903
4. Villar, J., González-Martín, J.M., Hernández-González, J., (...), Kacmarek, R.M. (2023) Predicting ICU Mortality in Acute Respiratory Distress Syndrome Patients Using Machine Learning: The Predicting Outcome and STratifiCation of severity in ARDS (POSTCARDS) Study. *Critical Care Medicine*. https://doi.org/10.1097/CCM.0000000000006030
5. Frondelius, T., Atkova, I., Miettunen, J., (...), Jansson, M. (2024) Early prediction of ventilator-associated pneumonia with machine learning models: A systematic review and meta-analysis of prediction model performance☆. *European Journal of Internal Medicine*. https://doi.org/10.1016/j.ejim.2023.11.009
6. Kim, J.H., Kwon, Y.S., Baek, M.S. (2021) Machine learning models to predict 30-day mortality in mechanically ventilated patients. *Journal of Clinical Medicine*. https://doi.org/10.3390/jcm10102172
7. Li, L., Tu, B., Xiong, Y., (...), Yao, Y. (2024) Machine Learning-Based Model for Predicting Prolonged Mechanical Ventilation in Patients with Congestive Heart Failure. *Cardiovascular Drugs and Therapy*. https://doi.org/10.1007/s10557-022-07399-9
8. Kim, J., Kim, Y.K., Kim, H., (...), Kim, H.-J. (2023) Machine Learning Algorithms Predict Successful Weaning From Mechanical Ventilation Before Intubation: Retrospective Analysis From the Medical Information Mart for Intensive Care IV Database. *JMIR Formative Research*. https://doi.org/10.2196/44763
9. Pearce, A.K., Nemati, S., Goligher, E.C., (...), Rogers, A. (2025) Can we predict the future of respiratory failure prediction?. *Critical Care*. https://doi.org/10.1186/s13054-025-05484-7
10. Stivi, T., Padawer, D., Dirini, N., (...), Ledot, S. (2024) Using Artificial Intelligence to Predict Mechanical Ventilation Weaning Success in Patients with Respiratory Failure, Including Those with Acute Respiratory Distress Syndrome. *Journal of Clinical Medicine*. https://doi.org/10.3390/jcm13051505
11. Villar, J., González-Martín, J.M., Férnandez, C., (...), Steyerberg, E.W. (2024) Predicting the Length of Mechanical Ventilation in Acute Respiratory Disease Syndrome Using Machine Learning: The PIONEER Study. *Journal of Clinical Medicine*. https://doi.org/10.3390/jcm13061811
12. Thiele, D., Rodseth, R., Friedland, R., (...), Nee, J. (2025) Machine Learning Models for the Early Real-Time Prediction of Deterioration in Intensive Care Units—A Novel Approach to the Early Identification of High-Risk Patients. *Journal of Clinical Medicine*. https://doi.org/10.3390/jcm14020350

13. Bassin, L., Raubenheimer, J., Bell, D. (2023) The implementation of a real time early warning system using machine learning in an Australian hospital to improve patient outcomes. *Resuscitation*. https://doi.org/10.1016/j.resuscitation.2023.109821
14. Kia, A., Timsina, P., Joshi, H.N., (...), Levin, M.A. (2020) MEWS++: Enhancing the prediction of clinical deterioration in admitted patients through a machine learning model. *Journal of Clinical Medicine*. https://doi.org/10.3390/jcm9020343
15. Salehinejad, H., Meehan, A.M., Rahman, P.A., (...), Caraballo, P.J. (2023) Novel machine learning model to improve performance of an early warning system in hospitalized patients: a retrospective multisite cross-validation study. *eClinicalMedicine*. https://doi.org/10.1016/j.eclinm.2023.102312
16. Akel, M.A., Carey, K.A., Winslow, C.J., (...), Edelson, D.P. (2021) Less is more: Detecting clinical deterioration in the hospital with machine learning using only age, heart rate, and respiratory rate. *Resuscitation*. https://doi.org/10.1016/j.resuscitation.2021.08.024
17. Zhu, Y., Zhang, J., Wang, G., (...), Yu, Q. (2021) Machine Learning Prediction Models for Mechanically Ventilated Patients: Analyses of the MIMIC-III Database. *Frontiers in Medicine*. https://doi.org/10.3389/fmed.2021.662340
18. Xu, H., Ma, Y., Zhuang, Y., (...), Zhou, X. (2024) Machine learning-based risk prediction model construction of difficult weaning in ICU patients with mechanical ventilation. *Scientific Reports*. https://doi.org/10.1038/s41598-024-71548-3
19. Zhang, Y., Chen, X., Tian, X., Zhao, H. (2026) A multicenter machine learning model for predicting ICU mortality in mechanically ventilated patients: development and external validation. *BMC medical informatics and decision making*. https://doi.org/10.1186/s12911-026-03590-2
20. Mendiratta, S., Mukkelli, V.G., Baidya Kayal, E., (...), Mehndiratta, A. (2026) Early prognosis prediction in mechanically ventilated patients using machine learning for tertiary care hospital settings. *Journal of Clinical Monitoring and Computing*. https://doi.org/10.1007/s10877-025-01387-8
21. Abujaber, A., Fadlalla, A., Gammoh, D., (...), El-Menyar, A. (2020) Prediction of in-hospital mortality in patients on mechanical ventilation post traumatic brain injury: machine learning approach. *BMC Medical Informatics and Decision Making*. https://doi.org/10.1186/s12911-020-01363-z
22. Villar, Jesús, González-Martín, Jesús M., Fernández, Cristina, (...), Szakmany, Tamas (2025) Early Prediction of ICU Mortality in Patients with Acute Hypoxemic Respiratory Failure Using Machine Learning: The MEMORIAL Study. *Journal of Clinical Medicine*. https://doi.org/10.3390/jcm14051711
23. Jiang, X., Li, J., Ju, J., (...), Yang, S. (2026) Construction and validation of a machine learning-based risk prediction model for invasive mechanical ventilation in AECOPD patients complicated with respiratory failure. *International Journal of Medical Informatics*. https://doi.org/10.1016/j.ijmedinf.2025.106244
24. Xinli, Ma, Jie, Zhao, Ming, Yan, (...), Lu, Ding (2025) Transformer-based multimodal precision intervention model for enhancing diaphragm function in elderly patients. *Frontiers in Computational Neuroscience*. https://doi.org/10.3389/fncom.2025.1615576
25. Neyra, J.A., Ortiz-Soriano, V., Liu, L.J., (...), Chen, J. (2023) Prediction of Mortality and Major Adverse Kidney Events in Critically Ill Patients With Acute Kidney Injury. *American Journal of Kidney Diseases*. https://doi.org/10.1053/j.ajkd.2022.06.004
26. Rockenschaub, P., Akay, E.M., Carlisle, B.G., (...), Madai, V.I. (2025) External validation of AI-based scoring systems in the ICU: a systematic review and meta-analysis. *BMC Medical Informatics and Decision Making*. https://doi.org/10.1186/s12911-024-02830-7
27. Yin, Chengfen, Xu, Lei, Li, Wenxiong, (...), Ma, Yunfeng (2025) Risk prediction model for difficulty in weaning from mechanical ventilation in critically ill patients: results from a multicentre retrospective study. *BMJ Open*. https://doi.org/10.1136/bmjopen-2024-097419
28. Zhu, Yibing, Zhang, Jin, Wang, Guowei, (...), Yu, Qian (2021) Machine Learning Prediction Models for Mechanically Ventilated Patients: Analyses of the MIMIC-III Database. *Frontiers in Medicine*. https://doi.org/10.3389/fmed.2021.662340
29. Ashrafi, N., Liu, Y., Xu, X., (...), Pishgar, M. (2024) Deep learning model utilization for mortality prediction in mechanically ventilated ICU patients. *Informatics in Medicine Unlocked*. https://doi.org/10.1016/j.imu.2024.101562

30. Zhu, J., Xie, Z., Liu, J., Li, C. (2026) Development and external validation of an ICU mortality prediction model using routinely collected clinical data. *Medicine (United States)*. https://doi.org/10.1097/MD.0000000000049009

31. Bashiri, F.S., Carey, K.A., Martin, J., (...), Churpek, M.M. (2024) Development and external validation of deep learning clinical prediction models using variable-length time series data. *Journal of the American Medical Informatics Association*. https://doi.org/10.1093/jamia/ocae088

32. Peng, J.-C., Liang, J.-R., Zhu, M.-L., (...), Gao, Y. (2026) Transformer-based deep learning for early detection of intensive care unit-acquired bloodstream infection through multivariate time-series analysis. *Digital Health*. https://doi.org/10.1177/20552076251412651

33. Zhang, Y. (2026) Respiratory Failure Risk Prediction Using Nursing Notes and Vital Signs. *Proceedings of 2026 2nd International Conference on Health Big Data, HBD 2026*. https://doi.org/10.1145/3833450.3833462

34. Wang, Y.-C., Zhang, S.-H., Lv, W.-H., (...), Ju, S. (2023) Added value of chest CT images to a personalized prognostic model in acute respiratory distress syndrome: a retrospective study. *Chinese Journal of Academic Radiology*. https://doi.org/10.1007/s42058-023-00116-x

35. Martin, J., Afshar, M., Afshar, A.S., (...), Churpek, M.M. (2026) Explainable multimodal deep learning models for variable-length sequences in critically ill patients. *Journal of Biomedical Informatics*. https://doi.org/10.1016/j.jbi.2026.105001

36. Mathis, M.R., Engoren, M.C., Williams, A.M., (...), Alam, H.B. (2022) Prediction of Postoperative Deterioration in Cardiac Surgery Patients Using Electronic Health Record and Physiologic Waveform Data. *Anesthesiology*. https://doi.org/10.1097/ALN.0000000000004345

37. Amanguli, M., Yaxiaerjiang, M., Guo, Y.-S., (...), Zhong, M. (2026) Artificial intelligence in mechanical ventilation: a narrative review of clinical applications and research gaps. *Journal of Thoracic Disease*. https://doi.org/10.21037/jtd-2026-1257

38. Viderman, D., Ayazbay, A., Kalzhan, B., (...), Abdildin, Y. (2024) Artificial Intelligence in the Management of Patients with Respiratory Failure Requiring Mechanical Ventilation: A Scoping Review. *Journal of Clinical Medicine*. https://doi.org/10.3390/jcm13247535

39. Ji, Q., Wu, Y., Yang, M., (...), Hu, W. (2026) Integration, challenges, and future of artificial intelligence in critical care medicine: comprehensive applications from predictive models to clinical integration. *Frontiers in Medicine*. https://doi.org/10.3389/fmed.2026.1874474

40. Lam, J.Y., Lu, X., Shashikumar, S.P., (...), Nemati, S. (2024) Development, deployment, and continuous monitoring of a machine learning model to predict respiratory failure in critically ill patients. *JAMIA Open*. https://doi.org/10.1093/jamiaopen/ooae141

41. Theilen, R., Huhle, R., Scharffenberg, M., (...), Wittenstein, J. (2026) Intelligent Lung Support in the Intensive Care Unit (IntelliLung): study protocol for an international observational, prospective, multicentre study. *BMJ open*. https://doi.org/10.1136/bmjopen-2025-114851

42. Ruiz-Botella, M., Manrique, S., Gómez, J., Bodí, M. (2024) Advancing ICU patient care with a Real-Time predictive model for mechanical Power to mitigate VILI. *International Journal of Medical Informatics*. https://doi.org/10.1016/j.ijmedinf.2024.105511

43. Prytherch, D.R., Smith, G.B., Schmidt, P.E., Featherstone, P.I. (2010) ViEWS-Towards a national early warning score for detecting adult inpatient deterioration. *Resuscitation*. https://doi.org/10.1016/j.resuscitation.2010.04.014

44. Gao, H., McDonnell, A., Harrison, D.A., (...), Harvey, S. (2007) Systematic review and evaluation of physiological track and trigger warning systems for identifying at-risk patients on the ward. *Intensive Care Medicine*. https://doi.org/10.1007/s00134-007-0532-3

45. Zeng, Xiang, Chen, Xiao Juan, Lai, Ping, (...), Qi, Xiyu (2025) Risk prediction models for extubation failure in critically ill patients on mechanical ventilation: a systematic review. *Frontiers in Medicine*. https://doi.org/10.3389/fmed.2025.1695394

46. Okabe, Y., Asaga, T., Bekku, S., (...), Shirakami, G. (2018) Lung-thorax compliance measured during a spontaneous breathing trial is a good index of extubation failure in the surgical intensive care unit: A retrospective cohort study. *Journal of Intensive Care*. https://doi.org/10.1186/s40560-018-0313-9

47. Bansal, V., Smischney, N.J., Kashyap, R., (...), Freeman, W.D. (2022) Reintubation Summation Calculation: A Predictive Score for Extubation Failure in Critically Ill Patients. *Frontiers in Medicine*. https://doi.org/10.3389/fmed.2021.789440

48. Chen, Wenqian, Lee, Benjamin, Zang, Zexi, (...), Ren, Yanli (2026) Data subdivision approach enhances machine learning-based mortality prediction in pediatric ICU patients. *PLoS ONE*. https://doi.org/10.1371/journal.pone.0349772

49. Briassoulis, George, Briassouli, Efrossini (2025) AI-Enabled Precision Nutrition in the ICU: A Narrative Review and Implementation Roadmap. *Nutrients*. https://doi.org/10.3390/nu18010110

50. Gallifant, J., Zhang, J., Arias López, M., (...), Formenti, F. (2022) Artificial intelligence for mechanical ventilation: systematic review of design, reporting standards, and bias. *British Journal of Anaesthesia*. https://doi.org/10.1016/j.bja.2021.09.025

51. Lu, Y., Chen, H., Li, M., (...), Lv, H. (2026) Machine learning-based prediction of early invasive mechanical ventilation in ICU patients with pneumonia: Development and external validation. *International Journal of Medical Informatics*. https://doi.org/10.1016/j.ijmedinf.2026.106541

52. Zeng, Xiang, Chen, Xiao Juan, Lai, Ping, (...), Qi, Xiyu (2025) Risk prediction models for extubation failure in critically ill patients on mechanical ventilation: a systematic review. *Frontiers in Medicine*. https://doi.org/10.3389/fmed.2025.1695394

53. Huang, C.-T., Yu, C.-J. (2013) Conventional weaning parameters do not predict extubation outcome in intubated subjects requiring prolonged mechanical ventilation. *Respiratory Care*. https://doi.org/10.4187/respcare.01773

54. Funk, G.-C., Anders, S., Breyer, M.-K., (...), Hartl, S. (2010) Incidence and outcome of weaning from mechanical ventilation according to new categories. *European Respiratory Journal*. https://doi.org/10.1183/09031936.00056909

55. Martin, J., Afshar, M., Afshar, A.S., (...), Churpek, M.M. (2026) Explainable multimodal deep learning models for variable-length sequences in critically ill patients. *Journal of Biomedical Informatics*. https://doi.org/10.1016/j.jbi.2026.105001

56. Cox, E.G.M., Wiersema, R., Eck, R.J., (...), Keus, F. (2023) External Validation of Mortality Prediction Models for Critical Illness Reveals Preserved Discrimination but Poor Calibration. *Critical Care Medicine*. https://doi.org/10.1097/CCM.0000000000005712

57. Churpek, M.M., Yuen, T.C., Winslow, C., (...), Edelson, D.P. (2016) Multicenter Comparison of Machine Learning Methods and Conventional Regression for Predicting Clinical Deterioration on the Wards. *Critical Care Medicine*. https://doi.org/10.1097/CCM.0000000000001571

58. Epstein, S. (2002) Decision to extubate. *Intensive Care Medicine*. https://doi.org/10.1007/s00134-002-1268-8

59. Knaus, W.A., Draper, E.A., Wagner, D.P., Zimmerman, J.E. (1985) APACHE II: A severity of disease classification system. *Critical Care Medicine*. https://doi.org/10.1097/00003246-198510000-00009

60. Clark, P.A., Inocencio, R.C., Lettieri, C.J. (2018) I-TRACH: Validating A Tool for Predicting Prolonged Mechanical Ventilation. *Journal of Intensive Care Medicine*. https://doi.org/10.1177/0885066616679974

61. Thambiraj, G., Bazoukis, G., Ghabousian, A., (...), Armoundas, A.A. (2025) Use of artificial intelligence in predicting in-hospital cardiac and respiratory arrest in an acute care environment—implications for clinical practice. *Frontiers in Medical Technology*. https://doi.org/10.3389/fmedt.2025.1681059

62. Aarab, Y., Pensier, J., Garnier, F., (...), Jaber, S. (2025) Incidence and outcomes of extubation failure in mechanically ventilated patients with cirrhosis: a post-hoc analysis of a prospective multicenter study. *Annals of Intensive Care*. https://doi.org/10.1186/s13613-025-01576-3

63. Choe, J., Nam, J., Lee, H.O., (...), Jeon, K. (2026) Negative inspiratory force predicts extubation failure after planned extubation in a medical ICU: a prospective cohort study. *Intensive and Critical Care Nursing*. https://doi.org/10.1016/j.iccn.2026.104531

64. Su, C.-L., Chiang, L.-L., Yang, S.-H., (...), Wu, C.-P. (2012) Preventive use of noninvasive ventilation after extubation: A prospective, multicenter randomized controlled trial. *Respiratory Care*. https://doi.org/10.4187/respcare.01141

65. Baptistella, Antuani Rafael, Mantelli, Laura Maito, Matte, Leandra, (...), Nunes Filho, João Rogério (2021) Prediction of extubation outcome in mechanically ventilated patients: Development and validation of the Extubation Predictive Score (ExPreS). *PLoS ONE*. https://doi.org/10.1371/journal.pone.0248868

66. https://www.sciencedirect.com/science/article/pii/S1364853521002337

67. Hssain, A.A., Pandian, V., Vahedian-Azimi, A., (...), Labib Shehatta, A.L. (2026) Long-term outcomes and prognostic factors of patients undergoing tracheostomy for prolonged mechanical ventilation: A retrospective study. *Intensive and Critical Care Nursing*. https://doi.org/10.1016/j.iccn.2026.104541

68. Koneru, A., Narne, N.K., Amirineni, S. (2025) Deep Learning Architectures for Predictive Healthcare Analytics: From Diagnosis to Prognosis. *2025 IEEE International Conference on Communication Networks and Computing, CNC 2025*. https://doi.org/10.1109/CNC68716.2025.11484661

69. Zappalà, S., Scaravilli, V., Rovati, L., (...), Grasselli, G. (2025) Development and validation of a machine learning model for real-time prediction of invasive mechanical ventilation weaning readiness. *Journal of Critical Care*. https://doi.org/10.1016/j.jcrc.2025.155105

70. Wang, Yeao, Rong, Jianke, Wei, Zhili, (...), Deng, YunDan (2025) Predicting ICU mortality in heart failure patients based on blood tests and vital signs. *Frontiers in Cardiovascular Medicine*. https://doi.org/10.3389/fcvm.2025.1590367

71. https://www.sciencedirect.com/science/article/pii/S1433305824012061

72. Tan, Y., Young, M., Girish, A., (...), Bergmann, J. (2023) Predicting respiratory decompensation in mechanically ventilated adult ICU patients. *Frontiers in Physiology*. https://doi.org/10.3389/fphys.2023.1125991

73. Jaadi, S.A., Al Wahaibi, L., Al-Hinai, M., (...), al-Alawi, A.M. (2026) Advanced Machine Learning Techniques for Predicting Inpatient Deterioration in General Medicine. *Computers*. https://doi.org/10.3390/computers15020123

74. Elbehery, M.M., Mohamed, M.S., Elshafey, M.A. (2026) Explainable ICD coding via MediThoth: A multimodal clinical decision support system. *Array*. https://doi.org/10.1016/j.array.2026.100875

75. Dao, B.T., Ma, V.D., Vu, T.A., (...), Truong, T.H. (2026) HyperShield: A personalized hypernetwork-based defense against poisoning attacks in federated learning for healthcare. *Expert Systems with Applications*. https://doi.org/10.1016/j.eswa.2025.130237

76. Zou, M., An, Y., Kuang, H., Wang, J. (2023) LGTRL-DE: Local and Global Temporal Representation Learning with Demographic Embedding for in-hospital mortality prediction. *Journal of Biomedical Informatics*. https://doi.org/10.1016/j.jbi.2023.104408

77. Gülbay, Mutlu, Başţuğ, Aliye, Özkan, Erdem, (...), Bodur, Hürrem (2022) Evaluation of the models generated from clinical features and deep learning-based segmentations: Can thoracic CT on admission help us to predict hospitalized COVID-19 patients who will require intensive care?. *BMC Medical Imaging*. https://doi.org/10.1186/s12880-022-00833-2

78. Nguyen, Q.T., Tran, M.P., Prabhakaran, V., (...), Nguyen, G.H. (2024) Compact machine learning model for the accurate prediction of first 24-hour survival of mechanically ventilated patients. *Frontiers in Medicine*. https://doi.org/10.3389/fmed.2024.1398565

79. Zhou, M. (2024) Early Prediction of Invasive Mechanical Ventilation Needs in Patients Admitted to Intensive Care Units: A Novel Ensemble Learning Approach. *6th International Conference on Intelligent Computing in Data Sciences, ICDS 2024*. https://doi.org/10.1109/ICDS62089.2024.10756474

80. Allyn, J., Oliver, M., Cerveau, R., (...), Ferdynus, C. (2026) Multi-source data integration through pooling and transfer learning improves generalizability and specialization of deep learning models for ICU mortality and length of stay prediction: a four-database external validation study. *Critical Care*. https://doi.org/10.1186/s13054-026-06034-5

81. van den Boogaard, M., Schoonhoven, L., Maseda, E., (...), Pickkers, P. (2014) Recalibration of the delirium prediction model for ICU patients (PRE-DELIRIC): A multinational observational study. *Intensive Care Medicine*. https://doi.org/10.1007/s00134-013-3202-7

82. Hwang, Hyeontaek, Choi, Sun Mi, Lee, Jinwoo, (...), Lee, Sang-Min (2000) *Respiratory Research*. https://doi.org/10.1186/s12931-020-01357-5

83. Piasecki, Adam, Adamczyk, Kamil, Śniegowski, Marcin, (...), Elbers, Paul W. G. (2025) Early prognostication for ICU patients with combined respiratory and circulatory failure: an interpretable machine learning approach. *Scientific Reports*. https://doi.org/10.1038/s41598-026-59064-y

84. Bansal, Vikas, Smischney, Nathan J., Kashyap, Rahul, (...), Freeman, William David (2021) Reintubation Summation Calculation: A Predictive Score for Extubation Failure in Critically Ill Patients. *Frontiers in Medicine*. https://doi.org/10.3389/fmed.2021.789440

85. Zhu, Yibing, Zhang, Jin, Wang, Guowei, (...), Yu, Qian (2021) Machine Learning Prediction Models for Mechanically Ventilated Patients: Analyses of the MIMIC-III Database. *Frontiers in Medicine*. https://doi.org/10.3389/fmed.2021.662340

86. Rothman, M.J., Rothman, S.I., Beals, J. (2013) Development and validation of a continuous measure of patient condition using the Electronic Medical Record. *Journal of Biomedical Informatics*. https://doi.org/10.1016/j.jbi.2013.06.011

87. Ishii, E., Nawa, N., Hashimoto, S., (...), Fujiwara, T. (2023) Development, validation, and feature extraction of a deep learning model predicting in-hospital mortality using Japan's largest national ICU database: a validation framework for transparent clinical Artificial Intelligence (cAI) development. *Anaesthesia Critical Care and Pain Medicine*. https://doi.org/10.1016/j.accpm.2022.101167

88. Figueroa-Casas, J.B., Dwivedi, A.K., Connery, S.M., (...), Galvis, J. (2015) Predictive models of prolonged mechanical ventilation yield moderate accuracy. *Journal of Critical Care*. https://doi.org/10.1016/j.jcrc.2015.01.020

89. Cifci, Mehmet Akif, Öney, Batuhan, Yildirim, Fazli, (...), Zontul, Metin (2025) Interpretable Adaptive Graph Fusion Network for Mortality and Complication Prediction in ICUs. *Diagnostics*. https://doi.org/10.3390/diagnostics15222825

90. https://www.sciencedirect.com/science/article/pii/S2297055X26016495

91. Figueroa-Casas, J.B., Dwivedi, A.K., Connery, S.M., (...), Galvis, J. (2015) Predictive models of prolonged mechanical ventilation yield moderate accuracy. *Journal of Critical Care*. https://doi.org/10.1016/j.jcrc.2015.01.020

92. Zhao, Y., Chen, S., Pang, C., (...), Lin, S. (2026) Development and external validation of a machine-learning model for classifying documented pressure injury status during ICU stays among patients receiving invasive mechanical ventilation: A MIMIC-IV and eICU study. *Journal of Tissue Viability*. https://doi.org/10.1016/j.jtv.2026.101058

93. https://www.sciencedirect.com/science/article/pii/S1471247424011080

94. Huang, Kuo-Yang, Hsu, Ying-Lin, Chen, Huang-Chi, (...), Hou, Ming-Hon (2023) Developing a machine-learning model for real-time prediction of successful extubation in mechanically ventilated patients using time-series ventilator-derived parameters. *Frontiers in Medicine*. https://doi.org/10.3389/fmed.2023.1167445

95. Hillman, K. (2005) Introduction of the medical emergency team (MET) system: A cluster-randomised controlled trial. *Lancet*. https://doi.org/10.1016/S0140-6736(05)66733-5

96. Taparugssanagorn, Attaphongse, Särestöniemi, Mariella, Hämäläinen, Matti, Iinatti, Jari (2026) Early Sepsis Detection Using Heterogeneous Structured ICU Data with Explainable Deep Learning. *Sensors (Basel, Switzerland)*. https://doi.org/10.3390/s26123648

97. Hadweh, Pierre, Niset, Alexandre, Salvagno, Michele, (...), Barrit, Sami (2025) Machine Learning and Artificial Intelligence in Intensive Care Medicine: Critical Recalibrations from Rule-Based Systems to Frontier Models. *Journal of Clinical Medicine*. https://doi.org/10.3390/jcm14124026

98. https://www.sciencedirect.com/science/article/pii/S147269472500336X

99. Bai, Yu, Xia, Jingen, Huang, Xu, (...), Zhan, Qingyuan (2022) Using machine learning for the early prediction of sepsis-associated ARDS in the ICU and identification of clinical phenotypes with

differential responses to treatment. *Frontiers in Physiology*. https://doi.org/10.3389/fphys.2022.1050849

100. https://www.sciencedirect.com/science/article/pii/S1471233425011331

101. Yang, Zhenyu, Cui, Xiaoju, Song, Zhe (2023) Predicting sepsis onset in ICU using machine learning models: a systematic review and meta-analysis. *BMC Infectious Diseases*. https://doi.org/10.1186/s12879-023-08614-0

102. Lim, L., Kim, M., Cho, K., (...), Lee, H.-C. (2025) Multicenter validation of a machine learning model to predict intensive care unit readmission within 48 hours after discharge. *eClinicalMedicine*. https://doi.org/10.1016/j.eclinm.2025.103112

103. Piasecki, Adam, Adamczyk, Kamil, Śniegowski, Marcin, (...), Elbers, Paul W. G. (2025) Early prognostication for ICU patients with combined respiratory and circulatory failure: an interpretable machine learning approach. *Scientific Reports*. https://doi.org/10.1038/s41598-026-59064-y

104. Bakos, Péter, Szabó, Bence, Laczkó, Dávid, (...), Molnár, Zsolt (2026) Machine learning models predicting extubation success in mechanically ventilated patients: a systematic review and meta-analysis. *Intensive Care Medicine Experimental*. https://doi.org/10.1186/s40635-026-00934-0

105. Patel, Anita K, Trujillo-Rivera, Eduardo, Morizono, Hiroki, Pollack, Murray M. (2022) The criticality Index-mortality: A dynamic machine learning prediction algorithm for mortality prediction in children cared for in an ICU. *Frontiers in Pediatrics*. https://doi.org/10.3389/fped.2022.1023539

106. Sheikhalishahi, S., Kaspar, M., Schwinn, J., (...), Hinske, L.C. (2026) Cross-Silo Federated Learning for Predicting Successful Mechanical Ventilation Weaning Across 5 Intensive Care Unit Databases: Retrospective Database Analysis. *JMIR Medical Informatics*. https://doi.org/10.2196/79713

107. Lin, S., Yang, M., Liu, C., (...), Long, X. (2024) A pretrain-finetune approach for improving model generalizability in outcome prediction of acute respiratory distress syndrome patients. *International Journal of Medical Informatics*. https://doi.org/10.1016/j.ijmedinf.2024.105397

108. Churpek, M.M., Carey, K.A., Snyder, A., (...), Edelson, D.P. (2025) Multicenter Development and Prospective Validation of eCARTv5: A Gradient-Boosted Machine-Learning Early Warning Score. *Critical Care Explorations*. https://doi.org/10.1097/CCE.0000000000001232

109. Wolff, R.F., Moons, K.G.M., Riley, R.D., (...), Mallett, S. (2019) PROBAST: A tool to assess the risk of bias and applicability of prediction model studies. *Annals of Internal Medicine*. https://doi.org/10.7326/M18-1376