

# Some important basis functions

We also solve some cool problems

---

18.303 Linear Partial Differential Equations: Analysis and Numerics

# Laplace equation in spherical coordinates

We solve for  $\Delta f = 0$  in spherical coordinates.

# Laplace equation in spherical coordinates

We solve for  $\Delta f = 0$  in spherical coordinates.

We have the coordinate chart between the Cartesian and spherical coordinates:

$$x = r \sin(\theta) \cos(\varphi)$$

$$y = r \sin(\theta) \sin(\varphi)$$

$$z = r \cos(\theta),$$

where  $r \in \mathbb{R}_+$ ,  $\theta \in [0, \pi]$ , and  $\varphi \in [0, 2\pi)$ .

# Laplace equation in spherical coordinates

We solve for  $\Delta f = 0$  in spherical coordinates.

We have the coordinate chart between the Cartesian and spherical coordinates:

$$x = r \sin(\theta) \cos(\varphi)$$

$$y = r \sin(\theta) \sin(\varphi)$$

$$z = r \cos(\theta),$$

where  $r \in \mathbb{R}_+$ ,  $\theta \in [0, \pi]$ , and  $\varphi \in [0, 2\pi)$ .

It requires a bit of work to calculate the change of coordinates but ultimately we get

$$\Delta f(r, \theta, \varphi) = \left[ \frac{1}{r^2} \partial_r (r^2 \partial_r) + \frac{1}{r^2 \sin(\theta)} \partial_\theta (\sin(\theta) \partial_\theta) + \frac{1}{r^2 \sin^2(\theta)} \partial_\varphi^2 \right] f(r, \theta, \varphi) = 0.$$

# Laplace equation in spherical coordinates

We solve for  $\Delta f = 0$  in spherical coordinates.

We have the coordinate chart between the Cartesian and spherical coordinates:

$$x = r \sin(\theta) \cos(\varphi)$$

$$y = r \sin(\theta) \sin(\varphi)$$

$$z = r \cos(\theta),$$

where  $r \in \mathbb{R}_+$ ,  $\theta \in [0, \pi]$ , and  $\varphi \in [0, 2\pi)$ .

It requires a bit of work to calculate the change of coordinates but ultimately we get

$$\Delta f(r, \theta, \varphi) = \left[ \frac{1}{r^2} \partial_r (r^2 \partial_r) + \frac{1}{r^2 \sin(\theta)} \partial_\theta (\sin(\theta) \partial_\theta) + \frac{1}{r^2 \sin^2(\theta)} \partial_\varphi^2 \right] f(r, \theta, \varphi) = 0.$$

We use separation by variables and write  $f(r, \theta, \varphi) = R(r)\Theta(\theta)\Phi(\varphi)$ .

Multiplying by  $r^2$  and reorganizing gives

$$\frac{\partial_r (r^2 R'(r))}{R(r)} + \frac{\sin^{-1}(\theta) \partial_\theta (\sin(\theta) \Theta'(\theta))}{\Theta(\theta)} + \frac{1}{\sin^2(\theta)} \frac{\Phi''(\varphi)}{\Phi(\varphi)} = 0.$$

Multiplying by  $r^2$  and reorganizing gives

$$\frac{\partial_r (r^2 R'(r))}{R(r)} + \frac{\sin^{-1}(\theta) \partial_\theta (\sin(\theta) \Theta'(\theta))}{\Theta(\theta)} + \frac{1}{\sin^2(\theta)} \frac{\Phi''(\varphi)}{\Phi(\varphi)} = 0.$$

This has to hold for all  $r$ ,  $\theta$ , and  $\varphi$  so some parts need to be constants. Probably the easiest is

$$\Phi_m''(\varphi) = m^2 \Phi_m(\varphi).$$

Multiplying by  $r^2$  and reorganizing gives

$$\frac{\partial_r (r^2 R'(r))}{R(r)} + \frac{\sin^{-1}(\theta) \partial_\theta (\sin(\theta) \Theta'(\theta))}{\Theta(\theta)} + \frac{1}{\sin^2(\theta)} \frac{\Phi''(\varphi)}{\Phi(\varphi)} = 0.$$

This has to hold for all  $r$ ,  $\theta$ , and  $\varphi$  so some parts need to be constants. Probably the easiest is

$$\Phi_m''(\varphi) = m^2 \Phi_m(\varphi).$$

This gives the azimuthal part of the function and  $\Phi$  is periodic. It is solved by

$$\Phi_m(\varphi) = e^{im\varphi},$$

where  $m$  is an integer.



Now we have

$$\begin{aligned}\partial_r (r^2 \partial_r R_\ell(r)) &= \lambda_\ell R_\ell(r), \\ \frac{1}{\sin(\theta)} \partial_\theta (\sin(\theta) \partial_\theta \Theta_\ell^m(\theta)) - \frac{m^2}{\sin^2(\theta)} \Theta_\ell^m &= -\lambda_\ell \Theta_\ell^m(\theta).\end{aligned}$$

Now we have

$$\begin{aligned}\partial_r (r^2 \partial_r R_\ell(r)) &= \lambda_\ell R_\ell(r), \\ \frac{1}{\sin(\theta)} \partial_\theta (\sin(\theta) \partial_\theta \Theta_\ell^m(\theta)) - \frac{m^2}{\sin^2(\theta)} \Theta_\ell^m &= -\lambda_\ell \Theta_\ell^m(\theta).\end{aligned}$$

Since  $\theta \in [0, \pi]$ , we can make a change of variables  $t = \cos(\theta) \in [-1, 1]$  ( $\cos$  is a bijection on this interval). We say  $\Theta_\ell^m(\theta) = P_\ell^m(\cos(\theta)) = P_\ell^m(t)$ .

Now we have

$$\begin{aligned}\partial_r (r^2 \partial_r R_\ell(r)) &= \lambda_\ell R_\ell(r), \\ \frac{1}{\sin(\theta)} \partial_\theta (\sin(\theta) \partial_\theta \Theta_\ell^m(\theta)) - \frac{m^2}{\sin^2(\theta)} \Theta_\ell^m &= -\lambda_\ell \Theta_\ell^m(\theta).\end{aligned}$$

Since  $\theta \in [0, \pi]$ , we can make a change of variables  $t = \cos(\theta) \in [-1, 1]$  ( $\cos$  is a bijection on this interval). We say  $\Theta_\ell^m(\theta) = P_\ell^m(\cos(\theta)) = P_\ell^m(t)$ .

The derivative with respect to  $\theta$  transforms as

$$\partial_\theta \rightarrow \frac{\partial t}{\partial \theta} \partial_t = -\sin(\theta) \partial_t.$$

Now we have

$$\begin{aligned}\partial_r (r^2 \partial_r R_\ell(r)) &= \lambda_\ell R_\ell(r), \\ \frac{1}{\sin(\theta)} \partial_\theta (\sin(\theta) \partial_\theta \Theta_\ell^m(\theta)) - \frac{m^2}{\sin^2(\theta)} \Theta_\ell^m &= -\lambda_\ell \Theta_\ell^m(\theta).\end{aligned}$$

Since  $\theta \in [0, \pi]$ , we can make a change of variables  $t = \cos(\theta) \in [-1, 1]$  ( $\cos$  is a bijection on this interval). We say  $\Theta_\ell^m(\theta) = P_\ell^m(\cos(\theta)) = P_\ell^m(t)$ .

The derivative with respect to  $\theta$  transforms as

$$\partial_\theta \rightarrow \frac{\partial t}{\partial \theta} \partial_t = -\sin(\theta) \partial_t.$$

Now the equation for the latitudinal part becomes

$$\partial_t (\sin^2(\theta) \partial_t P_\ell^m) - \frac{m^2}{\sin^2(\theta)} P_\ell^m = -\lambda_\ell P_\ell^m(t).$$

Writing  $\sin^2(\theta) = 1 - \cos^2(\theta) = 1 - t^2$  gives

$$\partial_t \left( (1 - t^2) \partial_t P_\ell^m \right) - \frac{m^2}{1 - t^2} P_\ell^m = -\lambda_\ell P_\ell^m(t).$$

Writing  $\sin^2(\theta) = 1 - \cos^2(\theta) = 1 - t^2$  gives

$$\partial_t ((1 - t^2) \partial_t P_\ell^m) - \frac{m^2}{1 - t^2} P_\ell^m = -\lambda_\ell P_\ell^m(t).$$

The only way  $P_\ell^m(\pm 1)$  is finite is if  $\lambda_\ell = \ell(\ell + 1)$ , where  $\ell$  is a non-negative integer (this is a long story why this happens). Because the differential operator to the left is negative semi-definite, we have to have  $|m| \leq \ell$ .

Writing  $\sin^2(\theta) = 1 - \cos^2(\theta) = 1 - t^2$  gives

$$\partial_t ((1 - t^2) \partial_t P_\ell^m) - \frac{m^2}{1 - t^2} P_\ell^m = -\lambda_\ell P_\ell^m(t).$$

The only way  $P_\ell^m(\pm 1)$  is finite is if  $\lambda_\ell = \ell(\ell + 1)$ , where  $\ell$  is a non-negative integer (this is a long story why this happens). Because the differential operator to the left is negative semi-definite, we have to have  $|m| \leq \ell$ .

The solution to this DE is given by the **Associated Legendre polynomials**. The usual Legendre polynomials  $P_\ell$  are given for  $m = 0$ . Now  $\Theta_\ell^m(\theta) = P_\ell^m(\cos(\theta))$ .

Writing  $\sin^2(\theta) = 1 - \cos^2(\theta) = 1 - t^2$  gives

$$\partial_t ((1 - t^2) \partial_t P_\ell^m) - \frac{m^2}{1 - t^2} P_\ell^m = -\lambda_\ell P_\ell^m(t).$$

The only way  $P_\ell^m(\pm 1)$  is finite is if  $\lambda_\ell = \ell(\ell + 1)$ , where  $\ell$  is a non-negative integer (this is a long story why this happens). Because the differential operator to the left is negative semi-definite, we have to have  $|m| \leq \ell$ .

The solution to this DE is given by the **Associated Legendre polynomials**. The usual Legendre polynomials  $P_\ell$  are given for  $m = 0$ . Now  $\Theta_\ell^m(\theta) = P_\ell^m(\cos(\theta))$ .

The polynomials  $P_\ell$  are degree  $\ell$  and are uniquely defined by  $P_\ell(1) = 1$  and the orthogonality condition

$$\int_{-1}^1 P_\ell(t) P_{\ell'}(t) dt = 0$$

if  $\ell \neq \ell'$ . If  $\ell = \ell'$ , this integral gives  $2/(2\ell + 1)$ .



The associated Legendre polynomials can be calculated from the Legendre polynomials through

$$P_{\ell}^m(t) = (-1)^m (1 - t^2)^{m/2} \frac{d^m}{dt^m} P_{\ell}(t).$$

The associated Legendre polynomials can be calculated from the Legendre polynomials through

$$P_{\ell}^m(t) = (-1)^m (1 - t^2)^{m/2} \frac{d^m}{dt^m} P_{\ell}(t).$$

The (normal) Legendre polynomials can also be obtained as

$$P_{\ell}(t) = \frac{1}{2^{\ell} \ell!} \frac{d^{\ell}}{dt^{\ell}} (t^2 - 1)^{\ell}.$$

The associated Legendre polynomials can be calculated from the Legendre polynomials through

$$P_{\ell}^m(t) = (-1)^m (1 - t^2)^{m/2} \frac{d^m}{dt^m} P_{\ell}(t).$$

The (normal) Legendre polynomials can also be obtained as

$$P_{\ell}(t) = \frac{1}{2^{\ell} \ell!} \frac{d^{\ell}}{dt^{\ell}} (t^2 - 1)^{\ell}.$$

These polynomials have many nifty properties but we will not spend too much time on that today.

Finally we have the radial part

$$\partial_r(r^2 R'_\ell(r)) = \ell(\ell + 1)R_\ell(r).$$

Finally we have the radial part

$$\partial_r(r^2 R'_\ell(r)) = \ell(\ell + 1)R_\ell(r).$$

This is solved by an ansatz  $R = r^\alpha$  giving

$$\alpha(\alpha + 1) = \ell(\ell + 1),$$

which implies  $\alpha = \ell$  or  $\alpha = -\ell - 1$ . Now

$$R(r) = A_\ell r^{\ell r} + B_\ell r^{-\ell-1}.$$

Finally we have the radial part

$$\partial_r(r^2 R'_\ell(r)) = \ell(\ell + 1)R_\ell(r).$$

This is solved by an ansatz  $R = r^\alpha$  giving

$$\alpha(\alpha + 1) = \ell(\ell + 1),$$

which implies  $\alpha = \ell$  or  $\alpha = -\ell - 1$ . Now

$$R(r) = A_\ell r^{\ell r} + B_\ell r^{-\ell-1}.$$

If we require that  $R$  is bounded at the origin, it implies that  $B_\ell = 0$ . Another usual boundary condition for  $R$  is that it is bounded at infinity. In that case  $A_\ell = 0$ . Alternatively you can add your favorite boundary condition at some fixed  $r_0$  and solve for the coefficients.

# Spherical harmonics

The functions

$$Y_{\ell}^m(\theta, \varphi) = \sqrt{\frac{(2\ell+1)(\ell-m)!}{4\pi(\ell+m)!}} P_{\ell}^m(\cos\theta) e^{im\varphi}$$

are the spherical harmonics that satisfy

$$r^2 \Delta Y_{\ell}^m = -\ell(\ell+1) Y_{\ell}^m$$

i.e. they are the eigenfunctions of the Laplacian on the surface of a sphere with a fixed radius  $r$ .

# Spherical harmonics

The functions

$$Y_{\ell}^m(\theta, \varphi) = \sqrt{\frac{(2\ell+1)(\ell-m)!}{4\pi(\ell+m)!}} P_{\ell}^m(\cos\theta) e^{im\varphi}$$

are the spherical harmonics that satisfy

$$r^2 \Delta Y_{\ell}^m = -\ell(\ell+1) Y_{\ell}^m$$

i.e. they are the eigenfunctions of the Laplacian on the surface of a sphere with a fixed radius  $r$ .

They are also normalized satisfying

$$\int_{\partial\Omega} Y_{\ell}^m Y_{\ell'}^{m'} dS = \int_0^{2\pi} \int_0^{\pi} \sin(\theta) Y_{\ell}^m Y_{\ell'}^{m'} d\theta d\varphi = \delta_{\ell}^{\ell'} \delta_m^{m'}.$$



# Spherical harmonics

The functions

$$Y_{\ell}^m(\theta, \varphi) = \sqrt{\frac{(2\ell+1)}{4\pi} \frac{(\ell-m)!}{(\ell+m)!}} P_{\ell}^m(\cos \theta) e^{im\varphi}$$

are the spherical harmonics that satisfy

$$r^2 \Delta Y_{\ell}^m = -\ell(\ell+1) Y_{\ell}^m$$

i.e. they are the eigenfunctions of the Laplacian on the surface of a sphere with a fixed radius  $r$ .

They are also normalized satisfying

$$\int_{\partial\Omega} Y_{\ell}^m Y_{\ell'}^{m'} dS = \int_0^{2\pi} \int_0^{\pi} \sin(\theta) Y_{\ell}^m Y_{\ell'}^{m'} d\theta d\varphi = \delta_{\ell}^{\ell'} \delta_m^{m'}.$$

They define a perfectly good basis for dealing with differential equations on a sphere.