

A bit of vector calculus

Some integration rules

18.303 Linear Partial Differential Equations: Analysis and Numerics

Divergence and gradient theorems

For $g : \mathbb{R}^N \rightarrow \mathbb{R}$ we have

$$\int_{\gamma} \nabla g(\mathbf{x}) \cdot d\mathbf{x} = g(\mathbf{x}_e) - g(\mathbf{x}_s).$$

Here γ is a differentiable path embedded in \mathbb{R}^N and $g(\mathbf{x}_s)$ and $g(\mathbf{x}_e)$ are the start and endpoints of the path. This is called the **gradient theorem** or the fundamental theorem of calculus for line integrals.

Divergence and gradient theorems

For $g : \mathbb{R}^N \rightarrow \mathbb{R}$ we have

$$\int_{\gamma} \nabla g(\mathbf{x}) \cdot d\mathbf{x} = g(\mathbf{x}_e) - g(\mathbf{x}_s).$$

Here γ is a differentiable path embedded in \mathbb{R}^N and $g(\mathbf{x}_s)$ and $g(\mathbf{x}_e)$ are the start and endpoints of the path. This is called the **gradient theorem** or the fundamental theorem of calculus for line integrals.

For a compact $\Omega \subset \mathbb{R}^N$ and a function $\mathbf{f} : \mathbb{R}^N \rightarrow \mathbb{R}^N$ we have

$$\int_{\Omega} \nabla \cdot \mathbf{f}(\mathbf{x}) dV = \int_{\partial\Omega} \mathbf{f}(\mathbf{x}) \cdot d\mathbf{S} = \int_{\partial\Omega} \mathbf{f}(\mathbf{x}) \cdot \hat{\mathbf{n}}(\mathbf{x}) dS,$$

where $\hat{\mathbf{n}}$ is the unit normal of a given point on the boundary of the region Ω . We assume here that the boundary is piecewise smooth. This is called the **divergence theorem** or Gauss's theorem.

Let $\mathbf{A} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be a sufficiently smooth vector field. Let Σ be a *simply connected* smooth two-dimensional subset of \mathbb{R}^3 with a piecewise smooth boundary.

Let $\mathbf{A} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be a sufficiently smooth vector field. Let Σ be a *simply connected* smooth two-dimensional subset of \mathbb{R}^3 with a piecewise smooth boundary.

We have

$$\int_{\Sigma} \nabla \times \mathbf{A} \cdot d\mathbf{S} = \oint_{\partial\Sigma} \mathbf{A} \cdot d\mathbf{x}.$$

Here the integral is contracted against the the tangent .

Green's first identity

Let $\varphi, \psi : \mathbb{R}^N \rightarrow \mathbb{R}$ be differentiable functions and $\Omega \subset \mathbb{R}^N$ a region with a piecewise differentiable boundary.

Green's first identity

Let $\varphi, \psi : \mathbb{R}^N \rightarrow \mathbb{R}$ be differentiable functions and $\Omega \subset \mathbb{R}^N$ a region with a piecewise differentiable boundary.

$$\int_{\Omega} \psi \Delta \varphi + \nabla \psi \cdot \nabla \varphi dV = \int_{\partial \Omega} \psi \nabla \varphi \cdot d\mathbf{S}.$$

Green's first identity

Let $\varphi, \psi : \mathbb{R}^N \rightarrow \mathbb{R}$ be differentiable functions and $\Omega \subset \mathbb{R}^N$ a region with a piecewise differentiable boundary.

$$\int_{\Omega} \psi \Delta \varphi + \nabla \psi \cdot \nabla \varphi dV = \int_{\partial \Omega} \psi \nabla \varphi \cdot d\mathbf{S}.$$

This can be also written as

$$\int_{\Omega} \nabla \psi \cdot \nabla \varphi dV = \int_{\partial \Omega} \psi \nabla \varphi \cdot d\mathbf{S} - \int_{\Omega} \psi \Delta \varphi dV,$$

so it becomes a rule for integration by parts for vector calculus.

Derive Green's first identity using the divergence theorem. Hint, choose $\mathbf{f} = \psi \nabla \varphi$.

We can use the divergence theorem to derive many other useful identities.

We can use the divergence theorem to derive many other useful identities.

E.g. choosing $\mathbf{f} = \psi \mathbf{I}$, where \mathbf{I} is the identity matrix gives

$$\int_{\Omega} \nabla \cdot (\psi \mathbf{I}) dV = \int_{\Omega} \nabla \psi dV = \int_{\partial\Omega} \psi \mathbf{I} \cdot d\mathbf{S} = \int_{\partial\Omega} \psi d\mathbf{S}.$$