

# Wave equation and resonances

How radio works etc...

---

18.303 Linear Partial Differential Equations: Analysis and Numerics

Last time we talked about the Fourier transform when we were dealing with a function (time evolution of the Fourier coefficients for the wave equation) that is defined on the real line  $\mathbb{R}$ .

Last time we talked about the Fourier transform when we were dealing with a function (time evolution of the Fourier coefficients for the wave equation) that is defined on the real line  $\mathbb{R}$ .

Often in problems like this we are actually interested in how a system responds to some sort of input starting at time  $t = 0$ . This gives the domain  $\mathbb{R}_+$  i.e. the positive real numbers (normally including  $t = 0$ ).

Last time we talked about the Fourier transform when we were dealing with a function (time evolution of the Fourier coefficients for the wave equation) that is defined on the real line  $\mathbb{R}$ .

Often in problems like this we are actually interested in how a system responds to some sort of input starting at time  $t = 0$ . This gives the domain  $\mathbb{R}_+$  i.e. the positive real numbers (normally including  $t = 0$ ).

The natural tool for studying these sort of systems e.g. the time evolution of the wave equation is the Laplace transform.

# Laplace transform

We define a linear transformation  $\mathcal{L} : V \rightarrow W$ , where  $V$  and  $W$  are suitable spaces of functions. We write

$$\mathcal{L}[f](s) = \int_0^{\infty} e^{-st} f(t) dt =: F(s).$$

# Laplace transform

We define a linear transformation  $\mathcal{L} : V \rightarrow W$ , where  $V$  and  $W$  are suitable spaces of functions. We write

$$\mathcal{L}[f](s) = \int_0^{\infty} e^{-st} f(t) dt =: F(s).$$

This transform has some nice properties. For example, for derivatives of a function we have

$$\mathcal{L}[f'](s) = \int_0^{\infty} e^{-st} f'(t) dt$$

# Laplace transform

We define a linear transformation  $\mathcal{L} : V \rightarrow W$ , where  $V$  and  $W$  are suitable spaces of functions. We write

$$\mathcal{L}[f](s) = \int_0^{\infty} e^{-st} f(t) dt =: F(s).$$

This transform has some nice properties. For example, for derivatives of a function we have

$$\mathcal{L}[f'](s) = \int_0^{\infty} e^{-st} f'(t) dt$$

Integrating by part gives

$$e^{-st} f(t) \Big|_{t=0}^{\infty} + s \int_0^{\infty} e^{-st} f(t) dt = sF(s) - f(0^-).$$

Here  $f(0^-) = \lim_{t \rightarrow 0^-} f(t)$  is the left limit at zero. For continuous functions it is just  $f(0)$ .

Can you generalize this result for higher order derivatives? Try it for the second order.



Can you generalize this result for higher order derivatives? Try it for the second order.

$$\begin{aligned}\mathcal{L}[f^{(n)}](s) &= \int_0^\infty e^{-st} f^{(n)}(t) dt \\ &= e^{-st} f^{(n-1)}(t) \Big|_{t=0}^\infty + s \int_0^\infty e^{-st} f^{(n-1)}(t) dt \\ &= s\mathcal{L}[f^{(n-1)}](s) - f^{(n-1)}(0).\end{aligned}$$

Can you generalize this result for higher order derivatives? Try it for the second order.

$$\begin{aligned}\mathcal{L}[f^{(n)}](s) &= \int_0^{\infty} e^{-st} f^{(n)}(t) dt \\ &= e^{-st} f^{(n-1)}(t) \Big|_{t=0}^{\infty} + s \int_0^{\infty} e^{-st} f^{(n-1)}(t) dt \\ &= s \mathcal{L}[f^{(n-1)}](s) - f^{(n-1)}(0).\end{aligned}$$

Using the above formula repeatedly gives

$$\mathcal{L}[f^{(n)}](s) = s^n F(s) - s^{n-1} f(0) - s^{n-2} f'(0) - \dots - f^{(n-1)}(0).$$

Can you generalize this result for higher order derivatives? Try it for the second order.

$$\begin{aligned}\mathcal{L}[f^{(n)}](s) &= \int_0^\infty e^{-st} f^{(n)}(t) dt \\ &= e^{-st} f^{(n-1)}(t) \Big|_{t=0}^\infty + s \int_0^\infty e^{-st} f^{(n-1)}(t) dt \\ &= s \mathcal{L}[f^{(n-1)}](s) - f^{(n-1)}(0).\end{aligned}$$

Using the above formula repeatedly gives

$$\mathcal{L}[f^{(n)}](s) = s^n F(s) - s^{n-1} f(0) - s^{n-2} f'(0) - \dots - f^{(n-1)}(0).$$

We see immediately that this is perfect for initial value problems where derivatives at 0 are given.

We see that

$$F(2\pi i\xi) = \int_0^\infty e^{-2\pi i\xi t} f(t) dt = \hat{f}(\xi)$$

for functions with  $f(t) = 0$  if  $t < 0$ .

We see that

$$F(2\pi i\xi) = \int_0^{\infty} e^{-2\pi i\xi t} f(t) dt = \hat{f}(\xi)$$

for functions with  $f(t) = 0$  if  $t < 0$ .

**Strengths of Laplace transform:**

- The transform is real. This is sometimes very handy.

We see that

$$F(2\pi i\xi) = \int_0^{\infty} e^{-2\pi i\xi t} f(t) dt = \hat{f}(\xi)$$

for functions with  $f(t) = 0$  if  $t < 0$ .

## Strengths of Laplace transform:

- The transform is real. This is sometimes very handy.
- The transform exists for a larger class of functions since the Laplace integral converges very strongly.

We see that

$$F(2\pi i\xi) = \int_0^{\infty} e^{-2\pi i\xi t} f(t) dt = \hat{f}(\xi)$$

for functions with  $f(t) = 0$  if  $t < 0$ .

## Strengths of Laplace transform:

- The transform is real. This is sometimes very handy.
- The transform exists for a larger class of functions since the Laplace integral converges very strongly.
- Can be evaluated with efficient algorithms for Fourier transform.

For example, let us calculate the Laplace transform for a function that is 0 for  $t < 0$  and  $\sin(\omega t)$  for  $t > 0$ :

$$\mathcal{L}[\sin(\omega t)](s) = \int_0^{\infty} e^{-st} \sin(\omega t) dt$$



For example, let us calculate the Laplace transform for a function that is 0 for  $t < 0$  and  $\sin(\omega t)$  for  $t > 0$ :

$$\mathcal{L}[\sin(\omega t)](s) = \int_0^{\infty} e^{-st} \sin(\omega t) dt$$

Casting the  $\sin$  function in the exponential form gives

$$\begin{aligned} \int_0^{\infty} e^{-st} \frac{1}{2i} (e^{i\omega t} - e^{-i\omega t}) dt &= \frac{1}{2i} \int_0^{\infty} e^{(i\omega-s)t} - e^{-(i\omega+s)t} dt \\ &= \frac{1}{2i} \left( \frac{1}{s-i\omega} - \frac{1}{s+i\omega} \right) \\ &= \frac{\omega}{s^2 + \omega^2}. \end{aligned}$$

We can try doing the same for the Fourier transform:

$$\begin{aligned}\mathcal{F}[\sin(\omega t)](\xi) &= \int_0^\infty e^{-2\pi i \xi t} \sin(\omega t) dt \\ &= \frac{1}{2i} \int_0^\infty e^{(\omega - 2\pi \xi)it} - e^{-(\omega + 2\pi \xi)it} dt.\end{aligned}$$

We can try doing the same for the Fourier transform:

$$\begin{aligned}\mathcal{F}[\sin(\omega t)](\xi) &= \int_0^\infty e^{-2\pi i \xi t} \sin(\omega t) dt \\ &= \frac{1}{2i} \int_0^\infty e^{(\omega - 2\pi \xi)it} - e^{-(\omega + 2\pi \xi)it} dt.\end{aligned}$$

Since the integrand is oscillating infinitely, this integral doesn't converge in the traditional sense.

We can try doing the same for the Fourier transform:

$$\begin{aligned}\mathcal{F}[\sin(\omega t)](\xi) &= \int_0^\infty e^{-2\pi i \xi t} \sin(\omega t) dt \\ &= \frac{1}{2i} \int_0^\infty e^{(\omega - 2\pi \xi)it} - e^{-(\omega + 2\pi \xi)it} dt.\end{aligned}$$

Since the integrand is oscillating infinitely, this integral doesn't converge in the traditional sense.

However, this integral can be regularized by considering  $\mathcal{F}[\sin(\omega t)](\xi - i\epsilon)$  and taking  $\epsilon \rightarrow 0$ .

We can try doing the same for the Fourier transform:

$$\begin{aligned}\mathcal{F}[\sin(\omega t)](\xi) &= \int_0^\infty e^{-2\pi i \xi t} \sin(\omega t) dt \\ &= \frac{1}{2i} \int_0^\infty e^{(\omega - 2\pi i \xi)it} - e^{-(\omega + 2\pi i \xi)it} dt.\end{aligned}$$

Since the integrand is oscillating infinitely, this integral doesn't converge in the traditional sense.

However, this integral can be regularized by considering  $\mathcal{F}[\sin(\omega t)](\xi - i\epsilon)$  and taking  $\epsilon \rightarrow 0$ .

This gives

$$\mathcal{F}[\sin(\omega t)](\xi) = \frac{1}{2i} \left( \frac{1}{2\pi i \xi - i\omega} - \frac{1}{2\pi i \xi + i\omega} \right) = \frac{\omega}{\omega^2 - (2\pi \xi)^2}.$$

We can try doing the same for the Fourier transform:

$$\begin{aligned}\mathcal{F}[\sin(\omega t)](\xi) &= \int_0^\infty e^{-2\pi i \xi t} \sin(\omega t) dt \\ &= \frac{1}{2i} \int_0^\infty e^{(\omega - 2\pi \xi)it} - e^{-(\omega + 2\pi \xi)it} dt.\end{aligned}$$

Since the integrand is oscillating infinitely, this integral doesn't converge in the traditional sense.

However, this integral can be regularized by considering  $\mathcal{F}[\sin(\omega t)](\xi - i\epsilon)$  and taking  $\epsilon \rightarrow 0$ .

This gives

$$\mathcal{F}[\sin(\omega t)](\xi) = \frac{1}{2i} \left( \frac{1}{2\pi i \xi - i\omega} - \frac{1}{2\pi i \xi + i\omega} \right) = \frac{\omega}{\omega^2 - (2\pi \xi)^2}.$$

We can recognize this as the Laplace transform evaluated at  $s = 2\pi i \xi$  just as it should be.

This brings us to the **downsides of Laplace transform**:

- Higher dimensional generalizations are not so useful since we hardly need the positive quadrant of  $\mathbb{R}^n$ .

This brings us to the **downsides of Laplace transform**:

- Higher dimensional generalizations are not so useful since we hardly need the positive quadrant of  $\mathbb{R}^n$ .
- Inverse transform requires extending the Laplace transform  $F(s)$  to the complex plane and sometimes requires some regularization magic.



# Inverse Laplace transform

Let's use the relationship between Fourier and Laplace transforms

$$F(2\pi i\xi) = \int_0^{\infty} e^{-2\pi i\xi t} f(t) dt = \hat{f}(\xi)$$

to derive the inverse transform.

# Inverse Laplace transform

Let's use the relationship between Fourier and Laplace transforms

$$F(2\pi i\xi) = \int_0^{\infty} e^{-2\pi i\xi t} f(t) dt = \hat{f}(\xi)$$

to derive the inverse transform.

Taking the inverse Fourier transform gives

$$f(t) = \int_{-\infty}^{\infty} e^{2\pi i\xi t} F(2\pi i\xi) d\xi.$$

# Inverse Laplace transform

Let's use the relationship between Fourier and Laplace transforms

$$F(2\pi i\xi) = \int_0^{\infty} e^{-2\pi i\xi t} f(t) dt = \hat{f}(\xi)$$

to derive the inverse transform.

Taking the inverse Fourier transform gives

$$f(t) = \int_{-\infty}^{\infty} e^{2\pi i\xi t} F(2\pi i\xi) d\xi.$$

Making the change of variables  $s = 2\pi i\xi$  gives

$$f(t) = \frac{1}{2\pi i} \int_{-i\infty}^{i\infty} e^{st} F(s) ds.$$

# Inverse Laplace transform

Let's use the relationship between Fourier and Laplace transforms

$$F(2\pi i\xi) = \int_0^{\infty} e^{-2\pi i\xi t} f(t) dt = \hat{f}(\xi)$$

to derive the inverse transform.

Taking the inverse Fourier transform gives

$$f(t) = \int_{-\infty}^{\infty} e^{2\pi i\xi t} F(2\pi i\xi) d\xi.$$

Making the change of variables  $s = 2\pi i\xi$  gives

$$f(t) = \frac{1}{2\pi i} \int_{-i\infty}^{i\infty} e^{st} F(s) ds.$$

We see that the integral has to be evaluated on the imaginary axis of the complex plane.

Let us try to calculate the inverse Laplace transform of the function

$$F(s) = \frac{\omega_0}{s^2 + \omega_0^2}.$$

Let us try to calculate the inverse Laplace transform of the function

$$F(s) = \frac{\omega_0}{s^2 + \omega_0^2}.$$

We know from before that this is the Laplace transform of  $\sin(\omega_0 t)$ . We have

$$\frac{1}{2\pi i} \int_{-i\infty}^{i\infty} e^{st} \frac{\omega_0}{s^2 + \omega_0^2} ds.$$

Let us try to calculate the inverse Laplace transform of the function

$$F(s) = \frac{\omega_0}{s^2 + \omega_0^2}.$$

We know from before that this is the Laplace transform of  $\sin(\omega_0 t)$ . We have

$$\frac{1}{2\pi i} \int_{-i\infty}^{i\infty} e^{st} \frac{\omega_0}{s^2 + \omega_0^2} ds.$$

Making the change of variable  $s = i\omega$  gives

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} e^{i\omega t} \frac{\omega_0}{\omega_0^2 - \omega^2} d\omega = -\frac{1}{2\pi} \int_{-\infty}^{\infty} e^{i\omega t} \frac{\omega_0}{(\omega - \omega_0)(\omega + \omega_0)} d\omega.$$

Let us try to calculate the inverse Laplace transform of the function

$$F(s) = \frac{\omega_0}{s^2 + \omega_0^2}.$$

We know from before that this is the Laplace transform of  $\sin(\omega_0 t)$ . We have

$$\frac{1}{2\pi i} \int_{-i\infty}^{i\infty} e^{st} \frac{\omega_0}{s^2 + \omega_0^2} ds.$$

Making the change of variable  $s = i\omega$  gives

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} e^{i\omega t} \frac{\omega_0}{\omega_0^2 - \omega^2} d\omega = -\frac{1}{2\pi} \int_{-\infty}^{\infty} e^{i\omega t} \frac{\omega_0}{(\omega - \omega_0)(\omega + \omega_0)} d\omega.$$

We notice that the integral has two singularities at  $\omega = \pm\omega_0$  and thus can't be in general calculated.



The answer comes from regularization. If we had instead

$$-\frac{1}{2\pi} \int_{-\infty-i\epsilon}^{\infty-i\epsilon} e^{i\omega t} \frac{\omega_0}{(\omega - \omega_0)(\omega + \omega_0)} d\omega$$

we wouldn't be integrating over the singularities on complex plane.

The answer comes from regularization. If we had instead

$$-\frac{1}{2\pi} \int_{-\infty-i\epsilon}^{\infty-i\epsilon} e^{i\omega t} \frac{\omega_0}{(\omega - \omega_0)(\omega + \omega_0)} d\omega$$

we wouldn't be integrating over the singularities on complex plane.

The above expression can be turned into a contour integral on complex plane so in fact it doesn't matter how large  $\epsilon$  is.

The answer comes from regularization. If we had instead

$$-\frac{1}{2\pi} \int_{-\infty-i\epsilon}^{\infty-i\epsilon} e^{i\omega t} \frac{\omega_0}{(\omega - \omega_0)(\omega + \omega_0)} d\omega$$

we wouldn't be integrating over the singularities on complex plane.

The above expression can be turned into a contour integral on complex plane so in fact it doesn't matter how large  $\epsilon$  is.

This integral is evaluated by the residues i.e.

$$f(t) = -\frac{1}{2\pi} \oint_{\gamma} g(w) dw = \frac{1}{i} [\text{Res}(g, \omega_0) + \text{Res}(g, -\omega_0)]$$

The answer comes from regularization. If we had instead

$$-\frac{1}{2\pi} \int_{-\infty-i\epsilon}^{\infty-i\epsilon} e^{i\omega t} \frac{\omega_0}{(\omega - \omega_0)(\omega + \omega_0)} d\omega$$

we wouldn't be integrating over the singularities on complex plane.

The above expression can be turned into a contour integral on complex plane so in fact it doesn't matter how large  $\epsilon$  is.

This integral is evaluated by the residues i.e.

$$f(t) = -\frac{1}{2\pi} \oint_{\gamma} g(w) dw = \frac{1}{i} [\text{Res}(g, \omega_0) + \text{Res}(g, -\omega_0)]$$

Evaluating this gives

$$f(t) = \frac{\omega_0}{i} \left( \frac{e^{i\omega_0 t}}{2\omega_0} - \frac{e^{-i\omega_0 t}}{2\omega_0} \right) = \sin(\omega_0 t).$$

If  $t < 0$  it turns out that this integral gives 0, as it should.

The formula we obtained can be modified to work for these difficult functions. In general we have

$$\mathcal{L}^{-1}[F](t) = \frac{1}{2\pi i} \int_{-i\infty+\Delta}^{i\infty+\Delta} e^{st} F(s) ds,$$

where  $\Delta$  is a real number s.t. it is larger than any real parts of singularities of  $F(s)$  on the complex plane.

The formula we obtained can be modified to work for these difficult functions. In general we have

$$\mathcal{L}^{-1}[F](t) = \frac{1}{2\pi i} \int_{-i\infty+\Delta}^{i\infty+\Delta} e^{st} F(s) ds,$$

where  $\Delta$  is a real number s.t. it is larger than any real parts of singularities of  $F(s)$  on the complex plane.

So, in short, inverse Laplace transforms are a bit more complicated. However, if  $F$  is an entire function on  $\mathbb{C}$  without singularities, we can set  $\Delta = 0$ .

# Resonances

Consider the ODE

$$u''(t) + \lambda^2 u(t) = \sin(\omega_0 t),$$

with initial conditions  $u'(0) = u(0) = 0$ . With different initial conditions we would have to be concerned with the solution to the homogeneous equation, which we know very well how to do at this point.

# Resonances

Consider the ODE

$$u''(t) + \lambda^2 u(t) = \sin(\omega_0 t),$$

with initial conditions  $u'(0) = u(0) = 0$ . With different initial conditions we would have to be concerned with the solution to the homogeneous equation, which we know very well how to do at this point.

Notice here that if  $\lambda \rightarrow \lambda_n$ , this is the equation we obtain for each eigenvalue of the inhomogeneous wave equation (possibly there's something time independent multiplying the RHS).



# Resonances

Consider the ODE

$$u''(t) + \lambda^2 u(t) = \sin(\omega_0 t),$$

with initial conditions  $u'(0) = u(0) = 0$ . With different initial conditions we would have to be concerned with the solution to the homogeneous equation, which we know very well how to do at this point.

Notice here that if  $\lambda \rightarrow \lambda_n$ , this is the equation we obtain for each eigenvalue of the inhomogeneous wave equation (possibly there's something time independent multiplying the RHS).

In general this is solved by an ansatz

$$u(t) = A \sin(\omega_0 t) + B \cos(\omega_0 t).$$

Plugging this in gives

$$(\lambda^2 - \omega_0^2)(A \sin(\omega_0 t) + B \cos(\omega_0 t)) = \sin(\omega_0 t).$$

From this we can solve

$$A = \frac{1}{\lambda^2 - \omega_0^2}, \quad B = 0,$$

as long as  $\omega_0 \neq \pm\lambda$ .

From this we can solve

$$A = \frac{1}{\lambda^2 - \omega_0^2}, \quad B = 0,$$

as long as  $\omega_0 \neq \pm\lambda$ .

What if  $\omega_0 = \lambda$ ?

From this we can solve

$$A = \frac{1}{\lambda^2 - \omega_0^2}, \quad B = 0,$$

as long as  $\omega_0 \neq \pm\lambda$ .

What if  $\omega_0 = \lambda$ ?

In this case this is solved by an ansatz

$u(t) = At \sin(\omega_0 t) + Bt \cos(\omega_0 t) = t(A \sin(\lambda t) + B \cos(\lambda t))$ . Plugging this in gives

$$-t\lambda^2(A \sin(\lambda t) + B \cos(\lambda t)) + 2\lambda(A \cos(\lambda t) - B \sin(\lambda t)) + t\lambda^2(A \sin(\lambda t) + B \cos(\lambda t)) = \sin(\lambda t).$$

From this we can solve

$$A = \frac{1}{\lambda^2 - \omega_0^2}, \quad B = 0,$$

as long as  $\omega_0 \neq \pm\lambda$ .

What if  $\omega_0 = \lambda$ ?

In this case this is solved by an ansatz

$u(t) = At \sin(\omega_0 t) + Bt \cos(\omega_0 t) = t(A \sin(\lambda t) + B \cos(\lambda t))$ . Plugging this in gives

$$-t\lambda^2(A \sin(\lambda t) + B \cos(\lambda t)) + 2\lambda(A \cos(\lambda t) - B \sin(\lambda t)) + t\lambda^2(A \sin(\lambda t) + B \cos(\lambda t)) = \sin(\lambda t).$$

Equating this with the RHS gives

From this we can solve

$$A = \frac{1}{\lambda^2 - \omega_0^2}, \quad B = 0,$$

as long as  $\omega_0 \neq \pm\lambda$ .

What if  $\omega_0 = \lambda$ ?

In this case this is solved by an ansatz

$u(t) = At \sin(\omega_0 t) + Bt \cos(\omega_0 t) = t(A \sin(\lambda t) + B \cos(\lambda t))$ . Plugging this in gives

$$-t\lambda^2(A \sin(\lambda t) + B \cos(\lambda t)) + 2\lambda(A \cos(\lambda t) - B \sin(\lambda t)) + t\lambda^2(A \sin(\lambda t) + B \cos(\lambda t)) = \sin(\lambda t).$$

Equating this with the RHS gives

$$A = 0, B = -\frac{1}{2\lambda}$$

and the solution is

$$u(t) = -\frac{t}{2\lambda} \cos(\lambda t).$$

Notice how the amplitude of this solution grows linearly in time. This is called a resonant frequency.

Notice how the amplitude of this solution grows linearly in time. This is called a resonant frequency.

Imagine that we would have a wave equation (we'll do this in detail later) with eigenvalues  $\lambda_n$ . If we have an input that is oscillating at a frequency  $f = \lambda_n/(2\pi)$  the system starts to vibrate uncontrollably.



Notice how the amplitude of this solution grows linearly in time. This is called a resonant frequency.

Imagine that we would have a wave equation (we'll do this in detail later) with eigenvalues  $\lambda_n$ . If we have an input that is oscillating at a frequency  $f = \lambda_n/(2\pi)$  the system starts to vibrate uncontrollably.

Radio is an example of a system where this phenomenon is made useful. Radio circuit is basically an electronic oscillator whose eigenmodes can be modified. By tuning the eigenmode to match the external frequency the oscillations increase greatly and we can listen to a radio channel at a given frequency.

Notice how the amplitude of this solution grows linearly in time. This is called a resonant frequency.

Imagine that we would have a wave equation (we'll do this in detail later) with eigenvalues  $\lambda_n$ . If we have an input that is oscillating at a frequency  $f = \lambda_n/(2\pi)$  the system starts to vibrate uncontrollably.

Radio is an example of a system where this phenomenon is made useful. Radio circuit is basically an electronic oscillator whose eigenmodes can be modified. By tuning the eigenmode to match the external frequency the oscillations increase greatly and we can listen to a radio channel at a given frequency.

Of course nothing blows up in nature. Next time we will talk about softening of resonances due to some sort of damping in the system. We will also derive the results above using the Laplace transform. It is recommended to revise complex analysis for this.