

# Inhomogeneous PDEs revisited

The inhomogeneous heat and wave equations

---

18.303 Linear Partial Differential Equations: Analysis and Numerics


# Basic Notions

Boundary value problems look basically like this:

$$\mathcal{L}u(\mathbf{x}) = f(\mathbf{x}), \mathbf{x} \in \omega;$$

$$\mathcal{G}u(\mathbf{x}) = g(\mathbf{x}), \mathbf{x} \in \partial\omega.$$

Now, if  $\mathbf{f} = 0$ , we say that the boundary value problem (the differential equation) is **homogeneous**. Otherwise the problem is said to be **inhomogeneous**.



An example of a 2d domain and its boundary.


# Basic Notions

Boundary value problems look basically like this:

$$\mathcal{L}u(\mathbf{x}) = f(\mathbf{x}), \mathbf{x} \in \omega;$$

$$\mathcal{G}u(\mathbf{x}) = g(\mathbf{x}), \mathbf{x} \in \partial\omega.$$

If the differential operator  $\mathcal{G}$  is just a (non-zero) function, these are **Dirichlet boundaries** and  $u$  is equal to some function on the boundary  $\partial\omega$ .



An example of a 2d domain and its boundary.

# Basic Notions


Boundary value problems look basically like this:

$$\mathcal{L}u(\mathbf{x}) = f(\mathbf{x}), \mathbf{x} \in \omega;$$

$$\mathcal{G}u(\mathbf{x}) = g(\mathbf{x}), \mathbf{x} \in \partial\omega.$$

If the differential operator  $\mathcal{G}$  is just a (non-zero) function, these are **Dirichlet boundaries** and  $u$  is equal to some function on the boundary  $\partial\omega$ .

If the differential operator is  $\hat{\mathbf{n}} \cdot \nabla = \frac{\partial}{\partial \hat{\mathbf{n}}}$  the boundary condition is called **Neumann type** or **flux boundary** condition. This denotes the derivative of function  $u$  in the normal direction of the boundary given by the normal vector  $\hat{\mathbf{n}}(\mathbf{x})$  (note that generally it depends on the point  $\mathbf{x} \in \partial\omega$ ).



An example of a 2d domain and its boundary.

Boundary conditions can be more complicated than Dirichlet or Neumann conditions (and will be e.g. for higher order differential operators). However, we have the following general property:

Boundary conditions can be more complicated than Dirichlet or Neumann conditions (and will be e.g. for higher order differential operators). However, we have the following general property:

Let  $u_0(\mathbf{x})$  solve the *homogeneous* problem

$$\mathcal{L}u_0(\mathbf{x}) = 0, \mathbf{x} \in \omega;$$

$$\mathcal{G}u_0(\mathbf{x}) = g(\mathbf{x}), \mathbf{x} \in \partial\omega.$$

Boundary conditions can be more complicated than Dirichlet or Neumann conditions (and will be e.g. for higher order differential operators). However, we have the following general property:

Let  $u_0(\mathbf{x})$  solve the *homogeneous* problem

$$\mathcal{L}u_0(\mathbf{x}) = 0, \mathbf{x} \in \omega;$$

$$\mathcal{G}u_0(\mathbf{x}) = g(\mathbf{x}), \mathbf{x} \in \partial\omega.$$

Now, consider the function  $u(\mathbf{x}) = v(\mathbf{x}) + u_0(\mathbf{x})$ . We have

$$\mathcal{L}u(\mathbf{x}) = \mathcal{L}v(\mathbf{x}) + \underbrace{\mathcal{L}u_0(\mathbf{x})}_{=0} = f(\mathbf{x}).$$

Boundary conditions can be more complicated than Dirichlet or Neumann conditions (and will be e.g. for higher order differential operators). However, we have the following general property:

Let  $u_0(\mathbf{x})$  solve the *homogeneous* problem

$$\begin{aligned}\mathcal{L}u_0(\mathbf{x}) &= 0, \quad \mathbf{x} \in \omega; \\ \mathcal{G}u_0(\mathbf{x}) &= g(\mathbf{x}), \quad \mathbf{x} \in \partial\omega.\end{aligned}$$

Now, consider the function  $u(\mathbf{x}) = v(\mathbf{x}) + u_0(\mathbf{x})$ . We have

$$\mathcal{L}u(\mathbf{x}) = \mathcal{L}v(\mathbf{x}) + \underbrace{\mathcal{L}u_0(\mathbf{x})}_{=0} = f(\mathbf{x}).$$

We notice that it suffices to require that

$$\mathcal{G}v(\mathbf{x}) = 0, \quad \mathbf{x} \in \partial\omega$$

because this will make sure that  $\mathcal{G}u(\mathbf{x}) = g(\mathbf{x})$ , when  $\mathbf{x} \in \partial\omega$ .



Now we have a boundary value problem

$$\mathcal{L}v(\mathbf{x}) = f(\mathbf{x}), \mathbf{x} \in \omega;$$

$$\mathcal{G}v(\mathbf{x}) = 0, \mathbf{x} \in \partial\omega.$$

This shows us that the general solution can be sought as

the solution to the homogeneous problem with inhomogeneous boundaries + the solution to the inhomogeneous problem with homogeneous boundaries.

# Heat equation

We have

$$\frac{\partial}{\partial t}u(t, \mathbf{x}) = \Delta u(t, \mathbf{x}) + f(t, \mathbf{x})$$

with some spatial boundary conditions for all  $t$  and in addition we are given

$$u(0, \mathbf{x}) = u^{(0)}(\mathbf{x}).$$

# Heat equation

We have

$$\frac{\partial}{\partial t}u(t, \mathbf{x}) = \Delta u(t, \mathbf{x}) + f(t, \mathbf{x})$$

with some spatial boundary conditions for all  $t$  and in addition we are given

$$u(0, \mathbf{x}) = u^{(0)}(\mathbf{x}).$$

We express  $u$  in the eigenbasis  $\{\phi_n\}$  of the Laplacian  $\Delta$  giving

$$u(t, \mathbf{x}) = \sum_n \hat{u}_n(t) \phi_n(\mathbf{x}).$$

Similarly,

$$f(t, \mathbf{x}) = \sum_n \hat{f}_n(t) \phi_n(\mathbf{x}).$$

Because the basis is linearly independent, we get

$$\frac{\partial}{\partial t} \hat{u}_n(t) = -\lambda_n^2 \hat{u}_n(t) + \hat{f}_n(t).$$

Because the basis is linearly independent, we get

$$\frac{\partial}{\partial t} \hat{u}_n(t) = -\lambda_n^2 \hat{u}_n(t) + \hat{f}_n(t).$$

Multiplying both sides by  $\exp(\lambda_n^2 t)$  and reorganizing gives

$$\frac{\partial}{\partial t} \hat{u}_n(t) e^{\lambda_n^2 t} + \lambda_n^2 \hat{u}_n(t) e^{\lambda_n^2 t} = \hat{f}_n(t) e^{\lambda_n^2 t}.$$

Because the basis is linearly independent, we get

$$\frac{\partial}{\partial t} \hat{u}_n(t) = -\lambda_n^2 \hat{u}_n(t) + \hat{f}_n(t).$$

Multiplying both sides by  $\exp(\lambda_n^2 t)$  and reorganizing gives

$$\frac{\partial}{\partial t} \hat{u}_n(t) e^{\lambda_n^2 t} + \lambda_n^2 \hat{u}_n(t) e^{\lambda_n^2 t} = \hat{f}_n(t) e^{\lambda_n^2 t}.$$

We notice that we can write this as

$$\frac{\partial}{\partial t} \left( \hat{u}_n(t) e^{\lambda_n^2 t} \right) = \hat{f}_n(t) e^{\lambda_n^2 t},$$

which we can integrate from 0 to  $t$  giving

$$\hat{u}_n(t) e^{\lambda_n^2 t} - \hat{u}_n(0) = \int_0^t \hat{f}_n(t') e^{\lambda_n^2 t'} dt'$$

We notice that  $\hat{u}_n(0)$  is given by the coefficients of the initial condition

$$\hat{u}_n(0) = \hat{u}_n^{(0)}.$$

We notice that  $\hat{u}_n(0)$  is given by the coefficients of the initial condition

$$\hat{u}_n(0) = \hat{u}_n^{(0)}.$$

Now we can write the complete solution for the coefficients as

$$\hat{u}_n(t) = \hat{u}_n^{(0)} e^{-\lambda_n^2 t} + \int_0^t \hat{f}_n(t') e^{-\lambda_n^2 (t-t')} dt'.$$

We see immediately that for  $f = 0$  we get the usual solution to the homogeneous problem.



We notice that  $\hat{u}_n(0)$  is given by the coefficients of the initial condition

$$\hat{u}_n(0) = \hat{u}_n^{(0)}.$$

Now we can write the complete solution for the coefficients as

$$\hat{u}_n(t) = \hat{u}_n^{(0)} e^{-\lambda_n^2 t} + \int_0^t \hat{f}_n(t') e^{-\lambda_n^2 (t-t')} dt'.$$

We see immediately that for  $f = 0$  we get the usual solution to the homogeneous problem.

What if  $f$  is independent of time?

This gives

$$\begin{aligned}\hat{u}_n(t) &= \hat{u}_n^{(0)} e^{-\lambda_n^2 t} + \hat{f}_n \int_0^t e^{-\lambda_n^2(t-t')} dt' \\ &= \hat{u}_n^{(0)} e^{-\lambda_n^2 t} + \frac{\hat{f}_n}{\lambda_n^2} \left( e^{-\lambda_n^2(t-t')} \right)_{t'=0}^t \\ &= \hat{u}_n^{(0)} e^{-\lambda_n^2 t} + \frac{\hat{f}_n}{\lambda_n^2} \left( 1 - e^{-\lambda_n^2 t} \right).\end{aligned}$$

This gives

$$\begin{aligned}\hat{u}_n(t) &= \hat{u}_n^{(0)} e^{-\lambda_n^2 t} + \hat{f}_n \int_0^t e^{-\lambda_n^2(t-t')} dt' \\ &= \hat{u}_n^{(0)} e^{-\lambda_n^2 t} + \frac{\hat{f}_n}{\lambda_n^2} \left( e^{-\lambda_n^2(t-t')} \right)_{t'=0}^t \\ &= \hat{u}_n^{(0)} e^{-\lambda_n^2 t} + \frac{\hat{f}_n}{\lambda_n^2} \left( 1 - e^{-\lambda_n^2 t} \right).\end{aligned}$$

What if  $t \rightarrow \infty$ ?

This gives

$$\begin{aligned}\hat{u}_n(t) &= \hat{u}_n^{(0)} e^{-\lambda_n^2 t} + \hat{f}_n \int_0^t e^{-\lambda_n^2(t-t')} dt' \\ &= \hat{u}_n^{(0)} e^{-\lambda_n^2 t} + \frac{\hat{f}_n}{\lambda_n^2} \left( e^{-\lambda_n^2(t-t')} \right)_{t'=0}^t \\ &= \hat{u}_n^{(0)} e^{-\lambda_n^2 t} + \frac{\hat{f}_n}{\lambda_n^2} (1 - e^{-\lambda_n^2 t}).\end{aligned}$$

What if  $t \rightarrow \infty$ ?

We obtain the solution to the Poisson equation

$$\hat{u}_n(t) = \frac{\hat{f}_n}{\lambda_n^2}.$$

This result is not too surprising if we go back to the original heat equation

$$\frac{\partial}{\partial t}u(t, \mathbf{x}) = \Delta u(t, \mathbf{x}) + f(t, \mathbf{x}).$$

This result is not too surprising if we go back to the original heat equation

$$\frac{\partial}{\partial t}u(t, \mathbf{x}) = \Delta u(t, \mathbf{x}) + f(t, \mathbf{x}).$$

Just assuming that we will reach a **steady state** with  $\partial_t u = 0$  gives

$$-\Delta u(t, \mathbf{x}) = f(t, \mathbf{x}).$$

# Wave equation

Now

$$\frac{\partial^2}{\partial t^2} u(t, \mathbf{x}) = \Delta u(t, \mathbf{x}) + f(t, \mathbf{x})$$

with some appropriate boundary conditions and initial conditions

$$\begin{aligned} u(0, \mathbf{x}) &= u^{(0)}(\mathbf{x}), \\ \left( \frac{\partial}{\partial t} u(t, \mathbf{x}) \right)_{t=0} &= v^{(0)}(\mathbf{x}). \end{aligned}$$

Now

$$\frac{\partial^2}{\partial t^2} u(t, \mathbf{x}) = \Delta u(t, \mathbf{x}) + f(t, \mathbf{x})$$

with some appropriate boundary conditions and initial conditions

$$\begin{aligned} u(0, \mathbf{x}) &= u^{(0)}(\mathbf{x}), \\ \left( \frac{\partial}{\partial t} u(t, \mathbf{x}) \right)_{t=0} &= v^{(0)}(\mathbf{x}). \end{aligned}$$

We proceed in a same way by writing the equation for the coefficients  $\hat{u}_n(t)$  as

$$\frac{\partial^2}{\partial t^2} \hat{u}_n(t) = -\lambda_n^2 \hat{u}_n(t) + \hat{f}_n(t).$$



# Wave equation

Now

$$\frac{\partial^2}{\partial t^2} u(t, \mathbf{x}) = \Delta u(t, \mathbf{x}) + f(t, \mathbf{x})$$

with some appropriate boundary conditions and initial conditions

$$\begin{aligned} u(0, \mathbf{x}) &= u^{(0)}(\mathbf{x}), \\ \left( \frac{\partial}{\partial t} u(t, \mathbf{x}) \right)_{t=0} &= v^{(0)}(\mathbf{x}). \end{aligned}$$

We proceed in a same way by writing the equation for the coefficients  $\hat{u}_n(t)$  as

$$\frac{\partial^2}{\partial t^2} \hat{u}_n(t) = -\lambda_n^2 \hat{u}_n(t) + \hat{f}_n(t).$$

This is an ordinary differential equation (ODE) but it can be a bit harder to solve.

Recall that the complete solution is the solution to the homogeneous problem with the necessary initial (boundary) conditions + the solution to the inhomogeneous equation with zero boundaries.

Recall that the complete solution is the solution to the homogeneous problem with the necessary initial (boundary) conditions + the solution to the inhomogeneous equation with zero boundaries.

We have already covered the solution to the homogeneous problem (here we write  $h(t, \mathbf{x})$ ) and the coefficients are given by

$$\hat{h}_n(t) = \alpha_n \cos(\lambda_n t) + \beta_n \sin(\lambda_n t),$$

where the coefficients  $\alpha_n$  and  $\beta_n$  can be solved from the initial conditions. (For complex equations we have  $\exp(\pm i\lambda_n t)$  instead of  $\sin$  and  $\cos$ ).

Recall that the complete solution is the solution to the homogeneous problem with the necessary initial (boundary) conditions + the solution to the inhomogeneous equation with zero boundaries.

We have already covered the solution to the homogeneous problem (here we write  $h(t, \mathbf{x})$ ) and the coefficients are given by

$$\hat{h}_n(t) = \alpha_n \cos(\lambda_n t) + \beta_n \sin(\lambda_n t),$$

where the coefficients  $\alpha_n$  and  $\beta_n$  can be solved from the initial conditions. (For complex equations we have  $\exp(\pm i\lambda_n t)$  instead of  $\sin$  and  $\cos$ ).

We can seek the particular solution to the inhomogeneous problem for the coefficients in the form

$$\hat{p}_n(t) = \hat{\xi}_n(t) \hat{h}_n(t).$$

Substituting this into the inhomogeneous equation for the coefficients gives

$$\begin{aligned}\hat{f}_n(t) &= \frac{\partial^2}{\partial t^2} \left( \hat{\xi}_n(t) \hat{h}_n(t) \right) + \lambda_n^2 \hat{\xi}_n(t) \hat{h}_n(t) \\ &= \hat{\xi}_n''(t) \hat{h}_n + 2 \hat{\xi}_n'(t) \hat{h}_n'(t) + \hat{h}_n''(t) \hat{\xi}_n(t) + \lambda_n(t)^2 \hat{h}_n(t) \hat{\xi}_n(t).\end{aligned}$$

Substituting this into the inhomogeneous equation for the coefficients gives

$$\begin{aligned}\hat{f}_n(t) &= \frac{\partial^2}{\partial t^2} \left( \hat{\xi}_n(t) \hat{h}_n(t) \right) + \lambda_n^2 \hat{\xi}_n(t) \hat{h}_n(t) \\ &= \hat{\xi}_n''(t) \hat{h}_n + 2 \hat{\xi}_n'(t) \hat{h}_n'(t) + \hat{h}_n''(t) \hat{\xi}_n(t) + \lambda_n(t)^2 \hat{h}_n(t) \hat{\xi}_n(t).\end{aligned}$$

The last two terms solve the homogeneous problem

$$\hat{\xi}_n(t) \underbrace{\left( \hat{h}_n''(t) + \lambda_n(t)^2 \hat{h}_n(t) \right)}_{=0}.$$

Substituting this into the inhomogeneous equation for the coefficients gives

$$\begin{aligned}\hat{f}_n(t) &= \frac{\partial^2}{\partial t^2} \left( \hat{\xi}_n(t) \hat{h}_n(t) \right) + \lambda_n^2 \hat{\xi}_n(t) \hat{h}_n(t) \\ &= \hat{\xi}_n''(t) \hat{h}_n(t) + 2 \hat{\xi}_n'(t) \hat{h}_n'(t) + \hat{h}_n''(t) \hat{\xi}_n(t) + \lambda_n(t)^2 \hat{h}_n(t) \hat{\xi}_n(t).\end{aligned}$$

The last two terms solve the homogeneous problem

$$\hat{\xi}_n(t) \underbrace{\left( \hat{h}_n''(t) + \lambda_n(t)^2 \hat{h}_n(t) \right)}_{=0}.$$

This gives

$$\hat{\xi}_n''(t) \hat{h}_n(t) + 2 \hat{\xi}_n'(t) \hat{h}_n'(t) = \hat{f}_n(t).$$

Substituting this into the inhomogeneous equation for the coefficients gives

$$\begin{aligned}\hat{f}_n(t) &= \frac{\partial^2}{\partial t^2} \left( \hat{\xi}_n(t) \hat{h}_n(t) \right) + \lambda_n^2 \hat{\xi}_n(t) \hat{h}_n(t) \\ &= \hat{\xi}_n''(t) \hat{h}_n + 2 \hat{\xi}_n'(t) \hat{h}_n'(t) + \hat{h}_n''(t) \hat{\xi}_n(t) + \lambda_n(t)^2 \hat{h}_n(t) \hat{\xi}_n(t).\end{aligned}$$

The last two terms solve the homogeneous problem

$$\hat{\xi}_n(t) \underbrace{\left( \hat{h}_n''(t) + \lambda_n(t)^2 \hat{h}_n(t) \right)}_{=0}.$$

This gives

$$\hat{\xi}_n''(t) \hat{h}_n(t) + 2 \hat{\xi}_n'(t) \hat{h}_n'(t) = \hat{f}_n(t).$$

Multiplying by  $\hat{h}_n$  gives

$$\hat{\xi}_n''(t) \hat{h}_n(t)^2 + 2 \hat{\xi}_n'(t) \hat{h}_n'(t) \hat{h}_n(t) = \hat{f}_n(t) \hat{h}_n(t),$$

which we can write as

$$\frac{\partial}{\partial t} \left( \hat{\xi}_n'(t) \hat{h}_n(t)^2 \right) = \hat{f}_n(t) \hat{h}_n(t).$$



Now this can be integrated from 0 to  $t$  giving

$$\hat{\xi}'_n(t)\hat{h}_n(t)^2 - \hat{\xi}'_n(0)\hat{h}_n(0)^2 = \int_0^t \hat{f}_n(t')\hat{h}_n(t')dt'.$$

Now this can be integrated from 0 to  $t$  giving

$$\hat{\xi}'_n(t)\hat{h}_n(t)^2 - \hat{\xi}'_n(0)\hat{h}_n(0)^2 = \int_0^t \hat{f}_n(t')\hat{h}_n(t')dt'.$$

**Question:** we know what  $\hat{h}_n(0)$  from the initial conditions but what is  $\hat{\xi}'_n(0)$  (or  $\hat{\xi}_n(0)$ )?

Now this can be integrated from 0 to  $t$  giving

$$\hat{\xi}'_n(t)\hat{h}_n(t)^2 - \hat{\xi}'_n(0)\hat{h}_n(0)^2 = \int_0^t \hat{f}_n(t')\hat{h}_n(t')dt'.$$

**Question:** we know what  $\hat{h}_n(0)$  from the initial conditions but what is  $\hat{\xi}'_n(0)$  (or  $\hat{\xi}_n(0)$ )?

They have to be 0 since the complete solution was the solution to the homogeneous problem with the given boundaries + the particular inhomogeneous solution with zero boundaries.

Now this can be integrated from 0 to  $t$  giving

$$\hat{\xi}'_n(t)\hat{h}_n(t)^2 - \hat{\xi}'_n(0)\hat{h}_n(0)^2 = \int_0^t \hat{f}_n(t')\hat{h}_n(t')dt'.$$

**Question:** we know what  $\hat{h}_n(0)$  from the initial conditions but what is  $\hat{\xi}'_n(0)$  (or  $\hat{\xi}_n(0)$ )?

They have to be 0 since the complete solution was the solution to the homogeneous problem with the given boundaries + the particular inhomogeneous solution with zero boundaries.

Now we can write the formal solution for the coefficients as

$$\hat{p}_n(t) = \hat{h}_n(t)\hat{\xi}_n(t) = \int_0^t \int_0^{t'} \hat{f}_n(t'') \frac{\hat{h}_n(t)\hat{h}_n(t'')}{\hat{h}_n(t')^2} dt'' dt'$$

Now this can be integrated from 0 to  $t$  giving

$$\hat{\xi}'_n(t)\hat{h}_n(t)^2 - \hat{\xi}'_n(0)\hat{h}_n(0)^2 = \int_0^t \hat{f}_n(t')\hat{h}_n(t')dt'.$$

**Question:** we know what  $\hat{h}_n(0)$  from the initial conditions but what is  $\hat{\xi}'_n(0)$  (or  $\hat{\xi}_n(0)$ )?

They have to be 0 since the complete solution was the solution to the homogeneous problem with the given boundaries + the particular inhomogeneous solution with zero boundaries.

Now we can write the formal solution for the coefficients as

$$\hat{p}_n(t) = \hat{h}_n(t)\hat{\xi}_n(t) = \int_0^t \int_0^{t'} \hat{f}_n(t'') \frac{\hat{h}_n(t)\hat{h}_n(t'')}{\hat{h}_n(t')^2} dt'' dt'$$

and the complete solution is

$$\hat{u}_n(t) = \hat{h}_n(t) + \hat{p}_n(t).$$

## The other way

That all seemed somewhat complicated so let's do something else. For the coefficients of the particular solution we have

$$\frac{\partial^2}{\partial t^2} \hat{p}_n(t) + \lambda_n^2 \hat{p}_n(t) = \hat{f}_n(t).$$

## The other way

That all seemed somewhat complicated so let's do something else. For the coefficients of the particular solution we have

$$\frac{\partial^2}{\partial t^2} \hat{p}_n(t) + \lambda_n^2 \hat{p}_n(t) = \hat{f}_n(t).$$

It turns out that the calculation of the Fourier coefficients can be extended to infinity and the function  $\hat{p}_n$  can be expressed as

$$\hat{p}_n(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \hat{p}_n(\omega) e^{i\omega t} d\omega.$$

(Note that now  $\omega$  is a continuous variable.)

## The other way

That all seemed somewhat complicated so let's do something else. For the coefficients of the particular solution we have

$$\frac{\partial^2}{\partial t^2} \hat{p}_n(t) + \lambda_n^2 \hat{p}_n(t) = \hat{f}_n(t).$$

It turns out that the calculation of the Fourier coefficients can be extended to infinity and the function  $\hat{p}_n$  can be expressed as

$$\hat{p}_n(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \hat{p}_n(\omega) e^{i\omega t} d\omega.$$

(Note that now  $\omega$  is a continuous variable.)

The function  $\hat{p}_n$  can be obtained as

$$\hat{p}_n = \int_{-\infty}^{\infty} \hat{p}_n(t) e^{-i\omega t} dt.$$



## The other way

That all seemed somewhat complicated so let's do something else. For the coefficients of the particular solution we have

$$\frac{\partial^2}{\partial t^2} \hat{p}_n(t) + \lambda_n^2 \hat{p}_n(t) = \hat{f}_n(t).$$

It turns out that the calculation of the Fourier coefficients can be extended to infinity and the function  $\hat{p}_n$  can be expressed as

$$\hat{p}_n(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \hat{p}_n(\omega) e^{i\omega t} d\omega.$$

(Note that now  $\omega$  is a continuous variable.)

The function  $\hat{p}_n$  can be obtained as

$$\hat{p}_n = \int_{-\infty}^{\infty} \hat{p}_n(t) e^{-i\omega t} dt.$$

This is the actual **Fourier transform** and here we have also defined the inverse transform.

We will write the coefficients using the Fourier expression giving the equation

$$\frac{\partial^2}{\partial t^2} \int_{-\infty}^{\infty} \hat{p}_n(\omega) e^{i\omega t} d\omega + \int_{-\infty}^{\infty} \lambda_n^2 \hat{p}_n(\omega) e^{i\omega t} d\omega = \int_{-\infty}^{\infty} \hat{f}_n(\omega) e^{i\omega t} d\omega.$$

We will write the coefficients using the Fourier expression giving the equation

$$\frac{\partial^2}{\partial t^2} \int_{-\infty}^{\infty} \hat{P}_n(\omega) e^{i\omega t} d\omega + \int_{-\infty}^{\infty} \lambda_n^2 \hat{P}_n(\omega) e^{i\omega t} d\omega = \int_{-\infty}^{\infty} \hat{F}_n(\omega) e^{i\omega t} d\omega.$$

Taking the derivative inside the integral gives

$$- \int_{-\infty}^{\infty} \omega^2 \hat{P}_n(\omega) e^{i\omega t} d\omega + \int_{-\infty}^{\infty} \lambda_n^2 \hat{P}_n(\omega) e^{i\omega t} d\omega = \int_{-\infty}^{\infty} \hat{F}_n(\omega) e^{i\omega t} d\omega.$$

We will write the coefficients using the Fourier expression giving the equation

$$\frac{\partial^2}{\partial t^2} \int_{-\infty}^{\infty} \hat{p}_n(\omega) e^{i\omega t} d\omega + \int_{-\infty}^{\infty} \lambda_n^2 \hat{p}_n(\omega) e^{i\omega t} d\omega = \int_{-\infty}^{\infty} \hat{f}_n(\omega) e^{i\omega t} d\omega.$$

Taking the derivative inside the integral gives

$$- \int_{-\infty}^{\infty} \omega^2 \hat{p}_n(\omega) e^{i\omega t} d\omega + \int_{-\infty}^{\infty} \lambda_n^2 \hat{p}_n(\omega) e^{i\omega t} d\omega = \int_{-\infty}^{\infty} \hat{f}_n(\omega) e^{i\omega t} d\omega.$$

It turns out that the functions  $\exp(i\omega t)$  form an orthogonal basis implying that the coefficients have to be the same. This gives

$$-\omega^2 \hat{p}_n(\omega) + \lambda_n^2 \hat{p}_n(\omega) e^{i\omega t} = \hat{f}_n(\omega).$$

Solving for  $\hat{P}_n$  gives

$$\hat{P}_n(\omega) = \frac{\hat{F}_n(\omega)}{\lambda_n^2 - \omega^2}.$$

Solving for  $\hat{P}_n$  gives

$$\hat{P}_n(\omega) = \frac{\hat{F}_n(\omega)}{\lambda_n^2 - \omega^2}.$$

We can use this to solve for  $\hat{p}_n$  resulting in

$$\hat{p}_n(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{\hat{F}_n(\omega)}{\lambda_n^2 - \omega^2} e^{i\omega t} d\omega.$$

Solving for  $\hat{P}_n$  gives

$$\hat{P}_n(\omega) = \frac{\hat{F}_n(\omega)}{\lambda_n^2 - \omega^2}.$$

We can use this to solve for  $\hat{p}_n$  resulting in

$$\hat{p}_n(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{\hat{F}_n(\omega)}{\lambda_n^2 - \omega^2} e^{i\omega t} d\omega.$$

What if  $\hat{F}_n(\lambda_n)$  is not zero? Does the integral diverge?

Solving for  $\hat{P}_n$  gives

$$\hat{P}_n(\omega) = \frac{\hat{F}_n(\omega)}{\lambda_n^2 - \omega^2}.$$

We can use this to solve for  $\hat{p}_n$  resulting in

$$\hat{p}_n(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{\hat{F}_n(\omega)}{\lambda_n^2 - \omega^2} e^{i\omega t} d\omega.$$

What if  $\hat{F}_n(\lambda_n)$  is not zero? Does the integral diverge?

That is something that is called a **resonance** but that will be a story for another day...