

Last time: $f(x) : \mathbb{R}^n \rightarrow \mathbb{R}^m$

The condition number $K_f(x) := \frac{\|Df(x)\| \|x\|}{\|f(x)\|}$

Today: Apply it to $Ax = b$

• Motivation: $A \in \mathbb{R}^{n \times n}$, $b \in \mathbb{R}^n$, A invertible

Solving $Ax = b$ exactly: $f(b) = A^{-1}b$

A backward stable algorithm satisfies $\hat{f}(b) = f(b) = A^{-1}(b + \Delta b)$
with $\|\Delta b\|/\|b\| = O(\epsilon_{\text{mach}})$

To analyse forward error, need to compute K_A

$$K_A(b) = \frac{\|Df\| \|b\|}{\|A^{-1}b\|} = \frac{\|A^{-1}\| \|b\|}{\|A^{-1}b\|}$$

We may want a condition number that is independent of input b ,

$$K(A) := \sup_{b \in \mathbb{R}^n} \frac{\|A^{-1}\| \|b\|}{\|A^{-1}b\|} = \|A^{-1}\| \|A\|$$

\uparrow
 $\|b\| = \|A A^{-1}b\| \leq \|A\| \|A^{-1}b\|$
"=" \uparrow attainable
for some $b \in \mathbb{R}^n$

$K(A)$ is called the condition number of A

• Properties of condition #s

1) $K(A) \geq 1$, $K(A^{-1}) = K(A)$

2) If A is normal with eigenvalues $|\lambda_1| \geq |\lambda_2| \geq \dots \geq |\lambda_n| > 0$

then $K_2(A) := \|A\|_2 \|A^{-1}\|_2 = \frac{|\lambda_1|}{|\lambda_n|}$

For general A , if the matrix norm is submultiplicative,

we have $K(A) \geq \frac{|\lambda_1|}{|\lambda_n|}$

Pf: When A is normal, $A = U^* \Lambda U$,

$$\|A\|_2 = \sqrt{\lambda_{\max}(A^*A)} = \sqrt{\lambda_{\max}(U^* \Lambda^2 U)} = |\lambda_1|$$

Similarly $\|A^{-1}\|_2 = |\lambda_n|^{-1}$

For general A , we need the following relation
between norm and maximum eigenvalue

Lemma: If $\|\cdot\|$ is a submultiplicative matrix norm,

then the spectral radius of A , denote by $\rho(A) = \{|\lambda| : \lambda \text{ is an eigenvalue of } A\}$
satisfies $\rho(A) \leq \|A\|$

Proof of lemma: exercise \square

Using this lemma, we have $\|A\| \geq |\lambda_1|$, $\|A^{-1}\| \geq |\lambda_n|^{-1}$
and the lower bound of A follows \square

3) condition #s of different norms are equivalent.

In particular, $\frac{1}{n} K_2(A) \leq K_1(A) \leq n K_2(A)$

4) 2-condition # measures the inverse of the relative distance
to the nearest singular matrix.

Prop. Let $A \in \mathbb{R}^{n \times n}$, A not singular, then

$$\frac{1}{K_2(A)} = \min_{\substack{\delta A: A + \delta A \\ \text{singular}}} \left\{ \frac{\|\delta A\|_2}{\|A\|_2} \right\}$$

Pf: exercise. \square

We have the following thm that tells you how the solution
is perturbed given an inaccurate right hand side.

Thm 1 Let $Ax = b$ and $r = b - A\hat{x}$
 \uparrow residual

then we have

$$\frac{1}{K(A)} \frac{\|r\|}{\|b\|} \leq \frac{\|\hat{x} - x\|}{\|x\|} \leq K(A) \frac{\|r\|}{\|b\|}$$

Pf: Let $\Delta x = \hat{x} - x$

$$\|\Delta x\| = \|A^{-1}r\| \leq \|A^{-1}\| \|r\|$$

use $\|b\| = \|Ax\| \leq \|A\| \|x\|$ we obtain inequ on the right.

The left inequ can be proved similarly \leftarrow exercise \square

$K(A)$ seems to only quantify the error in output for a perturbation in b , but it turns out that it also quantify the perturbation in A

Thm Let $Ax = b$, $(A + \Delta A)\hat{x} = b + \Delta b$

where $\|\Delta A\| \leq \varepsilon \|A\|$, $\|\Delta b\| \leq \varepsilon \|b\|$

and $\varepsilon K(A) < 1$. then

$$\frac{\|x - \hat{x}\|}{\|x\|} \leq \frac{K(A)}{1 - K(A) \cdot \|\Delta A\| / \|A\|} \left(\frac{\|\Delta A\|}{\|A\|} + \frac{\|\Delta b\|}{\|b\|} \right)$$

$$\approx K(A) \varepsilon \quad \text{for } \varepsilon < \frac{1}{K(A)}$$

form of eqn we get in backward error analysis of many algo for solving $Ax = b$

Pf: $\hat{x} = (A + \Delta A)^{-1}(b + \Delta b)$

$$x = (A + \Delta A)^{-1}(A + \Delta A)x = (A + \Delta A)^{-1}(b + (\Delta A)x)$$

$$\hat{x} - x = (A + \Delta A)^{-1}(\Delta b - (\Delta A)x)$$

$$= [A(I + A^{-1}\Delta A)]^{-1} (\Delta b - (\Delta A)x)$$

$$= (I + A^{-1}\Delta A)^{-1} A^{-1} (\Delta b - (\Delta A)x)$$

$$\|\hat{x} - x\| \leq \|(I + A^{-1}\Delta A)^{-1}\| \|A^{-1}\| (\|\Delta b\| + \|\Delta A\| \cdot \|x\|)$$

$$\text{we use } \|b\| = \|Ax\| \leq \|A\| \|x\|$$

$$\frac{\|\hat{x} - x\|}{\|x\|} \leq \|(I + A^{-1}\Delta A)^{-1}\| \kappa(A) \left(\frac{\|\Delta b\|}{\|b\|} + \frac{\|\Delta A\|}{\|A\|} \right) \quad (*)$$

Lemma: Let $\|\cdot\|$ be a submultiplicative matrix norm.

If $\|B\| < 1$, then $I+B$ is invertible

$$\text{and } \|(I+B)^{-1}\| \leq \frac{1}{1 - \|B\|}$$

Pf: exercise \square

Using this lemma, we have

$$\|(I + A^{-1}\Delta A)^{-1}\| \leq \frac{1}{1 - \|A^{-1}\Delta A\|} \leq \frac{1}{1 - \kappa(A) \|\Delta A\| / \|A\|} \quad (**)$$

The proof is finished after combining (*) (**). \square

ex. Matrix with large condition number (ill-posed problem)

$$\text{Hilbert matrix } H_n = \left(\frac{1}{i+j-1} \right)_{1 \leq i, j \leq n}$$

$$\begin{aligned} \kappa_2(H_n) &= \|H_n\|_2 \|H_n^{-1}\|_2 \\ &\approx \frac{(1+\sqrt{2})^{4n}}{\sqrt{n}} \approx \frac{34^n}{\sqrt{n}} \end{aligned}$$

$$\text{when } n=20, \quad \kappa_2(H_n) \approx 10^{29} \gg 10^{16} = 1/\epsilon_{mach}$$

Review: unitary matrix $Q^*Q = I$, $Q \in \mathbb{C}^{n \times n}$

$$Q = [\vec{q}_1 \dots \vec{q}_n], \quad q_i \in \mathbb{C}^n, \quad i = 1, \dots, n$$

$$Q \text{ is unitary} \iff q_i^* q_j = \delta_{ij}, \quad i, j = 1, \dots, n$$

i.e. $\{\vec{q}_1, \dots, \vec{q}_n\}$ forms an orthonormal basis on \mathbb{C}^n

Some properties: Let $x \in \mathbb{R}^n$, $A \in \mathbb{C}^{n \times n}$

$$\|Qx\|_2 = \|x\|_2$$

$$\|QA\|_2 = \|AQ\|_2 = \|A\|_2$$

$$\|QA\|_F = \|AQ\|_F = \|A\|_F$$

} unitary invariant norm

$$\text{Pf: } \|Qx\|_2^2 = x^* Q^* Q x = x^* x = \|x\|_2^2$$

$$\|QA\|_2^2 = \sqrt{\lambda_{\max}(A^* Q^* Q A)} = \sqrt{\lambda_{\max}(A^* A)} = \|A\|_2^2$$

$$\|QA\|_F^2 = \sqrt{\text{tr}(A^* Q^* Q A)} = \sqrt{\text{tr}(A^* A)} = \|A\|_F^2$$

Other equation can be proved similarly ← exercise

In Linear algebra course, we know when $A \in \mathbb{C}^{n \times n}$ is normal, then A is diagonalizable, i.e., can be factored as

$$A = U \Lambda U^*$$

where $U \in \mathbb{C}^{n \times n}$ is unitary, $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_n)$

• Q: A is not normal? $A \in \mathbb{C}^{m \times n}$?

• Singular value Decomposition

$$A \in \mathbb{C}^{m \times n},$$

$$\text{SVD: } A = U \Sigma V^*$$

$$\text{where } U = \begin{bmatrix} \dot{u}_1 & \dots & \dot{u}_m \end{bmatrix} \in \mathbb{C}^{m \times m}, \quad u_i \text{ left singular vector} \quad \left. \vphantom{\begin{matrix} U \\ V \end{matrix}} \right\} \text{unitary}$$

$$V = \begin{bmatrix} \dot{v}_1 & \dots & \dot{v}_n \end{bmatrix} \in \mathbb{C}^{n \times n}, \quad v_j \text{ right singular vector}$$

$$\Sigma = \begin{bmatrix} \sigma_1 & & \\ & \ddots & \\ & & \sigma_p \end{bmatrix} \in \mathbb{R}^{m \times n}, \quad p = \min\{m, n\} \quad \text{diagonal}$$

$$\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_p \geq 0 \quad \text{singular values}$$

it's not always possible to make $u_i^* v_j = \delta_{ij}$

Existence of SVD:

Since A^*A is Hermitian, positive-definite,

$$\text{we have } A^*A = V \begin{bmatrix} \sigma_1^2 & & \\ & \ddots & \\ & & \sigma_n^2 \end{bmatrix} V^*$$

$$\text{i.e. } A^*A v_i = \sigma_i^2 v_i, \quad i=1, \dots, n, \quad v_i^* v_j = \delta_{ij}$$

$$\text{Since } \text{rank}(A^*A) = \text{rank}(A) =: r$$

$$\text{we know } \sigma_1 \geq \dots \geq \sigma_r > \sigma_{r+1} = \dots = \sigma_n = 0$$

$$\text{Let } u_i = \frac{A v_i}{\sigma_i}, \quad i=1, \dots, r, \quad u_i \in \mathbb{C}^m$$

$$\text{Clearly } u_i^* u_j = \frac{v_i^* A^* A v_j}{\sigma_i \sigma_j} = \delta_{ij}, \quad i, j=1, \dots, r$$

$$\text{and } A v_i = \sigma_i u_i, \quad i=1, \dots, r \quad \text{by definition.}$$

$$\text{Since } \dim(\text{Null}(A)) = n-r, \text{ we know } A v_i = 0, \quad i=r+1, \dots, n$$

We can enlarge u_i to a set of orthonormal basis

$$\{u_1, \dots, u_r, u_{r+1}, \dots, u_m\}, \quad \text{s.t. } u_i^* u_j = \delta_{ij}, \quad i, j=1, \dots, m$$



Facts about SVD:

1) $r := \text{rank}(A) = \# \text{ non-zero sv's } \sigma_i$

2) $\text{range}(A) = \text{span}\{u_1, \dots, u_r\}$

$\text{null}(A) = \text{span}\{v_{r+1}, \dots, v_n\}$

3) $\|A\|_2 = \sigma_1$,

When $A \in \mathbb{C}^{n \times n}$ invertible, then

$$\kappa_2(A) = \sigma_1 / \sigma_n$$

Pf: $\|A\|_2 = \|U \Sigma V^*\|_2 = \|\Sigma\|_2 = \sqrt{\lambda_{\max}(\Sigma^* \Sigma)} = \sigma_1$

Similarly $\|A^{-1}\|_2 = \sigma_n^{-1}$ and hence $\kappa_2(A) = \sigma_1 / \sigma_n$

4) $A = \sum_{i=1}^r \sigma_i u_i v_i^*$,

Best rank k approximation: $A_k := \sum_{i=1}^k \sigma_i u_i v_i^*$ ($k \leq \min(m, n)$)

$$\min_{\text{rank}(B) \leq r} \|A - B\|_F = \|A - A_k\|_F = \sqrt{\sum_{i>k} \sigma_i^2} \quad (1)$$

$$\min_{\text{rank}(B) \leq r} \|A - B\|_2 = \|A - A_k\|_2 = \sigma_{k+1} \quad (2)$$

(Eckart-Young, 1936)

5) Point cloud alignment

Many tasks in computer vision involve the alignment of 3D shapes. Suppose we have a laser scanner that collects two point clouds of the same rigid objects from different views. how can we align these two

point clouds into a single coordinate frame.

Consider two sets of point clouds

$$X = \begin{bmatrix} x_1^T & \dots & x_n^T \end{bmatrix}, \quad Y = \begin{bmatrix} y_1^T & \dots & y_n^T \end{bmatrix} \in \mathbb{R}^{d \times n}$$

We aim to solve the following orthogonal Procrustes problem.

$$\begin{aligned} \min \quad & \|RX - Y\|_F^2 \\ \text{A: } & R^T R = I \\ & R \in \mathbb{R}^{d \times d} \end{aligned}$$

To find R , we note that

$$\begin{aligned} \|RX - Y\|_F^2 &= \text{tr}[(RX - Y)^T (RX - Y)] \\ &= \text{tr}(X^T X - Y^T R X - X^T R^T Y + Y^T Y) \end{aligned}$$

Thus, we want to maximize

$$\text{tr}(Y^T R X + X^T R^T Y) = 2 \text{tr}(Y^T R X) = 2 \text{tr}(R X Y^T)$$

Let $XY^T = U \Sigma V^T$, then

$$\begin{aligned} \text{tr}(R X Y^T) &= \text{tr}(R U \Sigma V^T) \\ &= \text{tr}(\underbrace{V^T R U}_{\tilde{R}} \Sigma) \\ &= \sum_{i=1}^d \tilde{r}_{ii} \sigma_i \end{aligned}$$

Since \tilde{R} is orthogonal, $|\tilde{r}_{ii}| \leq 1$, and hence

$$\text{tr}(R X Y^T) \leq \sum_{i=1}^d \sigma_i$$

"=" iff $\tilde{r}_{ii} = 1$, again since \tilde{R} is orthogonal, $\tilde{R} = I$

hence $R = V U^T$

b) Regularizing ill-posed problems

Consider solving the following problem

$$Ax = b \quad \text{with} \quad A \in \mathbb{R}^{n \times n}, \quad \lambda_i(A) > 0$$

$$\kappa_2(A) = \frac{\lambda_{\max}(A)}{\lambda_{\min}(A)} \gg 1$$

Since the problem is ill-posed, any algorithm that solves the problem is difficult.


How do we decrease the condition number?
(regularize the problem)

we may solve instead

$$(A + \alpha I) \hat{x} = b \quad (\alpha > 0)$$

$$\kappa_2(A) = \frac{\lambda_{\max}(A) + \alpha}{\lambda_{\min}(A) + \alpha} < \kappa_2(A)$$

How do we choose α ?



Since

$$\begin{aligned} x - \hat{x} &= (A + \alpha I)^{-1} (A + \alpha I - A) A^{-1} b \\ &= \alpha (A + \alpha I)^{-1} x \end{aligned}$$

$$\text{we have} \quad \|x - \hat{x}\|_2 \leq \alpha \| (A + \alpha I)^{-1} \|_2 \|x\|_2 = \frac{\alpha}{\lambda_{\min}(A) + \alpha} \|x\|_2$$

$$\text{we hope to balance} \quad \frac{\|x - \hat{x}\|_2}{\|x\|_2} \sim \kappa_2(A + \alpha I) \varepsilon_{\text{mach}}$$

that is $\frac{\alpha}{\lambda_{\min}(A) + \alpha} \sim \frac{\lambda_{\max}(A) + \alpha}{\lambda_{\min}(A) + \alpha} \epsilon_{mach}$

that is $\alpha \sim \lambda_{\max}(A) \epsilon_{mach}$