

## 18.335 Problem Set 2

There are four problems below (Problem 1 - Problem 4). You may complete as many problems as you want, but please submit **only three** problems for grading. **Due March 30, 2025 at 11:59pm.** Late submission will only be accepted (with mild penalty) up to **three days** after the due date. You should submit your problem set **electronically** on the 18.335 Gradescope page. Submit **both** a *scan* of any handwritten solutions (I recommend an app like TinyScanner or similar to create a good-quality black-and-white “thresholded” scan) and **also** a *PDF printout* of the Julia notebook of your computer solutions.

### Problem 0: Pset Honor Code

Include the following statement in your solutions:

*I will not look at 18.335 pset solutions from previous semesters. I may discuss problems with my classmates or others, but I will write up my solutions on my own. <your signature>*

### Problem 1: More on Matrices

Please prove the following facts:

- (a) Let  $A \in \mathbb{C}^{n \times n}$ . Show that

$$|\det(A)|^2 \leq \prod_{i=1}^n \sum_{j=1}^n |a_{ij}|^2.$$

Generate a random matrix and check this inequality on your own computer.

**Hint:** Use a suitable matrix factorization.

- (b) Let  $A \in \mathbb{C}^{n \times n}$ . Let  $A$  be Hermitian with positive diagonal entries, i.e.,  $a_{ii} > 0$ ,  $i = 1, \dots, n$ . Show that if  $A$  is either strictly diagonally dominant or irreducibly diagonally dominant, then  $A$  is positive definite. Generate a matrix satisfies these conditions, compute its eigenvalues, and verify that it is indeed positive definite.

**Hint:** Check the eigenvalues of  $A$ .

- (c) Let  $A \in \mathbb{C}^{n \times n}$  be an Hermitian positive definite matrix. Define  $\|x\|_A = \sqrt{x^* A x}$ . Show that  $\|\cdot\|_A$  is a norm over  $\mathbb{C}^n$ .

**Hint:** Use a suitable matrix factorization.

### Problem 2: Solving a System involving Triangular Matrices

Let  $U_1, U_2 \in \mathbb{R}^{n \times n}$  be upper triangular matrices and  $U_1 U_2 - \lambda I$  be invertible. Design an  $O(n^2)$  time complexity (i.e., an operation count of  $O(n^2)$ ) to solve the system:

$$(U_1 U_2 - \lambda I)x = b$$

Then implement your algorithm on your own computer.

**Hint:** Expand the matrix equation component-wise. What patterns do you see?

### Problem 3: Stationary Iterative Methods

- (a) Let  $A \in \mathbb{C}^{n \times n}$  be an Hermitian positive definite matrix. Split  $A$  as:

$$A = L + D + U,$$

where  $L$  is the strictly lower triangular part of  $A$ ,  $U$  is the strictly upper triangular part of  $A$ , and  $D$  is the diagonal part of  $A$ . Define a symmetric Gauss-Seidel method as follows:

$$\begin{aligned} (D + L)x_{k+\frac{1}{2}} &= -Ux_k + b \\ (D + U)x_{k+1} &= -Lx_{k+\frac{1}{2}} + b. \end{aligned}$$

Rewrite the method in the standard iterative form:  $x_{k+1} = Bx_k + f$ , and show that the spectral radius satisfies  $\rho(B) < 1$ , thereby establishing that the symmetric Gauss-Seidel method converges for all Hermitian positive definite matrices.

- (b) Let  $A \in \mathbb{C}^{n \times n}$  be Hermitian positive definite with eigenvalues  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n > 0$ . Prove that the iterative method

$$x_{k+1} = x_k + \omega(b - Ax_k), \quad k = 0, 1, \dots$$

converges when  $\omega \in (0, 2/\lambda_1)$ , and find the optimal  $\omega^*$  that achieves the fastest convergence rate. Implement the method on your own computer for a randomly generated Hermitian positive definite matrix. Experiment with different values of  $\omega$  and verify that the convergence rate is indeed fastest for  $\omega^*$ .

#### Problem 4: Ill-conditioned problems

Let  $H_n = (h_{ij}) \in \mathbb{R}^{n \times n}$  be the Hilbert matrix, defined as:

$$h_{ij} = \frac{1}{i+j-1}.$$

For this problem, you may want to choose an upper bound of approximately  $n \approx 20$ .

- (a) Define  $x_n = (1, \dots, 1)^\top \in \mathbb{R}^n$  and set  $b_n = H_n x_n$ . Use Gauss elimination (with partial pivoting) and Cholesky factorization to solve  $H_n x_n = b_n$ . Compute the relative error in any norm of your choice for different  $n$ :  $\frac{\|x_n - \hat{x}_n\|}{\|x_n\|}$ , where  $\hat{x}_n$  is the computed solution.
- (b) Compute the condition number  $\kappa_2(H_n)$  and compare it with the theoretical estimate:  $\kappa_2(H_n) \approx \frac{(1+\sqrt{2})^{4n}}{\sqrt{n}}$ . Use different methods to compute  $\kappa_2(H_n)$  and discuss any discrepancies.
- (c) Use the Tikhonov regularization to improve the result in (a), and numerically search for the optimal regularization parameter that minimizes the relative error in the chosen norm.
- (d) (Not for credit) Use stationary iterative solvers to solve the system and compare their performance with the direct methods used above.

#### Feedback (optional)

Please let me know how you're finding the course and the first problem set. What are you hoping to get out of the class? How is the pace of lecture? Please rate the difficulty and volume of the first problem set. You can submit an anonymous survey [here](#).