

Last time: $A \in \mathbb{R}^{n \times n}$, $b \in \mathbb{R}^n$, Solve $Ax = b$


$$\Leftrightarrow \min_{x \in \mathbb{R}^n} f(x), \quad f(x) = \frac{1}{2} x^T A x - b^T x$$

For quadratic $f(x)$, both GD and CG work

For generic $f(x)$, how to efficiently find minimizer?

$$x_* \in \operatorname{argmin}_{x \in \mathbb{R}^n} f(x) \Leftrightarrow \nabla f(x_*) = 0 \leftarrow \text{root finding problem}$$

Nonlinear root finder: Newton method



Idea:

- 1) Guess x_k
- 2) Find root of tangent line at x_k
- 3) Set $x_{k+1} = \text{root}$

equation of tangent line: $l(x) = g(x_k) + g'(x_k)(x - x_k)$

$$\text{root } l(x_{k+1}) = 0 \Leftrightarrow x_{k+1} = \varphi(x_k) := x_k - \frac{g(x_k)}{g'(x_k)}$$

Newton's method typically converges quadratically.

$$e_k := x_k - x_*, \text{ then } |e_{k+1}| \leq M |e_k|^2 = O(|e_0|^{2^k})$$

for smooth g and x_0 sufficiently close to root

If $g'(x_*) = 0$, then convergence is usually linear.

(exercise: if we know x_* is an m -fold root of $g(x)$

$$\text{we can modify } \varphi(x) \text{ so that } x_{k+1} = x_k - m \frac{g(x_k)}{g'(x_k)})$$

Newton's method in $n \geq 1$

Goal: $h: \mathbb{R}^n \rightarrow \mathbb{R}^n$, find $h(x) = 0$

$$h(x+\Delta x) = h(x) + J_h(x) \Delta x + o(\|\Delta x\|)$$

$$\Rightarrow \Delta x \approx J_h^{-1}(x) [h(x+\Delta x) - h(x)]$$

= 0 when

$x+\Delta x$ is root of linear approximation

$$\text{Newton step} \Rightarrow x_{k+1} = x_k + J_h^{-1}(x_k) h(x_k)$$

Note: adapts to non-square systems via pseudoinverse or least-squares solve.


Quasi-Newton Methods

Computing the Jacobian and solving $J_h(x) \Delta x_k = h(x_k)$

at each step is not always computationally tractable,

e.g. when n is very large or computing $h(x)$ is very expensive and J_h is not known analytically.

Idea: approximate $J_k \approx J_h(x_k)$ on the fly, using cheap updates at each iteration



If $x_k \rightarrow x_*$, secant line becomes better and better approximation to tangent line

• Secant line requires no explicit knowledge of derivatives of h

$$\text{Secant line equation: } \underbrace{h(x_{k+1}) - h(x_k)}_{=: \Delta h_k} = J_{k+1} \underbrace{(x_{k+1} - x_k)}_{=: \Delta x_k} \quad (*)$$

for $n > 1$, this does not uniquely determine J_{k+1} , so we need to impose more constraints.

No-change condition: A quasi-Newton update must not alter the curvature estimate in directions where the gradient provides no new information.

Formally, $J_{k+1} q = J_k q$ when $q^T \Delta x_k = 0$ (**)

(*) and (**) uniquely determine J_{k+1} from J_k :

From (*): $(J_{k+1} - J_k) q = u (\Delta x_k)^T q$ for some u , $\forall q$

From (**): $\Delta h_k = (J_k + u (\Delta x_k)^T) \Delta x_k$

$$\Rightarrow u = \frac{\Delta h_k - J_k \Delta x_k}{(\Delta x_k)^T \Delta x_k}$$

So we have

$$J_{k+1} = J_k + \underbrace{(\Delta h_k - J_k \Delta x_k) \frac{\Delta x_k^T}{(\Delta x_k)^T \Delta x_k}}_{\text{rank-1 update to } J_k}$$

equivalently, $J_{k+1} = \arg \min \|J - J_k\|_F$
s.t. (*) holds true

i.e. minimal information update.

Algorithm:

Given $x_0 \in \mathbb{R}^n$, $J_0 \in \mathbb{R}^{n \times n}$, $h_0 = h(x_0) \in \mathbb{R}^n$

for $k = 1, 2, 3, \dots$

$$x_{k+1} = x_k - J_k^{-1} h_k$$

$$J_{k+1} = J_k + (\Delta h_k - J_k \Delta x_k) \frac{\Delta x_k^T}{(\Delta x_k)^T \Delta x_k}$$

end

Since we actually need J_k^{-1} at each step, we can use the Sherman-Morrison formula to update J_{k+1}^{-1} from J_k^{-1}

$$J_{k+1}^{-1} = J_k^{-1} + (\Delta x_k - J_k^{-1} \Delta h_k) \frac{(\Delta x_k)^T J_k^{-1}}{(\Delta x_k)^T J_k^{-1} \Delta h_k}$$

This is called Broyden's first update

Broyden's second update comes from applying secant and no-change conditions directly to $G_{k+1} = J_{k+1}^{-1}$

$$\text{Secant condition: } \Delta x_k = G_{k+1} \Delta h_k \quad (\star)$$

$$\text{No-change: } G_{k+1} q = G_k q \quad \text{when } q^T \Delta h_k = 0 \quad (\star\star)$$

$$\Rightarrow G_{k+1} = G_k + (\Delta x_k - G_k \Delta h_k) \frac{\Delta h_k^T}{(\Delta h_k)^T \Delta h_k}$$

$$\text{or equivalently, } G_{k+1} = \arg \min \|G - G_k\|_F \\ \text{s.t. } (\star) \text{ holds true}$$

BFGS update

Recall in optimization problem $f: \mathbb{R}^n \rightarrow \mathbb{R}$, $\min_{x \in \mathbb{R}^n} f(x)$

- Since $x_* \in \arg \min_{x \in \mathbb{R}^n} f(x)$ necessarily requires $\nabla f(x_*) = 0$

Run Newton on ∇f using Hessian $H = \left(\frac{\partial^2 f}{\partial x_i \partial x_j} \right)_{1 \leq i, j \leq n}$

- Note that H is always symmetric, and at local minima,

H is symmetric positive definite (PSD).

- Notice that Broyden updates are not symmetric. Since we want approximation $H_k \rightarrow H$ when $x_k \rightarrow x^*$, can we adapt Broyden updates to keep H_k PSD?

- Idea: Use symmetric low-rank update to enforce symmetry while satisfying secant condition.

$$\Delta g_k = \nabla f(x_{k+1}) - \nabla f(x_k), \quad \Delta x_k = x_{k+1} - x_k$$

$$\text{Secant equation: } H_{k+1} \Delta x_k = \Delta g_k \quad (*)$$

$$\text{Symm. Rank-2 update: } H_{k+1} = H_k + \alpha u u^T + \beta v v^T \quad (**)$$

with $u = \Delta g_k$ and $v = H_k \Delta x_k$, i.e.

$$\Delta H_k = H_{k+1} - H_k = \underbrace{\alpha (\Delta g_k) (\Delta g_k)^T}_{\text{symm. type 2 update}} + \underbrace{\beta (H_k \Delta x_k) (H_k \Delta x_k)^T}_{\text{symm. type 1 update}}$$

choose α, β to satisfy secant equation $(*)$:

$$\Delta g_k = H_k \Delta x_k + \underbrace{\alpha (\Delta g_k) (\Delta g_k)^T \Delta x_k + \beta (H_k \Delta x_k) (H_k \Delta x_k)^T \Delta x_k}_{H_k \Delta x_k (\Delta x_k)^T \cdot H_k \Delta x_k}$$

$$\text{simply take } \alpha = [(\Delta g_k)^T \Delta x_k]^{-1}, \quad \beta = -[(\Delta x_k)^T H_k \Delta x_k]^{-1}$$

$$\begin{aligned} \text{then } & \beta (H_k \Delta x_k) (H_k \Delta x_k)^T \Delta x_k \\ &= - \frac{H_k \Delta x_k (\Delta x_k)^T H_k \Delta x_k}{(\Delta x_k)^T H_k \Delta x_k} = - H_k \Delta x_k \end{aligned}$$

and $(*)$ holds true

$$\Rightarrow H_{k+1} = H_k + \frac{\Delta g_k (\Delta g_k)^T}{(\Delta g_k)^T \Delta g_k} - \frac{H_k \Delta x_k (\Delta x_k)^T H_k}{(\Delta x_k)^T H_k \Delta x_k}$$

we can use the Sherman-Morrison-Woodbury formula to get a fast update formula directly for $H_k^{-1} \rightarrow H_{k+1}^{-1}$.

One can also show that BFGS update is PSD under appropriate restriction on Δx_k

Storing H_k could be memory intense, instead, store the most recent few $(\Delta g_k, \Delta x_k)$ and use them to compute H_k is call L-BFGS (limited memory BFGS).

Lemma (Sherman-Morrison-Woodbury)

Let $A \in \mathbb{R}^{n \times n}$, A invertible, $u, v \in \mathbb{R}^n$,

then $A + uv^T$ invertible iff $1 + v^T A u \neq 0$

and $(A + uv^T)^{-1} = A^{-1} - \frac{A^{-1} u v^T A^{-1}}{1 + v^T A^{-1} u}$