

LITERATURE REVIEW REPORT: DRUG DEVELOPMENT ANALYSIS

Target Study: Clinical Profile and Market Impact of Lecanemab (Leqembi)

- **Intern Name:** Mitushi Jain
- **Domain:** Pharmaceutical Research & Development
- **Task Module:** Task 1 – Literature Review on Drug Development
- **Submission Date:** June 11, 2026

1. Introduction & Overview

Lecanemab (marketed as Leqembi) is a humanized IgG1 monoclonal antibody directed against aggregated soluble and insoluble forms of amyloid-beta ($\text{A}\beta$). Developed jointly by Eisai and Biogen, Lecanemab received accelerated approval from the U.S. Food and Drug Administration (FDA) in January 2023 and achieved full traditional clinical approval in July 2023.

It is indicated for the treatment of mild cognitive impairment (MCI) or mild dementia stages of Alzheimer's disease. Lecanemab represents a significant milestone in neuropharmacology as one of the first disease-modifying therapies that actively targets the underlying pathophysiology of Alzheimer's disease rather than merely managing symptomatic expressions.

2. The Drug Discovery Process & Target Selection

The discovery pipeline for Lecanemab centered on the **Amyloid Cascade Hypothesis**, which posits that the accumulation of $\text{A}\beta$ plaques in the brain parenchyma is the primary driver of neurodegeneration in Alzheimer's disease.

- **Target Specificity:** Unlike earlier monoclonal antibodies that primarily targeted monomeric forms of amyloid or fully formed mature extracellular plaques, the molecular engineering of Lecanemab was optimized to target **soluble amyloid-beta protofibrils**.
- **Mechanism of Action (MoA):** Protofibrils are intermediate, highly toxic species of aggregated amyloid that precede large plaque structures. By selectively binding to these protofibrils with high affinity, Lecanemab neutralizes their intrinsic neurotoxicity and flags them for microglia-mediated clearance via phagocytosis. This prevents downstream synaptic injury, hyperphosphorylation of tau proteins, and subsequent neuronal cell death.

3. Clinical Trial Phases & Efficacy Analysis

The clinical development and validation of Lecanemab progressed through robust trial evaluations to establish safety margins and primary endpoint efficacy:

Phase I & II Clinical Trials

- **Phase I Evaluation:** Focused on safety, tolerability, and pharmacokinetics across ascending single and multiple doses. It demonstrated predictable dose-dependent plasma concentrations and established an acceptable safety profile.
- **Phase II Trial (Study 201):** A randomized, double-blind, placebo-controlled dose-finding study involving 856 patients. It utilized Bayesian adaptive randomization to determine optimal dosing. The results showed that a bi-weekly dose of 10 mg/kg achieved substantial reduction in brain amyloid burden and slowed clinical decline, establishing the framework for Phase III protocols.

Phase III Pivotal Trial (Clarity AD)

The global, multicenter Phase III **Clarity AD trial** was a randomized, double-blind, placebo-controlled study that enrolled 1,795 patients with early-stage Alzheimer's disease. Patients received either intravenous infusions of Lecanemab (10 mg/kg every two weeks) or a placebo over an 18-month duration.

- **Primary Endpoint Result:** The trial successfully met its primary endpoint, measuring changes on the Clinical Dementia Rating-Sum of Boxes (**CDR-SB**) scale. Lecanemab demonstrated a statistically significant **27% reduction in clinical decline** compared to the placebo over 18 months (p-value < 0.001).
- **Secondary Endpoints:** Significant preservation of cognitive function and daily living capabilities was observed via secondary metrics, including the Alzheimer's Disease Assessment Scale–Cognitive Subscale (ADAS-Cog14) and the Alzheimer's Disease Cooperative Study-Activities of Daily Living Scale for Mild Cognitive Impairment (ADCS-ADL-MCI).
- **Safety & Adverse Events:** The primary safety concern identified during trials was **Amyloid-Related Imaging Abnormalities (ARIA)**, specifically ARIA-E (edema/effusion) and ARIA-H (microhemorrhage). ARIA-E occurred in roughly 12.6% of Lecanemab-treated patients, though the majority were asymptomatic and resolved with temporary treatment suspension, emphasizing the need for regular MRI monitoring during clinical administration.

4. Therapeutic Applications

Lecanemab is restricted to patients in the **early stages** of the Alzheimer's disease spectrum:

- **Mild Cognitive Impairment (MCI):** Patients exhibiting subtle but measurable memory impairment who retain independent functional abilities.
- **Mild Dementia Stage:** Patients with confirmed early-stage cognitive deficits where neurodegeneration has begun impacting complex daily routines.

Administration Protocol: The therapeutic standard requires intravenous (IV) infusion at a dosage of 10 mg/kg administered every two weeks. Initiation of therapy requires positive confirmation of amyloid-beta pathology via Positron Emission Tomography (PET) imaging or cerebrospinal fluid (CSF) biomarkers.

5. Market Impact & Future Projections

The commercial introduction of Lecanemab has significantly transformed the landscape of neurodegenerative healthcare and healthcare economics:

- **Healthcare System Transformation:** The requirement for biomarker verification (PET scans/CSF draws) and regular safety monitoring via MRI has accelerated infrastructure upgrades in memory clinics and diagnostic imaging departments globally.
- **Economic Variables:** Priced at an initial wholesale acquisition cost of approximately \$26,500 annually in the United States, the drug has generated widespread discussion regarding affordability, insurance coverage (such as U.S. Centers for Medicare & Medicaid Services), and equitable global healthcare access.
- **Market Competitive Position:** Lecanemab has established itself as a first-line disease-modifying standard, outperforming previous controversial approvals and paving the way for next-generation subcutaneous delivery forms and combination treatments with anti-tau therapies.

6. Conclusion & References

Lecanemab represents a major clinical advancement in translating the molecular biology of amyloid aggregation into a validated disease-modifying pharmaceutical intervention. By slowing cognitive decline by 27% in early-stage patients, it provides clinically meaningful preservation of independence, marking a turning point in managing Alzheimer's disease.

References

1. Eisai Co., Ltd. & Biogen Inc. (2023). *Leqembi (lecanemab-irmb) Full Prescribing Information*. U.S. Food and Drug Administration.
2. van Dyck, C. H., Swanson, C. J., Leoutsakos, J. M., et al. (2023). Lecanemab in Early Alzheimer's Disease. *New England Journal of Medicine*, 388(1), 9-21.
3. Swanson, C. J., Zhang, Y., Dhadda, S., et al. (2021). A randomized, double-blind phase 2b proof-of-concept clinical trial in early Alzheimer's disease with lecanemab, an anti-A β protofibril antibody. *Alzheimer's Research & Therapy*, 13(1), 80.

