

Dantzig-Wolfe and Fenchel decompositions : hybridation and degeneracy

François Lamothe, Alain Hait, Emmanuel Rachelson, Claudio Contardo, Bernard Gendron

2 juillet 2025




Introduction

$$\begin{array}{ll}\min & cx \\ & x \in \mathbb{Z} \\ \text{s.t.} & Ax \leq a \\ & Bx \leq b\end{array}$$


Introduction

$$\begin{aligned} \min_{x \in \mathbb{Z}} \quad & cx \\ \text{s.t.} \quad & Ax \leq a \\ & Bx \leq b \end{aligned}$$



Introduction


$$\begin{array}{ll}\min & cx \\ \text{s.t.} & Ax \leq a \\ & Bx \leq b\end{array}$$



Linear relaxation

Introduction

$$\begin{array}{ll}\min & cx \\ \text{s.t.} & Ax \leq a \\ & x \in P\end{array}$$




A better relaxation

Introduction

$$\begin{array}{ll}\min & cx \\ \text{s.t.} & Ax \leq a \\ & x \in P\end{array}$$

$$\begin{array}{ll}\max & \pi x \\ \text{s.t.} & x \in P\end{array}$$




Polyhedron known only implicitly : oracle

- Dantzig-Wolfe decomposition
- Fenchel decomposition
- Proposed hybridation
- An illustration of degeneracy

Dantzig Wolfe decomposition

$$\begin{aligned} \min_{x \in \mathbb{R}} \quad & cx \\ \text{s.t.} \quad & Ax \leq a \\ & x \in P \end{aligned}$$



Dantzig Wolfe decomposition


$$\min_{x \in \mathbb{R}} cx$$

$$\text{s.t. } Ax \leq a$$

$$x = \sum \lambda_i x_i$$

$$\sum \lambda_i = 1$$

$$\lambda_i \geq 0$$



Point = convex combination of vertices

Dantzig Wolfe decomposition


$$\min_{x \in \mathbb{R}} cx$$

$$\text{s.t. } Ax \leq a$$

$$x = \sum \lambda_i x_i$$

$$\sum \lambda_i = 1$$

$$\lambda_i \geq 0$$



Start with few vertices

Dantzig Wolfe decomposition


$$\min_{x \in \mathbb{R}} cx$$

$$\text{s.t. } Ax \leq a$$

$$x = \sum \lambda_i x_i$$

$$\sum \lambda_i = 1$$

$$\lambda_i \geq 0$$



Add new vertex \rightarrow better inner approx

Dantzig Wolfe decomposition


$$\min_{x \in \mathbb{R}} cx$$

$$\text{s.t. } Ax \leq a$$

$$x = \sum \lambda_i x_i$$

$$\sum \lambda_i = 1$$

$$\lambda_i \geq 0$$



Dual vars of master pb \rightarrow certify solution
is optimal

Dantzig Wolfe decomposition


$$\min_{x \in \mathbb{R}} cx$$

$$\text{s.t. } Ax \leq a$$

$$x = \sum \lambda_i x_i$$

$$\sum \lambda_i = 1$$

$$\lambda_j \geq 0$$



Dual vars of master pb \rightarrow certify solution is optimal

Dantzig Wolfe decomposition

$$\min_{x \in \mathbb{R}} cx$$

$$\text{s.t. } Ax \leq a$$


$$x = \sum \lambda_i x_i$$

$$\sum \lambda_i = 1$$

$$\lambda_i \geq 0$$

$$\max \pi x$$

$$\text{s.t. } x \in P$$



Dual direction + oracle \rightarrow new vertex

Dantzig Wolfe decomposition

$$\min_{x \in \mathbb{R}} cx$$

$$\text{s.t. } Ax \leq a$$


$$x = \sum \lambda_i x_i$$

$$\sum \lambda_i = 1$$

$$\lambda_i \geq 0$$

$$\max \pi x$$

$$\text{s.t. } x \in P$$




Repeat until optimal

Summary

- Dantzig-Wolfe decomposition
- Fenchel decomposition
- Hybridation
- Degeneracy

Fenchel decomposition




Fenchel decomposition = outer
approximation

Fenchel decomposition

$$\min_{x \in \mathbb{R}} cx$$

$$\text{s.t. } Ax \leq a$$

$$\pi^j x \leq \pi_0^j \quad \forall j \in J$$




Fenchel decomposition = outer approximation

Fenchel decomposition

$$\min_{x \in \mathbb{R}} cx$$

$$\text{s.t. } Ax \leq a$$

$$\pi^j x \leq \pi_0^j \quad \forall j \in J$$



New cuts generated with separation problem

Fenchel decomposition


$$\min_{x \in \mathbb{R}} cx$$

$$\text{s.t. } Ax \leq a$$

$$\pi^j x \leq \pi_0^j \quad \forall j \in J$$

$$\max_{\pi, \pi_0} \pi x^* - \pi_0$$

$$\text{s.t. } \pi x_i \leq \pi_0 \quad \forall i \in I$$



New cuts generated with separation problem

Fenchel decomposition


$$\min_{x \in \mathbb{R}} cx$$

$$\text{s.t. } Ax \leq a$$

$$\pi^j x \leq \pi_0^j \quad \forall j \in J$$

$$\max_{\pi, \pi_0} \pi x^* - \pi_0$$

$$\text{s.t. } \pi x_i \leq \pi_0 \quad \forall i \in I$$



Adding cut \rightarrow tighter outer approx

Fenchel decomposition


$$\min_{x \in \mathbb{R}} cx$$

$$\text{s.t. } Ax \leq a$$

$$\pi^j x \leq \pi_0^j \quad \forall j \in J$$

$$\max_{\pi, \pi_0} \pi x^* - \pi_0$$

$$\text{s.t. } \pi x_i \leq \pi_0 \quad \forall i \in I$$



Repeat until optimal

Fenchel decomposition


$$\min_{x \in \mathbb{R}} cx$$

$$\text{s.t. } Ax \leq a$$

$$\pi^j x \leq \pi_0^j \quad \forall j \in J$$

$$\max_{\pi, \pi_0} \pi x^* - \pi_0$$

$$\text{s.t. } \pi x_i \leq \pi_0 \quad \forall i \in I$$



Repeat until optimal

Normalization

$$\max_{\pi, \pi_0} \pi x^* - \pi_0$$

$$\text{s.t. } \pi x_i \leq \pi_0 \quad \forall i \in I$$

Normalization

$$\max_{\pi, \pi_0} \pi x^* - \pi_0$$

$$\text{s.t. } \pi x_i \leq \pi_0 \quad \forall i \in I$$

$$\max_{\lambda_i} 0$$

$$x^* = \sum \lambda_i x_i$$

$$\sum \lambda_i = 1$$

$$\lambda_i \geq 0$$

Normalization

$$\max_{\pi, \pi_0} \pi x^* - \pi_0$$

$$\text{s.t. } \pi x_i \leq \pi_0 \quad \forall i \in I$$


Model is unbounded

Normalization

$$\max_{\pi, \pi_0} \pi x^* - \pi_0$$

$$\text{s.t. } \pi x_i \leq \pi_0 \quad \forall i \in I$$

$$\|\pi\|_2 \leq 1$$




Normalisation with L2 norm

Normalization

$$\max_{\pi, \pi_0} \pi x^* - \pi_0$$

$$\text{s.t. } \pi x_i \leq \pi_0 \quad \forall i \in I$$

$$\|\pi\|_2 \leq 1$$




No guarantee to get a facet

Normalization

$$\max_{\pi, \pi_0} \pi x^* - \pi_0$$

$$\text{s.t. } \pi x_i \leq \pi_0 \quad \forall i \in I$$

$$|\pi(\hat{x} - x)| \leq 1$$




Directional normalisation : always a facet

Summary


- Dantzig-Wolfe decomposition
- Fenchel decomposition
- Hybridation
- Degeneracy

Hybridation




Both inner and outer approx in one algo

Hybridation




Cut outer solution : directional normalization toward inner sol
Obtain a cut + supporting vertices

Hybridation




Update both approximations

Hybridation



Repeat until optimal




Unsplittable flow problem :
root node

Hybridation

Best setup for our method

- Many oracle calls \rightarrow Fast oracle required
example : knapsack




Unsplittable flow problem :
root node

Hybridation

Best setup for our method

- Many oracle calls \rightarrow Fast oracle required
example : knapsack
- Symetries/several optimal solution : Fenchel is slow
Our method suffers less from symetries




Unsplittable flow problem :
root node

Hybridation

Best setup for our method

- Many oracle calls \rightarrow Fast oracle required
example : knapsack
- Symetries/several optimal solution : Fenchel is slow
Our method suffers less from symetries
- Degeneracy :
Dantzig-Wolfe is slow
Fenchel methods don't use duals




Unsplittable flow problem :
root node


Summary

- Dantzig-Wolfe decomposition
- Fenchel decomposition
- Hybridation
- Degeneracy

Degeneracy




Degeneracy




Dual problem = find a bound

Degeneracy




Dual bound decomposes in two parts

Degeneracy




Dual bound decomposes in two parts

Degeneracy




What if there are several decompositions?

Degeneracy




What if there are several decompositions?

Degeneracy




What if there are several decompositions ?

Degeneracy




What if there are several decompositions ?

Degeneracy



What if there are several decompositions ?

Degeneracy



Solving degeneracy ?

Recap :

- Inner approximation : Dantzig-Wolfe
- Outer approximation : Fenchel
- Hybridation can be good ! Use inner and outer approx
- Degeneracy is Dantzig-Wolfe bane
- How to prevent degeneracy in Dantzig-Wolfe ?



Lamothe et al. On the integration of Dantzig-Wolfe and Fenchel decompositions via directional normalizations.

<https://hal.science/hal-04875156v1>

Thank you for your attention !



Lamothe et al. On the integration of Dantzig-Wolfe and Fenchel decompositions via directional normalizations.

<https://hal.science/hal-04875156v1>